

C-6200

Base Reg. (4042)

U.S. LABEL-(S)

Reason to Issue: To show "RESTRICTED USE PESTICIDE" classification; to update label.

Date of Draft 9/13/79 (S)
Supersedes Draft Dated: 10/10/72

EPA Reg. No. 3125-182

RESTRICTED USE PESTICIDE

For retail sale to, and use only by, Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

(R)

GOPHACIDE

0.1% Bait

RODENTICIDE

ACCEPTED
APR 16 1981
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 3125-182

FOR CONTROL OF POCKET GOPHERS

NET CONTENTS POUNDS

WARNING

KEEP OUT OF REACH OF CHILDREN

(See rear panel for warning statements)

ACTIVE INGREDIENT:

O,O-Bis(4-chlorophenyl) (1-iminoethyl) phosphoramidothioate 0.1%

INERT INGREDIENTS: 99.9%
100.0%

U.S. Patent No. 3,281,321
Canadian Patent No. 734,492

EPA Reg. No. 3125-182

STOP - READ THE LABEL BEFORE USE

MOBAY CHEMICAL CORPORATION
AGRICULTURAL CHEMICALS DIVISION
Box 4913, Kansas City, Mo. 64120

Note: Underlined letters in chemical nomenclature should be italicized when printed, not underlined.

41

GOPHACIDE 0.1% Bait

DIRECTIONS FOR USE

IMPORTANT: Read these entire Directions and Conditions of Sale before using GOPHACIDE 0.1% Bait.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE THE DIRECTIONS FOR CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED FOR A RANGE OF WEATHER CONDITIONS SIMILAR TO THOSE WEATHER CONDITIONS THAT ARE ORDINARY AND CUSTOMARY IN THE GEOGRAPHIC AREA WHERE THE PRODUCT IS USED. INSUFFICIENT CONTROL OF PESTS MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL WEATHER, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO THE CROP TO WHICH THE PRODUCT IS APPLIED, OTHER CROPS, ANIMALS, MAN, OR THE ENVIRONMENT. MOBAY AND THE SELLER OFFER, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL WEATHER, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF MOBAY AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

GOPHACIDE 0.1% Bait is effective for the control of the mountain (Thomomys talpoides) and plains (Geomys bursarius) pocket gophers. Bait preparations are readily accepted and death usually occurs within 24 to 48 hours after the bait is eaten.

How to Apply:

Place in artificial burrows made with suitable burrow building equipment. Use 1-1/2 lbs. of GOPHACIDE 0.1% Bait per acre in burrows spaced at 20-foot intervals or 1 lb. per acre in burrows spaced at 30-foot intervals. Treatment may also be made to natural burrows at the rate of 1-1/2 lbs. of GOPHACIDE 0.1% Bait per acre of area to be treated.

RESTRICTIONS

For use only by personnel experienced in rodent control. Consult your local, county, or state rodent control experts for advice on how to best use this product safely and efficiently. Apply only in burrows. Do not use above ground surface. Control of other rodent species is not recommended, since evaluations have not yet been completed.

FISH AND WILDLIFE CAUTION

This product is toxic to fish and wildlife. Keep out of any body of water. Treated bait exposed on soil surface will be hazardous to birds and other wildlife. Apply this product only as specified on this label.

WARNING

May be harmful if swallowed or absorbed through the skin. Do not get in eyes or on skin. This material is cumulative. Repeated contact with the skin may increase hazard of handling. Wear cotton or natural rubber gloves while handling. If cotton gloves are used, they must be laundered or discarded after each day's use. Always wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse.

Do not store or use in an area where there is possibility of children or animals gaining access to the material or to the carcasses of animals poisoned by the bait.

GOPHACIDE 0.1% Bait

WARNING (Cont.)

If poisoning occurs, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression. If swallowed, vomiting should be induced. Administer milk or water freely and induce vomiting by giving one dose (1/2 oz. or 15 ml.) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Never give anything by mouth to an unconscious person. If eyes are contaminated, wash with flowing water for at least 15 minutes.

ANTIDOTE: Although atropine may be less effective as an antidote with this than with some other organic phosphorus compounds, it should be used in large therapeutic doses in cases of acute poisoning. Repeat as necessary to the point of tolerance.

CONTAINER DISPOSAL

DO NOT REUSE THIS CONTAINER - Completely empty the contents and bury the unused chemical at least 18 inches deep in an isolated location away from water supplies.

Burn the empty container completely. Exercise extreme caution and stay well away from the smoke.

Do not use the container in any connection with feed, food, or drinking water.

GOPHACIDE is a Reg. TM of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen.

43