

193-6468.YLD

U.S. LABEL


Base Pre-Reg. (6468)

Reason to Issue: To propose use on Christmas trees, Southern Pine Orchards and Ornamentals.

Date of Draft: 6/20/88 (Pre-Reg) (B)

EPA Reg. No. 3125-193

GUTHION®
50% WETTABLE POWDER
CROP INSECTICIDE




ACTIVE INGREDIENT:
O,O-Dimethyl S-[(4-oxo-1,2,3-benzotriazin-3(4H)-yl)methyl] phosphorodithioate 50%

AMENDMENT

To the Previously Registered Labeling

Add the Following:



RECOMMENDED APPLICATIONS

CROP	INSECT	Pounds GUTHION 50% W.P.	REMARKS
ORNAMENTALS			
Southern Pine Orchards	Coneworm Seedworm	See Remarks	Use 4.5 pounds per 100 gallons of water (0.2% dilution) as a full coverage spray*. Use 2.25 pounds per 10 gallons of water (1% dilution) for low volume sprayers. Apply first application within 30 days following conelet closure, followed by 3 to 5 applications (at least 30 days apart). Apply approximately 5 to 10 gallons of the 0.2% dilution with high volume sprayers or approximately 1 to 2 gallons of the 1% dilution with low volume sprayers per tree. Thorough coverage of cones is necessary for maximum control.

*This concentration is calculated for conventional hydraulic-type sprayers. When lower volumes of spray are applied per acre with concentrate sprayers, increase the concentration of GUTHION 50% WP in the spray mixture in order to apply amount of GUTHION 50% WP per acre equivalent to a full coverage spray. Where conditions dictate an air application, apply dosage per acre equivalent to a full coverage ground spray in not less than one gallon of water per acre.

