


U.S. LABEL

Reason to Issue: To revise walnut use pattern and add split application.

Date of Pre-Reg Draft: 7/27/92 (C)
Supercedes Draft Dated: 10/24/91

EPA Reg. No. 3125-193

GUTHION
50% WETTABLE POWDER
CROP INSECTICIDE



ACTIVE INGREDIENT:
O,O-Dimethyl S-[(4-oxo-1,2,3-benzotriazin-3(4H)-yl)methyl] phosphorodithioate 50%

INERT INGREDIENTS: 50%
100%

EPA Reg. No. 3125-193

AMENDMENT

To Previously Registered Labeling

Revise as Follows:

RECOMMENDED APPLICATIONS

CROPS	INSECTS	POUNDS GUTHION 50% WP	REMARKS
<u>NUTS</u>			
Walnuts	Codling moth Filbertworm Navel orangeworm Walnut husk fly Red-humped caterpillar	3 to 4	Apply specified dosage per acre as a full coverage spray by air or ground. A total of 3 applications may be made per crop season regardless of dosage or formulation used. Allow at least 14 days between applications except in the case of split application described below. Allow at least 21 days between last application and harvest. Do not graze livestock in treated groves for 21 days after treatment.
		2 + 2	<u>Split Application:</u> On large trees, where coverage is difficult to obtain, an aerial application followed by a ground application may be made. In order to achieve control at this rate, the two applications must be made within 10 days. Allow 21 days from the last application until harvest.

BEST AVAILABLE COPY