

PM-19

193-8519

Reg # 3125-193


U.S. LABEL

Base Pre-Reg. (8519)

Reason to Issue: To revise almond use pattern by deleting dormant spray and adding 25 ft buffer.

Date of Pre-Reg Draft: 05/03/95 (C)

# GUTHION 50% WETTABLE POWDER CROP INSECTICIDE



**ACTIVE INGREDIENT:**

O,O-Dimethyl S-[(4-oxo-1,2,3-benzotriazin-3(4H)-yl)methyl] phosphorodithioate ..... 50%

**INERT INGREDIENTS:** ..... 50%

100%

EPA Reg. No. 3125-193

## AMENDMENT To Previously Registered Labeling

Revise as Follows:

RECOMMENDED APPLICATIONS			
CROP	INSECT	POUNDS GUTHION 50% WP	REMARKS
NUTS  Almonds	Peach twig borer Navel orangeworm	3 to 4	Apply specified dosage per acre as an aerial, ground concentrate or dilute application. Use at least 20 gallons per acre by aircraft and up to 400 gallons per acre by ground equipment. Applications are limited to <del>one dormant spray and/or</del> two post-bloom sprays per season.  Alternate Row Spraying: Alternate rows may be sprayed. However, due to the presence of untreated areas, satisfactory control may not be obtained. For reduction of pest numbers to occur, all rows must be treated within 10 days or less.
		2 + 2	Split Application: On large trees where coverage is difficult to obtain, an aerial application followed by a ground application may be made. In order to achieve control at this rate, the two applications must be made within 10 days.
<p>All Applications: Allow <del>120 days between the dormant application and the first post bloom spray.</del> Allow at least 28 days between post-bloom sprays at 3 to 4 lbs per acre or the first spray of a split application. The last application may be made up to 28 days before harvest. Do not graze livestock in treated groves for 21 days after treatment. Applications may not be made within 25 feet of an aquatic site (lakes, reservoirs, rivers, permanent streams, marshes, estuaries, natural ponds, or ponds used for commercial fishing).</p>			

# GUTHION 50% WP