


U.S. LABEL-(S)

Reason to Issue: To limit the amount of product that may be applied per 1000 square feet per year.

Date of Draft: 4/20/89 (Pre-Reg)(S)

EPA Reg. No. 3125-216



DYRENE[®]
LAWN AND GARDEN
FUNGICIDE

ACTIVE INGREDIENT:
4,6-Dichloro-N-(2-chlorophenyl)
-1,3,5-triazin-2-amine 50%

AMENDMENT

To Previously Registered Labeling

Revise as Follows:

RECOMMENDED APPLICATIONS

DISEASE	OUNCES PER 1000 sq. ft.*	REMARKS
Leafspot Melting-out Leaf blight	4 to 8 ozs.	Begin applications in the spring before disease symptoms appear and repeat at <u>10-to-14-day intervals</u> . <u>During extended</u>
Root, Crown, and Rhizome Rots (Helminthosporium spp.) (Drechslera spp.) Dollar Spot Brown Patch Copper Spot Rusts		periods of wet weather, applications every 7 days may be required. Continue treatments throughout the the spring and summer months. Use the higher rate in areas with a history of severe infestations or when disease symptoms are visible. Use the low rate on a protective schedule prior to any symptoms visible on the turfgrass.
Gray Snow Mold (Typhula incarnata)	4 ozs	Apply DYRENE Lawn and Garden fungicide at the specified rate after the first killing frost in the fall. A second application should be made when the turf is free of snow in mid-winter. Apply a third application after final melting of snow in late winter or early spring.

*NOTE: Apply the specified amount of DYRENE Lawn and Garden fungicide in 2 to 4 gallons of water per 1,000 sq. ft. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. Do not apply more than one (1) pound (16 ounces) of DYRENE Lawn and Garden Fungicide per 1000 sq. ft. (43.5 pounds) per year.

ALL COPY EXCEPT THAT WHICH IS UNDERLINED HAS BEEN PREVIOUSLY REGISTERED.

BEST AVAILABLE COPY