

NET WEIGHT 5 POUNDS



DYLOX

5% BAIT INSECTICIDE

EPA Reg. No. 3125-217

FOR EFFECTIVE CONTROL OF SURFACE FEEDING CUTWORMS,
AND SIMILAR INSECTS INFESTING MANY CROPS

ACTIVE INGREDIENT
Chlorpyrifos (2,2-dimethyl-
3-hydroxyethyl-
phosphorothioate) 95%
INERT INGREDIENTS 5%
EPA Reg. No. 3125-217
Federal Patent No. 3,129,379

CAUTION

KEEP OUT OF REACH
OF CHILDREN

See Side Panel for Caution Statements

Chemagro

1000 North 17th Street
P.O. Box 1000
Wilmington, Delaware 19801

PLANTING AND APPLICATION

CROP	Pounds DYLOX 5% Bait	REMARKS
FIELD CROPS		
<u>Alfalfa</u>	20	Do not apply more than 4 times in 14 days
<u>Barley</u> <u>Flax</u> <u>Oats</u> <u>Wheat</u>	20	Do not apply within 20 days of treated fields
<u>Corn</u>	20	Apply once per season high. Do not use corn within 40 days of treat-
<u>Cotton</u>	30	Do not apply within 70 days of livestock in treat- ed areas after application.
<u>Peanuts</u>	20	Applications may be made or broadcast by air or ground applications may be made additional applications digging and harvest.
<u>Safflower</u>	30	Do not apply after plan-
<u>Sugarbeets</u>	30	Do not apply within 28 days of harvest if tops are not to be fed to livestock
<u>Tobacco</u>	20	Do not apply after plan-
SEED FIELD CROPS		
<u>Soybeans</u>	30	Do not pasture or use for food or forage
VEGETABLES		
<u>Beans (Dry & Snap)</u>	20	Do not apply within 14 days of treated vines for use as food
<u>Beans (Green)</u>	30	Do not apply more than once in 14 days of harvest if used for food
<u>Cucumbers (Southern peas, black-eyed peas, crowder peas)</u>	20	Do not apply more than once in 14 days of harvest if used for food

...the use of this product is restricted to the control of surface feeding cutworms and similar insects infesting many crops. The user must understand that the Buyer assumes all such risks.

DYLOX 5% Bait gives excellent control of surface feeding cutworms, cutworms, armyworms and crickets infesting many crops.

Applications may be made in a band over planted row or around the stems of individual plants. If such methods are not practical, apply the bait as a broadcast treatment over the infested area. As an alternate treatment DYLOX 5% Bait may be applied to the soil in a band approximately 6 inches wide and 1/8 to 1/4 inch deep around the field to be protected from worms moving into the crop.

If the field has a history of previous infestations or if cutworms are suspected to be present, apply DYLOX 5% Bait immediately after planting or transplanting.

RESTRICTIONS

PROTECT WILDLIFE!

This product is toxic to birds and other wildlife and must be applied in strict accordance with label directions. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. In cleaning of equipment or disposal of wastes, do not contaminate water.


Do not use on other crops used for food or forage. Use only according to label directions. Applications at rates above those shown may result in illegal crop residues. Do not treat food crops grown in the greenhouse.

CONTAINER AND WASTE DISPOSAL

Rinse equipment and containers and dispose of wastes by burning in non-crop land away from water systems. Dispose of empty container by burying with wastes or by burning. Keep out of smoke.

CAUTION

May be harmful if swallowed. Avoid contact with skin. Avoid contact with skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse. Do not eat, drink, or use tobacco. Keep out of reach of children. If you experience a slight or severe medical ailment, discontinue use. Wash thoroughly with soap and water. To Physician: Atropine sulfate and 1 mg of PAM is dose and state and may be administered as needed with atropine.



RECOMMENDED APPLICATIONS

CROP	Pounds DYLOX 5% Bait	REMARKS
FIELD CROPS		
<u>Alfalfa</u>	20	Do not apply more than one application per cutting nor within 14 days of cutting or grazing
<u>Barley</u> <u>Flax</u> <u>Oats</u> <u>Wheat</u>	20	Do not apply within 21 days of harvest. Do not graze treated fields
<u>Corn</u>	20	Apply once per season when plants are 3-12 inches high. Do not use corn for food, feed, or ensilage within 40 days of treatment
<u>Cotton</u>	30	Do not apply within 7 days of picking. Do not graze livestock in treated fields within 14 days after application. Applications may be made in a band over the row or broadcast by air or ground equipment. Three applications may be made before digging. One additional application may be made between digging and harvest.
<u>Peanuts</u>	20	
<u>Safflower</u>	30	Do not apply after plants begin to bloom
<u>Sugarbeets</u>	30	Do not apply within 28 days of harvest if tops are to be fed to livestock nor within 14 days of harvest if tops are not to be fed
<u>Tobacco</u>	20	Do not apply after plants are 2 feet tall
SEED FIELD CROPS		
Soybeans	30	Do not pasture or use treated crop for feed, food or forage
VEGETABLES		
<u>Beans (Dry & Snap)</u>	30	Do not apply within 14 days of harvest. Do not feed treated vines to livestock or swine
<u>Beans (Lima)</u>	30	Do not apply more than two applications per harvest for food, feed or forage within 30 days of treatment
<u>Cowpeas (Southern peas, black-eyed peas, crowder peas)</u>	30	Do not apply more than two applications per harvest for food, feed or forage within 14 days of harvest. Do not feed treated vines to livestock or swine

RECOMMENDED APPLICATIONS

CROP	Pounds DYLOX 5% Bait	REMARKS
VEGETABLES (Cont'd)		
<u>Brussels Sprouts</u>		
<u>Cabbage</u>	20	Do not apply within 21 days of harvest
<u>Cauliflower</u>		
<u>Collards</u> <u>Lettuce</u>	20	Do not apply within 28 days of harvest
<u>Peppers</u>	20	Do not apply within 21 days of harvest
<u>Pumpkins</u>	20	Do not apply more than once per season nor within 14 days of harvest
<u>Table Beets</u>	30	Do not apply within 28 days of harvest. Do not eat tops
<u>Tomatoes</u>	20	Do not apply within 28 days of harvest

WARRANTY AND LIMITATION OF DAMAGES

Chemagro warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the risks referred to therein. CHEMAGRO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY, and no agent of Chemagro is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

DYLOX is a Reg. TM of Farbenfabriken Bayer AG

ACCEPTED

MAR 21 1972

DYLOX
INSECTICIDE

KILLING CUTWORMS
IN MANY CROPS