

YLD26

U.S. LABEL

Reason to Issue: To propose use on  
non-bearing strawberry nursery stock.

Date of Draft: 8/20/85 (Pre-Reg) (H)

EPA Reg. No. 3125-236-AA

---

® NEMACUR  
15% Granular

SYSTEMIC INSECTICIDE NEMATOCIDE

ACTIVE INGREDIENT:

Ethyl 3-methy-4-(methylthio)phenyl  
(1-methylethyl)phosphoramidate ..... 15%


INERT INGREDIENTS: ..... 85%  
100%

AMENDMENT TO  
Previously Registered Labeling

Add the following:

DIRECTIONS FOR USE


CROP	PEST	DOSAGE NEMACUR 15% G BAND: Ozs/1000 ft. row	REMARKS
Non-bearing Strawberry nursery stock	Nematodes	17 ozs for any row spacing (or 13.8 pounds per acre on 40 inch rows)	BAND: Apply specified dosage in ozs. per 1000 ft. of row in a 12 inch band over the row immediately after transplanting and incorporate to a depth of 1-2 inches with cultivation equipment. Apply a second application in a 12 inch band over the row 8 weeks after transplanting. Incorporate by cultivation or by sprinkler irrigation (1 in. of water) immediately following application. Do not apply more than two applications to non-bearing nursery stock. Use only on strawberry nursery stock that will not be harvested for fruit within 20 months after application.



to include tight heading varieties of  
Chinese cabbage.  
EPA Reg. No. 3125-236-AA

®  
**NEMACUR**  
15% Granular  
**SYSTEMIC INSECTICIDE - NEMATOCIDE**

Only for agricultural use, or for sale to, use, and storage  
by commercial applicators, and other service persons



ACTIVE INGREDIENT:  
Ethyl 3-methyl-4-(methylthio)phenyl  
(1-methylethyl)phosphoramidate ..... 15%

INERT INGREDIENTS: ..... 85%  
100%

AMENDMENT TO  
Previously Registered Labeling.

Revise as Follows:

**RECOMMENDED APPLICATIONS**

CROP	PEST	DOSAGE NEMACUR 15% GRANULAR		REMARKS
		BAND: OZS./ 1000 Ft. of Row	BROADCAST: LBS./ACRE	
<b>VEGETABLES</b>				
Cabbage (includes tight heading varieties of Chinese Cabbage)  (direct seeded and trans- planted)	Nematodes	11 to 22 ozs. for any row spacing (or 10 to 20 lbs. per acre on 36 inch rows)	26.7 to 40 pounds	<b>FOR BAND OR BROADCAST APPL CATION</b>  <b>BAND:</b> Apply specified dos in ozs. per 1000 ft. of ro in front of the planter sh as a 12 to 15 inch band. Incorporate the granules in the soil to a depth of 2 t inches. Use the high rate fields with high populatio of nematodes or in fields having a history of seriou nematode damage. WHEN BAN APPLICATIONS ARE USED ON NARROW ROW CROPS, DO NOT U BAND WIDTHS THAT WILL ALLO TREATED AREAS TO OVERLAP.  <b>BROADCAST:</b> Distribute the granules uniformly over th entire area to be treated immediately incorporate to depth of 2 to 6 inches by disking or tilling. Use t high rate in fields with h populations of nematodes in fields having a history serious nematode damage.
	Nematodes	11 to 22 ozs. for any row spacing (or 10 to 20 lbs. per acre on 36 inch rows)	26.7 to 40 pounds	

**BEST AVAILABLE COPY**

YLD26

U.S. LABEL

Reason to Issue: To propose use on  
non-bearing strawberry nursery stock.

Date of Draft: 8/20/85 (Pre-Reg) (H)

EPA Reg. No. 3125-236-AA

® NEMACUR  
15% Granular


## SYSTEMIC INSECTICIDE NEMATICIDE

## ACTIVE INGREDIENT:

Ethyl 3-methy-4-(methylthio)phenyl  
(1-methylethyl)phosphoramidate ..... 15%INERT INGREDIENTS: ..... 85%  
100%AMENDMENT TO  
Previously Registered LabelingAdd the following:

## DIRECTIONS FOR USE

CROP	PEST	DOSAGE NEMACUR 15% G BAND: Ozs/1000 ft. row	REMARKS
Non-bearing Strawberry nursery stock	Nematodes	17 ozs for any row spacing (or 13.8 pounds per acre on 40 inch rows)	BAND: Apply specified dosage in ozs. per 1000 ft. of row in a 12 inch band over the row immediately after transplanting and incorporate to a depth of 1-2 inches with cultivation equipment. Apply a second application in a 12 inch band over the row 8 weeks after transplanting. Incorporate by cultivation or by sprinkler irrigation (1 in. of water) immediately following application. Do not apply more than two applications to non-bearing nursery stock. Use only on strawberry nursery stock that will not be harvested for fruit within 20 months after application.



BEST AVAILABLE COPY