

PM 22

3125-236

5-12-87

10/1

YLD10

Base Pre-Reg (6022)

U.S. LABEL

Reason To Issue: To reduce pre-harvest  
interval to 30 days for all citrus fruits.


Date of Draft: 10/28/86 (Pre-Reg) (H)

EPA Reg. No. 3125-236-AA

® NEMACUR

15% Granular

SYSTEMIC INSECTICIDE NEMATICIDE



ACTIVE INGREDIENT:

Ethyl 3-methyl-4-(methylthio)phenyl

(1-methylethyl)phosphoramidate ..... 15%

AMENDMENT TO PREVIOUSLY REGISTERED LABELING

REVISE TO READ AS FOLLOWS:

RECOMMENDED APPLICATIONS

CROP	PEST	DOSAGE NEMACUR 15% G	REMARKS
FRUIT			
Citrus	Nematodes	67 to 133 pounds	<p><b>BROADCAST:</b> Apply specified dosage per acre as a broadcast application. Immediately after application, incorporate the granules into the soil by light raking or tilling.</p> <p><b>BAND:</b> Apply proportionally less product per acre based on band width. (Example: to treat trees planted on 20 foot centers with a 10 foot band centered on the row, apply one-half of the broadcast rate). Center the treated band on the tree row using a minimum band width of "to the drip line". Incorporate immediately after application.</p> <p>Under dry conditions, follow with 1/2 to 1 inch of overhead or flood irrigation. Do not apply more than once per season nor within 30 days of harvest. Do not graze livestock on treated areas. On non-bearing citrus make initial applications when trees have recovered from transplanting shock and have initiated new growth, provided that one or more irrigations have been applied and soil has settled around the roots. Phytotoxicity may occur if applications are made before soil has settled and plants are growing.</p>