

## U.S. LABEL

**Reason to Issue:** To add pest to citrus. **6/19/89 Draft:** Revise to read "Except Florida" and "Florida Only." **12/5/89 Draft:** Remove imported fire ant.

**Date of Draft:** 12/5/89 (Pre-Reg (T))  
**Supersedes Pre. Reg Draft Dated:** 6/19/89.

EPA Reg. No. 3125-236

## RESTRICTED USE PESTICIDE

For Sale to and Use Only by a Certified Applicator for Uses Authorized By His Certification, or by Persons Under His Direct Supervision.

**NEMACUR®**  
**15% Granular**  
**SYSTEMIC INSECTICIDE - NEMATOCIDE**

FOR EFFECTIVE CONTROL OF  
 NEMATODES IN CERTAIN FIELD,  
 FRUIT AND VEGETABLE CROPS

## ACTIVE INGREDIENT:

Ethyl 3-methyl-4-(methylthio)phenyl  
 (1-methylethyl)phosphoramidate ..... 15%

INERT INGREDIENTS: ..... 85%  
 100%

AMENDMENT TO PREVIOUSLY REGISTERED LABELING  
 RECOMMENDED APPLICATIONS

CROP	PEST	DOSAGE OF NEMACUR 15% G	REMARKS
FRUIT			
citrus (Except Florida) (In California do not apply to Kumquat, Tangelo or Citrus hybrids)	Nematodes  Citrus root weevil complex including Fuller rose beetle (Suppression)	33 1/2 to 67 pounds	<p>Apply specified dosage per acre in a band in up to 2 applications per season, not to exceed 67 pounds of NEMACUR 15% Granular per acre per season. Apply to the soil with ground equipment and incorporate immediately using mechanical incorporation or irrigation. Center the treated band on the tree row using a band width of 50% of the row spacing and covering the feeder root system of the plant.</p> <p>Under dry conditions, follow with overhead or flood irrigation. Do not apply more than 10 pounds active ingredient (67 pounds of NEMACUR 15% Granular) per acre per season. Do not apply within 30 days of harvest.; Do not graze livestock on treated areas. On non-bearing citrus, make initial applications when trees have recovered from transplanting shock and have initiated new growth, provided that one or more irrigations have been applied and soil has settled around the roots. Phytotoxicity may occur if applications are made before soil has settled and plants are growing.</p>

ACCEPTED

APR 24 1990

Under the Federal Insecticide, Fungicide, and  
 Rodenticide Act, as amended, the  
 pesticide registered under  
 EPA Reg. No. 3125-236

**NEMACUR 15% G**

**RECOMMENDED APPLICATIONS**

CROP FRUIT	PEST	DOSAGE OF NEMACUR 15% G	REMARKS
Citrus (Florida only)	Nematodes  Citrus root weevil complex including Fuller rose beetle (Suppression)	33 1/2 to 67 pounds	<p>Apply the specified dosage per acre in a band in up to 2 applications per season, not to exceed 67 pounds of NEMACUR 15% Granular per acre per season. Apply to the soil with ground equipment and incorporate immediately using mechanical incorporation or irrigation. Center the treated band on the tree row using a band width of 50% of the row spacing and covering the feeder root system of the plant.</p> <p>Under dry conditions, follow with overhead or flood irrigation. Do not apply more than 10 pounds active ingredient (67 pounds or NEMACUR 15% Granular) per acre per season. Do not apply within 30 days of harvest. Do not graze livestock on treated areas. On non-bearing citrus, make initial applications when trees have recovered from transplanting shock and have initiated new growth, provided that one or more irrigations have been applied and soil has settled around the roots. Phytotoxicity may occur if applications are made before soil has settled and plants are growing.</p>

**BEST AVAILABLE COPY**

236-6884.YLD

Pre-Reg (6884)

U.S. LABEL

Reason to Issue: To propose use on strawberries. 1/4/90 Draft: Revise per internal comments.

Date of Draft: 1/4/90 (Pre Reg) (T)  
Supersedes Pre-Reg Draft Dated: 11/21/89

EPA Reg. No. 3125-236

**RESTRICTED USE PESTICIDE**

For Sale to and Use Only By a Certified Applicator for Uses Authorized By His Certification, or By Persons Under His Direct Supervision.


NEMACUR®  
15% Granular  
SYSTEMIC INSECTICIDE - NEMATOCIDE

ACTIVE INGREDIENT:  
Ethyl 3-methyl-4-(methylthio)phenyl  
(1-methylethyl)phosphoramidate ..... 15%

INERT INGREDIENTS: ..... 85%  
100%

AMENDMENT TO PREVIOUSLY REGISTERED LABELING

ADD THE FOLLOWING:



• NEMACUR 15% Granular Systemic Insecticide - Nematocide

---

**RECOMMENDED APPLICATIONS**

CROP	PEST	DOSAGE NEMACUR 15% G	REMARKS										
Straw- berries	Nematodes	14.7 to 22 oz for any row spacing (or 12 to 18 pounds per acre on 40-inch rows).	<p>Apply specified dosage per 1000 ft of row in a 12- to 18-inch band over the row and incorporate immediately by cultivation or by sprinkler irrigation prior to transplanting. Do not apply more than one application. Do not apply within 110 days of harvest. Apply by ground equipment only.</p> <p><u>BAND WIDTH</u></p> <table><tr><td><u>12-INCH</u></td><td><u>18-INCH</u></td></tr><tr><td>14.7 oz/1000 ft</td><td>22 oz/1000 ft</td></tr><tr><td>OR</td><td>OR</td></tr><tr><td>12 lb/acre</td><td>18 lb/acre</td></tr><tr><td>(40-inch rows)</td><td>(40-inch rows)</td></tr></table> <p>Use the maximum band width and rate in fields with high populations of nematodes or in fields having a history of serious nematode damage.</p>	<u>12-INCH</u>	<u>18-INCH</u>	14.7 oz/1000 ft	22 oz/1000 ft	OR	OR	12 lb/acre	18 lb/acre	(40-inch rows)	(40-inch rows)
<u>12-INCH</u>	<u>18-INCH</u>												
14.7 oz/1000 ft	22 oz/1000 ft												
OR	OR												
12 lb/acre	18 lb/acre												
(40-inch rows)	(40-inch rows)												

236-6857.YLD

Pre-Reg (6857)

U.S. LABEL

Reason to Issue: To propose use on strawberries.

Date of Draft: 11/21/89 (Pre Reg) (I)

EPA Reg. No. 3125-236

-----

**RESTRICTED USE PESTICIDE**

For Sale to and Use Only By a Certified Applicator for Uses Authorized  
By His Certification, or By Persons Under His Direct Supervision.

**NEMACUR®**  
**15% Granular**  
**SYSTEMIC INSECTICIDE - NEMATICIDE**


**ACTIVE INGREDIENT:**

Ethyl 3-methyl-4-(methylthio)phenyl  
(1-methylethyl)phosphoramidate ..... 15%

INERT INGREDIENTS: ..... 85%  
100%

**AMENDMENT TO PREVIOUSLY REGISTERED LABELING**

ADD THE FOLLOWING:



# NEMACUR 15% Granular Systemic Insecticide - Nematicide

## RECOMMENDED APPLICATIONS

CROP	PEST	DOSAGE	REMARKS							
		NEMACUR 15% G								
		BAND: OZ/ 1000 FT OF ROW								
Straw- berries	Nematodes	14.7 to 22 oz for any row spacing (or 12 to 18 pounds per acre on 40-inch rows).	Apply specified dosage per 1000 ft of row in a 12- to 18-inch band over the row at transplanting and incorporate immediately by cultivation or by sprinkler irrigation. Do not apply more than one application. Do not apply within 110 days of harvest. Apply by ground equipment only.							
			<u>BAND WIDTH</u>							
			<table><tr><td><u>12-INCH</u></td><td><u>18-INCH</u></td></tr><tr><td>14.7 oz/1000 ft</td><td>22 oz/1000 ft</td></tr><tr><td>OR</td><td>OR</td></tr><tr><td>12 lb/acre</td><td>18 lb/acre</td></tr><tr><td>(40-inch rows)</td><td>(40-inch rows)</td></tr></table>	<u>12-INCH</u>	<u>18-INCH</u>	14.7 oz/1000 ft	22 oz/1000 ft	OR	OR	12 lb/acre
<u>12-INCH</u>	<u>18-INCH</u>									
14.7 oz/1000 ft	22 oz/1000 ft									
OR	OR									
12 lb/acre	18 lb/acre									
(40-inch rows)	(40-inch rows)									
			Use the maximum band width and rate in fields with high populations of nematodes or in fields having a history of serious nematode damage.							