

Reg # 3125-236

PM-22

184

236-7293.YLD

Pre-Reg (7293)

U.S. LABEL

Reason to Issue: To propose use on
bok choy.

Date of Draft: 7/10/91 (Pre Reg) (T)

EPA Reg. No. 3125-236

RESTRICTED USE PESTICIDE

For Sale to and Use Only By a Certified Applicator for Uses Authorized
By His Certification, or By Persons Under His Direct Supervision.


NEMACUR®
15% Granular
SYSTEMIC INSECTICIDE - NEMATOCIDE

ACTIVE INGREDIENT:
Ethyl 3-methyl-4-(methylthio)phenyl
(1-methylethyl)phosphoramidate 15%

INERT INGREDIENTS: 85%
100%

AMENDMENT TO PREVIOUSLY REGISTERED LABELING

ADD THE FOLLOWING:



RECOMMENDED APPLICATIONS

CROP	PEST	DOSAGE NEMACUR 15% G	REMARKS						
<u>VEGETABLES</u>									
Bok Choy (direct seeded only) (California only)	Nematodes	14.7 to 18.4 oz per 1000 ft of row for any row spacing	Apply specified dosage in a 12- to 15-inch band over the row and incorporate. Refer to the table below for specific band width application rates. Treatment may be applied immediately before or at planting of direct seeded crop. ON NARROW ROW CROPS, DO NOT ALLOW BANDS TO OVERLAP.						
			<table><tr><th><u>Bandwidth</u></th><th><u>oz/1000 ft of Row</u></th></tr><tr><td>12"</td><td>14.7</td></tr><tr><td>15"</td><td>18.4</td></tr></table>	<u>Bandwidth</u>	<u>oz/1000 ft of Row</u>	12"	14.7	15"	18.4
<u>Bandwidth</u>	<u>oz/1000 ft of Row</u>								
12"	14.7								
15"	18.4								

384

236-7294.YLD

Pre-Reg (7294)

U.S. LABEL

Reason to Issue: To add narrow band applications on cabbage and brussels sprouts.

Date of Draft: 7/10/91 (Pre Reg) (T)

EPA Reg. No. 3125-236

RESTRICTED USE PESTICIDE

For Sale to and Use Only By a Certified Applicator for Uses Authorized
By His Certification, or By Persons Under His Direct Supervision.


NEMACUR®
15% Granular
SYSTEMIC INSECTICIDE - NEMATOCIDE

ACTIVE INGREDIENT:
Ethyl 3-methyl-4-(methylthio)phenyl
(1-methylethyl)phosphoramidate 15%

INERT INGREDIENTS: 85%
100%

AMENDMENT TO PREVIOUSLY REGISTERED LABELING

ADD THE FOLLOWING:



RECOMMENDED APPLICATIONS

CROP	PEST	DOSAGE NEMACUR 15% G	REMARKS																		
<u>VEGETABLES</u>																					
Cabbage (includes tight heading varieties of Chinese cabbage) (direct seeded and trans- planted) (Except cabbage grown for seed)	Nematodes	7.3 to 18.4 oz per 1000 ft of row for any row spacing	Apply specified dosage in a 6- to 15-inch band over the row. Refer to table below for specific band width application rates. Application may be made prior to planting, during planting, after planting but prior to crop emergence (direct seeded cabbage), or immediately following transplanting. Treat a minimum band width of 6 inches per row for multiple rows on a single bed or a minimum 12-inch band width for a single row per bed. Utilize wider band widths in fields with a history of nematode damage. Incorporate mechanically or with overhead irrigation. On narrow row spacing, do not allow bands to overlap. Do not treat more than 50% of the total field area. Do not make more than one application per crop season.																		
Brussels Sprouts (trans- planted) (Except brussels sprouts grown for seed)			<table><tr><th><u>Bandwidth</u></th><th><u>oz/1000 ft of Row</u></th></tr><tr><td>6"</td><td>7.3</td></tr><tr><td>7"</td><td>8.6</td></tr><tr><td>8"</td><td>9.8</td></tr><tr><td>9"</td><td>11.0</td></tr><tr><td>9.5"</td><td>11.6</td></tr><tr><td>10"</td><td>12.2</td></tr><tr><td>12"</td><td>14.7</td></tr><tr><td>15"</td><td>18.4</td></tr></table>	<u>Bandwidth</u>	<u>oz/1000 ft of Row</u>	6"	7.3	7"	8.6	8"	9.8	9"	11.0	9.5"	11.6	10"	12.2	12"	14.7	15"	18.4
<u>Bandwidth</u>	<u>oz/1000 ft of Row</u>																				
6"	7.3																				
7"	8.6																				
8"	9.8																				
9"	11.0																				
9.5"	11.6																				
10"	12.2																				
12"	14.7																				
15"	18.4																				