



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

DEC 21 1993

Melvin K. Tolliver
MILES INC.
AGRICULTURE DIVISION
8400 Hawthorn Road
P.O. Box 4913
Kansas City, MO 64120

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Subject: Label Amendment Submission of 06/16/93 in Response to PR Notice 93-7
EPA Reg. No. 3125-236 ✓
NEMACUR 15% GRANULAR

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.



Recycled/Recyclable
Printed with Soy/Canola ink on paper that
contains at least 50% recycled fiber

20/12

Page 2


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

MILES INC
NEMACUR 15% GRANULAR
3125-236 06/16/93
Original Submission

Add the following statement to your label: "Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them." Refer to page 20 of the Main Labeling Guidance for placement.

NEMACUR 15% Granular

236-7821.YLD

U. S. LABEL

Base Pre-Reg. (7821)

4042

Reason to Issue: To revise in accordance with
Worker Protection Standard - PR Notice 93-3.

Date of Draft: 05/24/93(Pre-Reg)(T)

RESTRICTED USE PESTICIDE

Due to high acute oral, dermal and inhalation toxicity
For Retail Sale to and Use Only by a Certified Applicator for Uses Authorized by His
Certification, or by Persons Under His Direct Supervision.

NEMACUR®

15% Granular

SYSTEMIC INSECTICIDE - NEMATICIDE

FOR EFFECTIVE CONTROL OF NEMATODES IN CERTAIN FIELD, FRUIT AND VEGETABLE CROPS

ACTIVE INGREDIENT:

Ethyl 3-methyl-4-(methylthio)phenyl(1-methylethyl)phosphoramidate 15%

INERT INGREDIENTS..... 85%

100%

EPA Reg. No. 3125-236

EPA Est. No. 5905-GA-1

Net Weight _____ Pounds

STOP - Read the label before use.

KEEP OUT OF REACH OF CHILDREN

DANGER PELIGRO



POISON

See rear panel for Statements of Practical Treatment and Other Precautionary Statements.

Si usted no entiende la etiqueta, busque a
alguien para que se la explique a usted en
detalle.

(If you do not understand the label, find
someone to explain it to you in detail.)

~~PRECAUCION AL USUARIO: Si usted no puede
leer o entender inglés, no use este producto
hasta que la etiqueta le haya sido explicada
ampliamente.~~

~~(TO THE USER: If you cannot read or
understand English, do not use this product
until the label has been fully explained to you.)~~

ACCEPTED
with COMMENTS
In EPA Letter Dated

MILES

DEC 22 1993

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

3125-236

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER



POISON

Fatal if swallowed, or absorbed through the skin. Causes eye irritation. May be fatal if inhaled. Do not get in eyes, on skin, or on clothing. Do not breathe dust. ~~Wear protective clothing, rubber gloves, and goggles or face shield when handling. Wash thoroughly with soap and water after handling and before eating or smoking. Remove and wash contaminated clothing before reuse.~~

Do not contaminate feed or food. ~~Keep all unprotected persons out of the operating area or vicinity. Wash hands, arms, and face thoroughly with soap and warm water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Keep out of the reach of children.~~

Not for use or storage in or around the home.

Personal Protective Equipment
Applicators and other handlers must wear:

- Chemical-resistant protective suit
- Water-proof gloves
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- For exposures in enclosed areas, a respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G)
- For exposures outdoors, dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements: When handlers use enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

SYMPTOMS OF POISONING: A sense of "tightness" in the chest. Sweating. Contracted pupils. Stomach pains. Vomiting and diarrhea.

STATEMENTS OF PRACTICAL TREATMENT

Organophosphate

In case of poisoning call a physician immediately. Have patient lie down and keep quiet. If **swallowed**, **vomiting should be induced**. Administer water freely and induce vomiting by giving one dose (1/2 oz or 15 ml) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Never give anything by mouth to an unconscious person. Professional medical assistance should be secured immediately. If on **skin**, remove contaminated clothing and wash skin immediately with soap and warm water. If **eyes are contaminated**, wash with flowing water for at least 15 minutes.

If inhaled, remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth.

NOTE TO PHYSICIAN: ANTIDOTE - Administer atropine sulfate in large therapeutic doses. Repeat as necessary to the point of tolerance. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Compound inhibits cholinesterase resulting in stimulation of the central nervous system, the parasympathetic nervous system, and the somatic motor nerves. Do not give morphine. Watch for pulmonary edema, which may develop in serious cases of poisoning even after 12 hours. At first sign of pulmonary edema, the patient should be placed in an oxygen tent and treated symptomatically.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Cover, incorporate, or pick up spilled granules at row ends or turn areas. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Ground Water Advisory: Fenamiphos is known to leach through soil and has been found in ground water as a result of agriculture use. Users are advised not to apply in areas where soils are permeable, particularly where ground water is used for drinking water. Consult with the pesticide state lead agency for information regarding soil permeability and aquifer vulnerability in your area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Chemical-resistant protective suit
- Waterproof gloves
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

IMPORTANT: Read these entire Directions and Conditions of Sale before using NEMACUR 15% Granular systemic insecticide-nematicide.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE THE DIRECTIONS FOR CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED FOR A RANGE OF WEATHER CONDITIONS SIMILAR TO THOSE WEATHER CONDITIONS THAT ARE ORDINARY AND CUSTOMARY IN THE GEOGRAPHIC AREA WHERE THE PRODUCT IS USED. INSUFFICIENT CONTROL OF PESTS AND/OR INJURY TO THE CROP TO WHICH THE PRODUCT IS APPLIED MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL WEATHER, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO OTHER CROPS, ANIMALS, MAN, OR THE ENVIRONMENT. MILES OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL WEATHER, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF MILES AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

Consult your State Agricultural Experiment Station or Extension Service Specialist or Miles representative for guidance on the selection of dosage rates and timing of applications appropriate for your specific crop and pest. State regulations may contain additional restrictions or requirements. Check the "Remarks" and "Restrictions" sections of this label for additional directions.

Incorporation Recommendations: Unless otherwise specified, NEMACUR 15% Granular must be incorporated into the soil for optimum results. Incorporation may be accomplished by irrigation or mechanical equipment. For applications made at planting, the action of some planters during the planting operation may provide adequate incorporation. When utilizing planters which do not provide adequate incorporation of NEMACUR 15% Granular into the soil, additional incorporation may be achieved by use of equipment specifically designed for incorporation purposes behind the planter. Precautions should be taken to avoid disturbing seed or transplant replacement.

RECOMMENDED APPLICATIONS									
CROP	PEST	DOSAGE NEMACUR 15% G	REMARKS						
FIELD CROPS Cotton	Thrips (early season reduction)	6 to 8 oz per 1000 ft of row for any row spacing (5 to 6.5 lb per acre on 40 inch row).	In furrow: Apply specified dosage per 1000 ft of row.						
	Nematodes Thrips	8 to 12 oz per 1000 ft of row for any row spacing (6.5 to 9.8 lb per acre on 40 inch row spacing).	<p>In furrow: Apply specified dosage per 1000 ft of row.</p> <p>Suggested use rates of NEMACUR 15% G in oz/1000 feet of row based on nematode infestation levels:*</p> <table><tr><td>Low</td><td>Moderate</td><td>High*</td></tr><tr><td>8</td><td>10</td><td>12</td></tr></table> <p>*Use the 12 oz/1000 feet rate in fields with a history of nematode problems.</p>	Low	Moderate	High*	8	10	12
	Low	Moderate	High*						
8	10	12							
IMPORTANT: Do not overlap bands on narrow row crops. Do not feed or graze cotton foliage. Under very heavy thrip pressure, additional control measures may be needed.									
Peanuts	Nematodes Thrips (early season reduction)	11 to 18.7 oz for any row spacing (or 10 to 17 lb per acre on 36-inch rows)	<p>Apply specified dosage per 1000 ft of row as a 12-inch band at planting. Incorporate the granules into the soil. ON NARROW ROW CROPS, DO NOT ALLOW BANDS TO OVERLAP.</p> <p>Use the high rate in fields with high populations of nematodes, in fields where root-knot species other than southern root-knot, <i>Meloidogyne incognita</i>, are present or in fields having a history of serious nematode damage.</p> <p>Do not feed or graze green peanut vines or peanut vine hay. Do not hog down treated peanut fields.</p>						
VEGETABLES Bok Choy (direct seeded only) (California only)	Nematodes	14.7 to 18.4 oz per 1000 ft of row for any row spacing	<p>Apply specified dosage in a 12- to 15-inch band over the row and incorporate. Refer to the table below for specific band width application rates. <i>Treatment may be applied immediately before or at planting of direct seeded crop.</i> ON NARROW ROW CROPS, DO NOT ALLOW BANDS TO OVERLAP.</p> <table><tr><td>Bandwidth</td><td>oz/1000 ft of Row</td></tr><tr><td>12"</td><td>14.7</td></tr><tr><td>15"</td><td>18.4</td></tr></table>	Bandwidth	oz/1000 ft of Row	12"	14.7	15"	18.4
Bandwidth	oz/1000 ft of Row								
12"	14.7								
15"	18.4								

RECOMMENDED APPLICATIONS																					
CROP	PEST	DOSAGE NEMACUR 15% G	REMARKS																		
VEGETABLES (Continued) Cabbage (includes tight heading varieties of Chinese cabbage) (direct seeded and transplanted) (Except cabbage grown for seed) Brussels sprouts (transplanted) (Except brussels sprouts grown for seed)	Nematodes	7.3 to 18.4 oz per 1000 ft of row for any row spacing	Apply specified dosage in a 6- to 15-inch band over the row. Refer to table below for specific band width application rates. Application may be made prior to planting, during planting, after planting but prior to crop emergence (direct seeded cabbage), or immediately following transplanting. Treat a minimum band width of 6 inches per row for multiple rows on a single bed or a minimum 12-inch band width for a single row per bed. Utilize wider band width in fields with a history of nematode damage. Incorporate mechanically or with overhead irrigation. On narrow row spacing, do not allow bands to overlap. Do not treat more than 50% of the total field area. Do not make more than one application per crop season. <table><tr><td>Bandwidth</td><td>oz/1000 ft of Row -</td></tr><tr><td>6"</td><td>7.3</td></tr><tr><td>7"</td><td>8.6</td></tr><tr><td>8"</td><td>9.8</td></tr><tr><td>9"</td><td>11.0</td></tr><tr><td>9.5"</td><td>11.6</td></tr><tr><td>10"</td><td>12.2</td></tr><tr><td>12"</td><td>14.7</td></tr><tr><td>15"</td><td>18.4</td></tr></table>	Bandwidth	oz/1000 ft of Row -	6"	7.3	7"	8.6	8"	9.8	9"	11.0	9.5"	11.6	10"	12.2	12"	14.7	15"	18.4
Bandwidth	oz/1000 ft of Row -																				
6"	7.3																				
7"	8.6																				
8"	9.8																				
9"	11.0																				
9.5"	11.6																				
10"	12.2																				
12"	14.7																				
15"	18.4																				
Eggplant	Nematodes	14.7 oz (or 13.4 lb per acre for 36-inch rows)	Apply specified dosage per 1000 ft of row in a 12-inch band over the row at transplanting and immediately incorporate the granules into the soil. ON NARROW ROW CROPS, DO NOT ALLOW BANDS TO OVERLAP.																		
Garlic	Bulb and Stem Nematode	9.2 to 18.4 oz/1000 ft of planting furrow for any rows spacing (15 to 30 lb per acre on 40-inch beds with two seed lines [rows] per bed)	Apply specified dosage in-furrow with the garlic seed (cloves) at planting. For best results, plant the crop within 7 to 10 days after cracking (seed preparation).																		
Okra (Except California)	Nematodes	14.7 to 18.4 oz (or 13.4 to 16.7 lb per acre on 36-inch rows)	Apply specified dosage per 1000 ft of row as a 12- to 15-inch band at planting. Incorporate the granules into the soil. Use low rate in a 12-inch band. Use high rate in a 15-inch band. <table><tr><td colspan="2">BAND WIDTH (36-inch row spacing)</td></tr><tr><td>12-inch</td><td>15-inch</td></tr><tr><td>14.7 oz/1000 ft of row or 13.4 lb/acre</td><td>18.4 oz/1000 ft of row or 16.7 lb/acre</td></tr></table> ON NARROW ROW CROPS, DO NOT ALLOW BANDS TO OVERLAP.	BAND WIDTH (36-inch row spacing)		12-inch	15-inch	14.7 oz/1000 ft of row or 13.4 lb/acre	18.4 oz/1000 ft of row or 16.7 lb/acre												
BAND WIDTH (36-inch row spacing)																					
12-inch	15-inch																				
14.7 oz/1000 ft of row or 13.4 lb/acre	18.4 oz/1000 ft of row or 16.7 lb/acre																				

RECOMMENDED APPLICATIONS									
CROP	PEST	DOSAGE NEMACUR 15% G	REMARKS						
VEGETABLES (Continued) Non-Bell Peppers (California, Georgia and Puerto Rico only)	Nematodes	14.7 to 18.4 oz (or 13.4 to 16.7 lb per acre on 36-inch rows)	Apply specified dosage per 1000 ft of row in a 12-inch band over the row at planting and incorporate the granules into the soil. Use the high rate in fields with high populations of nematodes or in fields having a history of serious nematode damage.						
FRUIT Citrus (In California do not apply to Kumquat, Tangelo or Citrus hybrids)	Nematodes Citrus root weevil complex including Fuller rose beetle (Suppression)	33-1/2 to 67 lb	<p>Apply specified dosage per acre in a band in up to 2 applications per season, not to exceed 67 pounds of NEMACUR 15% Granular per acre per season. Apply to the soil with ground equipment and incorporate immediately using mechanical incorporation or irrigation. Center the treated band on the tree row using a band width of 50% of the row spacing and covering the feeder root system of the plant.</p> <p>Under dry conditions, follow with overhead or floor irrigation. Do not apply more than 10 pounds active ingredient (67 pounds of NEMACUR 15% Granular) per acre per season. Do not apply within 30 days of harvest. Do not graze livestock on treated areas. On non-bearing citrus make initial applications when trees have recovered from transplanting shock and have initiated new growth, provided that one or more irrigations have been applied and soil has settled around the roots. Phytotoxicity may occur if applications are made before soil has settled and plants are growing.</p>						
Strawberries	Nematodes	14.7 to 22 oz for any row spacing (or 12 to 18 lb per acre on 40-inch rows)	<p>Apply specified dosage per 1000 ft of row in a 12- to 18-inch band over the row and incorporate immediately by cultivation or by sprinkler irrigation prior to transplanting. Do not apply more than one application. Do not apply within 110 days of harvest. Apply by ground equipment only.</p> <table><tr><th colspan="2">BAND WIDTH</th></tr><tr><th>12-INCH</th><th>18-INCH</th></tr><tr><td>14.7 oz/1000 ft or 12 lb/acre (40-inch rows)</td><td>22 oz/1000 ft or 18 lb/acre (40-inch rows)</td></tr></table> <p>Use the maximum band width and rate in fields with high populations of nematodes or in fields having a history of serious nematode damage.</p>	BAND WIDTH		12-INCH	18-INCH	14.7 oz/1000 ft or 12 lb/acre (40-inch rows)	22 oz/1000 ft or 18 lb/acre (40-inch rows)
BAND WIDTH									
12-INCH	18-INCH								
14.7 oz/1000 ft or 12 lb/acre (40-inch rows)	22 oz/1000 ft or 18 lb/acre (40-inch rows)								

RECOMMENDED APPLICATIONS			
CROP	PEST	DOSAGE NEMACUR 15% G	REMARKS
FRUIT (Continued) Pineapple (Puerto Rico only)	Nematodes	66-2/3 to 133-1/3 pounds	Apply specified dosage (10 to 20 lb active ingredient) per acre over the entire area to be treated before planting and thoroughly incorporate into the soil mechanically or with sufficient rainfall or irrigation (not to exceed the depth of the root zone).
	<p>NOTE: Postplanting applications of NEMACUR 3 can be made in addition to a preplant application of NEMACUR 15% Granular. Refer to the labeling for NEMACUR 3 for directions and dosage rates.</p> <p>Do not apply more than a total of 20 lb active ingredient per acre per crop season regardless of the formulation or method of application. Do not use forage or fodder for animal feed.</p>		
Non-bearing Strawberry nursery stock	Nematodes	17 oz for any row spacing (or 13.8 pounds per acre on 40- inch rows)	Apply specified dosage per 1000 ft of row in a 12- inch band over the row immediately after transplanting and incorporate with cultivation equipment. Apply a second application in a 12-inch band over the row 8 weeks after transplanting. Incorporate by cultivation or by sprinkler irrigation (1 inch of water) immediately following application. Do not apply more than two applications to non-bearing nursery stock. Use only on strawberry nursery stock that will not be harvested for fruit within 20 months after application.

RESTRICTIONS

Any food crop not specified on this label may be planted into treated areas 120 days after the last application. Any cover crops that are planted during the 120 day period must be plowed under and not grazed.

Do not use on other crops. Use only according to label directions.

Do not apply this product with a knapsack or similar equipment that is placed on the user's body.

PROTECTIVE CLOTHING AND EQUIPMENT

~~USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT DURING MIXING/LOADING, APPLICATION, REPAIR AND CLEANING OF MIXING, LOADING AND APPLICATION EQUIPMENT, DISPOSAL OF THE PESTICIDE, AND EARLY REENTRY INTO TREATED AREAS: Protective suit of one or two pieces covering all parts of the body except head, hands, and feet; chemical-resistant gloves; chemical-resistant shoes (or chemical-resistant shoe covers or chemical-resistant boots); and a NIOSH or MSHA approved respirator. In addition, mixer/loaders must wear a chemical-resistant apron. During equipment repair and cleaning, the respirator need not be worn. During early reentry after dust has settled and vapors have dispersed, the respirator need not be worn.~~

~~IF MIXING/LOADING IS PERFORMED USING A CLOSED SYSTEM, THE FOLLOWING PROTECTIVE~~

~~CLOTHING AND EQUIPMENT MAY BE WORN AS AN ALTERNATIVE: Long-sleeved shirt and long-legged pants; chemical-resistant gloves; chemical-resistant apron; shoes and socks.~~

~~IF APPLICATION IS PERFORMED USING AN ENCLOSED CAB, THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT MAY BE WORN AS AN ALTERNATIVE: Long-sleeved shirt and long-legged pants; shoes, and socks. Chemical-resistant gloves must be available in the cab and must be worn when exiting. This clothing is inadequate to protect you during equipment repair or cleaning, reentry, or pesticide disposal work.~~

~~IMPORTANT! If pesticide comes in contact with skin, wash off with soap and water. Always wash hands, face, and arms with soap and water before smoking, eating, drinking, or toileting.~~

PROTECTIVE CLOTHING AND EQUIPMENT (Continued)

~~AFTER WORK: Before removing gloves, wash them with soap and water. Take off all work clothes and shoes. Shower using soap and water. Wear clean clothes. Do not reuse contaminated clothing. Personal clothing worn during work must be laundered separately from household articles. Store protective clothing separately from personal clothing. Clean or~~

~~launder protective clothing after each use. Respirators must be cleaned and filters replaced according to instructions included with the respirator. Protective clothing and equipment that becomes heavily contaminated must be destroyed according to state and local regulations. HEAVILY CONTAMINATED CLOTHING CANNOT BE ADEQUATELY DECONTAMINATED.~~

REENTRY STATEMENTS

~~Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not reenter treated area for 48 hours unless appropriate protective clothing is worn. Any person coming in direct contact with treated soil prior to incorporation or irrigation, or with treated wet soil after the initial irrigation, must wear protective clothing and equipment as described in the section "Protective Clothing and Equipment" on this label. Wash thoroughly with soap and water after handling and before eating or smoking.~~

~~Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.~~

~~(The following may appear on supplemental labeling accompanying the product)~~

~~Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings should include these statements found under the label heading "Precautionary Statements" and "Re-entry Statements" included on this label. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: DANGER, area treated with NEMACUR 15% Granular on (date of application). See "Protective Clothing and Equipment" for use of proper protective clothing. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash with flowing water for at least 15 minutes. If swallowed, vomiting should be induced. (See Precautionary Statements on label).~~

STORAGE AND DISPOSAL

Pesticide Disposal: Do not contaminate water, food or feed by storage or disposal.

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Do not use container in connection with food, feed, or drinking water. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticide as directed above. In spill or leak incidents, keep unauthorized people away. You may contact the Miles Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Miles Kansas City Emergency Response Telephone No. is 816-242-2582, or contact Chemtrec at 800-424-9300.

Miles Inc.
Crop Protection Products
P.O. Box 4913, Kansas City, MO 64120-0013