

Staffel's



ACTIVE INGREDIENTS

Carbaryl
(1-naphthyl
methylcarba-
mate) 4% by wt.
Metalde-
hyde 1% by wt.

**INERT
INGREDIENTS** 95% by wt.

Contains SEVIN*
carbaryl insecticide
and metaldehyde

USDA Reg. No. 3286-19

WARRANTY

1. The formulation guarantees and warrants (a) that the active ingredient content and the total net weight are as stated within lawful limits and (b) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states.
2. The formulator makes no other express or implied guarantee, warranty, or representation.
3. No modification of this warranty and disclaimer is authorized, except by specific references to them in writing by an employee of the formulator.

*SEVIN is the registered trade mark of Union Carbide Corporation for carbaryl insecticide.

| Crop | Dosage Limitations to Meet Residue Tolerances | Days Between Final Application and Harvest |
|---|---|--|
| Cabbage, broccoli, cauliflower, lettuce (head), beets (roots) | 50 lbs. | 3 |
| Lettuce (leaf), spinach, beets (tops) | 50 lbs. | 14 |
| Beans, carrots, tomatoes, peppers, potatoes | 50 lbs. | None |
| Cucumbers, melons, squash | 25 lbs. | None |
| Pears | 25 lbs. | None |
| Asparagus (Do not repeat application within 3 days.) | 50 lbs. | 1 |
| Citrus | 50-60 lbs. | 5 |
| Apples, cherries, peaches, pears, plums | 50-60 lbs. | 1 |
| Apricots, nectarines, grapes | 50-60 lbs. | 3 |
| Ornamentals, borders | 50-100 lbs. | None |

Special instructions for improved performance are:

1. Slugs and snails favor dark, damp places. Bait where these pests stay during daylight hours.
2. Be sure area to be baited is damp. In dry weather, water area well before baiting.
3. Apply bait in evening as these pests feed mainly at night.
4. Until under control bait should be renewed at least every 7-10 days or more frequently if necessary.
5. Since snails and slugs migrate extensively, when invasions from adjacent areas are anticipated, a barrier of the bait will aid in control.

Mfd. by THE FERD STAFFEL COMPANY
SAN ANTONIO, TEXAS



BUG BAIT

**WARNING: Keep out of reach of children.
See side panels for other cautions.**

**WARNING: Keep out of reach of children and animals.
WARNING: Harmful dust. Avoid contact with skin, eyes and clothing. Do not take internally. Avoid contact with skin. Wear regular long-sleeve work clothes. Change to clean clothes daily. Wash hands and face before eating. Wash thoroughly after handling.**

WARNING: May be fatal if inhaled or swallowed.

Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Apply this product only as specified on this label.

NOTE TO PHYSICIAN:

Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

ACCEPTED
Dec-20-1971
UNDER THE FEDERAL INSECTICIDE
AND RODENTICIDE ACT

