

ACCEPTED


SEP 1 29 1975

UNDER... TRADE
FUNGICIDE... ACT
FOR ECOR...
ED UNDER NO. 513-277...
TO ATTACHED...



SENCOR™

50% Wettable Powder
HERBICIDE



For control
of certain grasses
and broadleaf weeds

IMPORTANT

SUPPLEMENT TO PACKAGE LABEL

Directions for use,
precautions,
conditions of sale

SENCOR™

50% Wettable Powder
HERBICIDE



CAUTION
Keep Out of Reach of
Children

(See other pages
of booklet for
additional caution statements)

Guarantee

ACTIVE INGREDIENT:	
4-Amino-6-(1, 1-dimethylethyl)-3-	
(methylthio)-1,2,4-triazin-	
5(4H)-one	50%
INERT INGREDIENTS:	50%
	<hr/>
	100%

Container Size: 5 lbs.

EPA Reg. No. 3125-277-AA

U.S. Patent No. 3,671,523

SENCOR is a TM of the Parent Company of
Farbenfabriken Bayer GmbH, Leverkusen

STORE IN A COOL DRY PLACE



CHEMAGRO AGRICULTURAL DIVISION
MOBAY CHEMICAL CORPORATION
Box 4913 Kansas City Mo 64120

Directions for use

Important:

Read these entire Directions and Conditions of Sale, including the Warranty and Limitation of Damages provision, before using SENCOR 50% Wettable Powder

Conditions of sale:

These Directions for Use reflect the opinion of experts based on tests of effectiveness, of toxicity to plants and laboratory animals and of residues on or in food and feed under normal conditions of use. However, because of the wide range of conditions under which this product may be used, even though label directions are followed, it is impossible to eliminate all risks associated with its use because of abnormal conditions beyond the control of the seller. These include, but are not limited to the presence of other products or materials, not recommended on this label, that may be in or on the soil. Chemagro and the Seller offer, and the Buyer uses, this product subject to the understanding that the Buyer assumes all such risks.

General Information

Mixing:

Fill the spray tank half full with clean water and add SENCOR 50% Wettable Powder at the recommended rate. Mix thoroughly and add water to fill spray tank. Maintain sufficient agitation during mixing and spraying operations to insure a uniform spray mixture.

It is essential that the sprayer be accurately calibrated before applying SENCOR 50% Wettable Powder. Avoid boom-overlaps that will increase dosages above those recommended. Check the sprayer frequently during application to

be sure it is working properly and delivering a uniform spray pattern.

Sprayer Cleanup:

Spray equipment must be thoroughly cleaned to remove remaining traces of herbicide that might injure other crops to be sprayed. Drain any remaining spray solution of SENCOR from the spray tank and discard in non-crop areas away from water supplies. Rinse the spray tank and refill with water, adding a heavy-duty detergent at the rate of one cup per 20 gallons of water. Recycle this mixture through the equipment for 5 minutes and spray out. Repeat this procedure twice. Fill the spray tank with clean water, recycle for 5 minutes, and spray out. Clean pump and nozzle screens thoroughly. Wash away any spray mixture from the outside of spray tank, nozzles or spray rig.

Soybeans

SENCOR 50% Wettable Powder is recommended for use in ground or aircraft spray equipment as a preemergence surface application to soybeans.

A minimum amount of soil moisture is required to activate SENCOR 50% Wettable Powder. In areas of low rainfall, pre-emergence applications to dry soil should be followed with light irrigation. Do not apply a heavy irrigation immediately after application. As with many surface-applied herbicides, weed control and crop tolerance may vary with rainfall and/or soil texture.

The soil surface and seed bed should be well-prepared and free of trash and clods. Seed must be planted at least 1½ inches below the soil surface. Distribute the chemical uniformly over the area to be treated.

Ground Application:

SENCOR 50% Wettable Powder is recommended for use with ground equipment as a preemergence surface application for the control of certain grasses and broadleaf weeds in soybeans. Make a single application as a band or broadcast spray during the planting or in a separate operation after planting but before the crop emerges.

Use a standard low pressure (20 to 40 psi) herbicide boom sprayer equipped with fan tips and screens no finer than 50-mesh in nozzle and in-line strainers. Spray equipment should be calibrated to deliver 10 to 40 gallons of spray mixture per acre broadcast or, for band application 1/4 to 1 gallon per inch of band width, regardless of row spacing.

Aerial Application:

SENCOR 50% Wettable Powder is recommended for use in aerial spray equipment as a preemergence surface application for the control of certain grasses and broadleaf weeds in soybeans. Make a single application after planting but before the crop emerges.

Use a standard low pressure (20 to 40 psi) aerial herbicide boom sprayer equipped with fan tips and screens no finer than 50-mesh in nozzle and in-line strainers. Aerial spray equipment should be calibrated to apply the proper amount of SENCOR 50% Wettable Powder in 2 to 10 gallons of spray mixture per acre. SENCOR 50% Wettable Powder mixes readily with water for these concentrate aerial sprays. However, if the spray mix is allowed to stand unagitated, redispersion of the settled powder becomes difficult. Therefore, maintain agitation or recirculation of the spray mix until tank is empty.

Weeds Controlled:

SENCOR 50% Wettable Powder applied preemergence to soybeans, according to directions, will effectively control

Grass Weeds:

Crabgrass	Yellow foxtail
Crowfootgrass	Seedling johnsongrass
Giant foxtail	Fall panicum
Green foxtail	Signalgrass (Brachiaria)

Broadleaf Weeds:

Spurred anoda	Pennsylvania smartweed
Beggartlice	Spotted spurge
Buffalobur	Sunflower
Carpetweed	Velvetleaf
Cocklebur	
Jimsonweed	
Lambsquarters	
Venice mallow	
Wild mustard	
Prostrate pigweed	
Redroot pigweed (Carelessweed)	
Smooth pigweed	
Florida purslane	
(Florida pusley, <i>Richardia scabra</i>)	
Common ragweed	
Sesbania (Coffeebean, Coffeeweed)	
Shepherdspurse	
Sicklepod (Cassia)	
Prickly sida (Spiny sida, Teaweed)	

Hard to control weeds:

Although SENCOR 50% Wettable Powder may not provide commercially acceptable control in every instance, it will suppress growth of the following weeds and reduce their competition with soybean plants.

Grass Weeds:

Barnyardgrass (watergrass)
Yellow nutsedge

Broadleaf Weeds:

Common purslane

Recommended Applications Preemergence Soy beans

Remarks

BROADCAST: Select the proper dosage as shown below based on soil type and percent organic matter. Apply proper dosage in 10 to 40 gallons of water per acre with ground equipment or 2 to 10 gallons of water per acre with aircraft equipment as a broadcast spray to the soil surface. Application may be made during planting or as a separate operation after planting but before the crop emerges. Use flat fan spray tips or suitable nozzles designed for broadcast application of herbicides with boom sprayers. Do not apply to sandy soils. Do not apply to light soils (sandy loam and loamy sand) containing less than 2% organic matter. Seed must be planted at least 1½ inches below soil surface. Do not incorporate into the soil. Do not apply more than once per season. Do not use treated vines for feed or forage.

Pounds of SENCOR 50% W.P. per acre			
	ORGANIC MATTER		
	Less than 2%	2 to 4%	Over 4%
LIGHT SOILS (Sandy loam, loamy sand)	DO NOT USE	¾	1
MEDIUM SOILS (Silt loam, silt loam, sandy clay, sandy clay loam)	¾ to 1	1 to 1¼	1¼ to 1½
HEAVY SOILS (Silt clay, silty clay, loam clay, clay loam)	1 to 1¼	1¼ to 1½	1½ to 1¾
Mississippi Delta Only	1½	1¾	2

BAND: Use proportionally less than 50% W.P. per acre.

Examples: To treat a 15-inch band on rows 30 inches apart, use one-half of the broadcast rate. To treat a 14-inch band on rows 42 inches apart, use one-third of the broadcast rate.

Use flat fan spray tips mounted behind press or row wheel for application during planting. 50% W.P. may be applied in a separate operation after planting but before the crop emerges. Weeds between treated bands should be removed by cultivating as needed, using protective fenders to keep freshly turned soil off treated bands.

Do not apply to sandy soils. Do not apply to light soils (sandy loam and loamy sand) containing less than 2% organic matter. Seed must be planted at least 1½ inches below soil surface. Do not incorporate into the soil. Do not apply more than once per season. Do not use treated vines for feed or forage.

Special Precautions

Injury may occur to **soybeans** following the application of SENCOR 50% Wettable Powder on soils having a calcareous surface area or a pH of 7.5 or higher.

Due to the sensitivity of certain soybean varieties, SENCOR 50% Wettable Powder is not recommended for use on Tracy, Semmes, Altona, Vansoy or Coker 102.

If initial planting does not produce a satisfactory stand, fields treated with SENCOR 50% Wettable Powder alone, following Treflan, or in combination with Lasso or Treflan may be replanted to soybeans. If replanting is necessary, do not rework soil. Do not apply a second treatment as injury to soybeans may occur.

Uses of SENCOR in Combinations with Other Herbicides

SENCOR/TREFLAN[®] Overlay Application

SENCOR may be applied as preemergence broadcast or band overlay application following a pre-plant incorporated treatment of TREFLAN. Consult the TREFLAN label for specific directions for use, recommendations, restrictions and any additional weeds not specified on this label.

SENCOR/TREFLAN Tank Mix Application

A single application of a tank mixed combination of SENCOR 50% WP and TREFLAN EC will control more weeds and grasses in soybeans than when either herbicide is used alone.

Prepare the soil surface by deep plowing, offset disking or tandem disking prior to the application of the herbicide combination. The soil surface should be well prepared and free of clods and trash.

Mixing and Spray Equipment:

Fill the spray tank half full of clean water add the recommended amount of SENCOR and TREFLAN. Mix thoroughly and add water to fill spray tank. Apply in 10 to 40 gallons of water per acre using any properly calibrated low pressure herbicide sprayer equipped with herbicide tips and screens no finer than 50 mesh in-nozzle and in-line strainers. Agitate thoroughly before and during application with good bypass agitation.

Sprayers should be accurately calibrated before applying SENCOR/TREFLAN. Avoid boom overlaps that will increase rates above those recommended. Check the

sprayer frequently during the application to be sure it is working properly and delivering a uniform spray pattern. As the amount of water used (spray volume) decreases, the importance of accurate calibration and uniform application increases.

Application:

Apply SENCOR/TREFLAN to the soil surface and incorporate in the same operation, if possible. Variable weed control may result from delayed incorporation if SENCOR/TREFLAN are applied to a wet, warm soil surface or if the wind velocity is 10 miles per hour or higher. Use machinery that mixes SENCOR/TREFLAN thoroughly with the soil. Incorporation may be delayed up to 8 hours after application. Shallow incorporation with implements set to cut less than 2 inches deep may result in erratic weed control. Do not use spike or spring-tooth harrows alone for incorporation.

Broadcast Rates

Soil Texture	TREFLAN EC Pints per Acre	SENCOR 50% WP Pounds per Acre
Light (sandy loam, loamy sandy) ^a	1 pint	½ lb
Medium (loam, silt loam, silt, sandy clay, sandy clay loam)	1½ pints	¾ lb
Heavy (Silty clay, silty clay loam, ^b clay, clay loam)	2 pints	1 lb

^aDo not use on light soils with less than 1% organic matter.

^bSilty clay loam soils are transitional soils and may be classified as medium textured soils in some regions of the U.S.

Incorporation Equipment:

1. Set PTO-driven equipment (tillers, cultivators, hoes) to cut 2 to 3 inches deep and space rotors to provide a clean sweep of the soil. PTO equipment should not be operated at a speed greater than 4 miles per hour.

2. Set disk to cut 3 to 4 inches deep and operate twice in different directions at 4 to 6 miles per hour.

3. Set mulch treader and other similar disk-type implements to cut 3 to 4 inches deep and operate twice in different directions at 5 to 8 miles per hour.

For Light and Medium Textured Soils Only

4. Set rolling cultivator to cut 2 to 4 inches deep and operate twice at 6 to 8 miles per hour. Set bed conditioner (Do-All) to cut 2 to 4 inches deep and operate at 4 to 6 miles per hour.

Weeds Controlled:

SENCOR 50% Wettable Powder and TREFLAN EC tank mix combination properly applied and soil incorporated up to 10 days before planting will effectively control the following weeds:

Annual Grasses	Annual Broadleaves
Annual bluegrass	Carelessweed
Barnyardgrass (watergrass)	Carpetweed
Bromegrass	Chickweed
Brachiaria (Signalgrass)	Common ragweed
Cheat	Florida purslane (Florida pusley)
Crabgrass	Goosefoot
Foxtails (including giant foxtail)	Hemp sesbania
Goosegrass	Jimsonweed
Johnsongrass (from seed)	Kochia
glerice	Knotweed
Jbur	Lambsquarters
Sprangletop	Pigweed (spiny redroot)
Stinkgrass	Prickly sida
Texas panicum	Purslane
Wild cane (Shattercane)	Smartweed
	Stinging nettle
	Russian thistle
	Velvetleaf
	Venice mallow
	Wild mustard

NOTE: Where cocklebur is a serious problem, use the preemergence rates recommended for SENCOR 50% WP alone as an overlay application following a preplant incorporated treatment of TREFLAN (see "SENCOR/TREFLAN Overlay Application" section on this label.)

Control of cocklebur, morningglory and giant ragweed with SENCOR/TREFLAN tank mixes may be erratic, ranging from poor to excellent depending upon soil temperature, time of germination, depth of weed seed in soil, amount of timing of soil moisture. Control may be improved with timely cultivation.

Special Precautions

Applied according to directions and under normal growing conditions, SENCOR/TREFLAN will not reduce soybean yields. Over-application may result in stunting, crop injury or soil residue. Uneven application or improper soil incorporation of the tank mix can result in erratic weed control or crop injury. Seedling disease, cold weather, deep planting, excessive moisture, high soil pH, high salt concentration or drought may weaken soybean seedlings and increase possibility of damage from the tank mix. Under these conditions delayed crop development or reduced yields may result.

In the Central United States, do not plant sorghum or oats for 12 months where the tank mix has been applied unless 25 inches or more of irrigation and/or rainfall (total) was used to produce the crop. If less than 25 inches total water was used to produce the crop during the year, do not plant either crop for 18 months after the tank mix application. Cool wet weather conditions during the early stage of growth may increase the possibility of injury to sorghum.

Do not use treated vines for feed or forage. For additional precautions, restrictions, limitations and sprayer clean-up information, refer to the appropriate section of this label.

SENCOR/LASSO*
Tank Mix Application

SENCOR 50% Wettable Powder may be applied in a tank mix combination with LASSO as a preemergence broadcast or band application to soybeans in accordance with the specified soil types and dosages recommended. Apply proper dosage in 10 to 40 gallons of water per acre with ground equipment or 2 to 10 gallons of water per acre with aircraft equipment as a broadcast spray to the soil surface.

For specific information regarding spray equipment, dilution rates, mixing, directions for use, methods of application, limitations and restrictions refer to the appropriate section of this label.

Refer to the LASSO label for pertinent recommendations, directions for use, restrictions and any additional weeds not specified on this label.

SENCOR Pounds 50% W.P.
plus LASSO Quarts 4 E.C.
SOYBEANS

SOIL TEXTURE	2% or Less Organic Matter	Greater than 2% Organic Matter
	SENCOR + LASSO LBS./ACRE QTS./ACRE	SENCOR + LASSO LBS./ACRE QTS./ACRE
T SOILS (Sandy Loam)	½ + 1½ to 2	¾ + 1½ to 2
MEDIUM SOILS (Loam, silt loam, silt, sandy clay, sandy clay loam)	¾ + 1½ to 2	1 + 1½ to 2
HEAVY SOILS (Silty clay, silty clay loam, clay, clay loam)	1 + 2	1½ + 2 to 2½
Mississippi Delta Only (Silty clay to heavy clay)	SENCOR LBS./ACRE + Lasso QTS./ACRE 2 + 2 to 2½	

Do not use Lasso plus SENCOR on sand or loamy sand soils with less than 2% organic matter. Do not use on muck soils.

SENCOR/SURFLAN
Tank Mix Application

A single application of a tank-mixed combination of SENCOR 50 WP and Surflan 75 WP will control more weeds and grasses in soybeans than when either herbicide is used alone. The combination, properly applied as pre-emergence herbicides, will control the following:

- | | |
|-----------------------|---|
| Grasses | Broadleaves |
| Barnyardgrass | Common purslane |
| Crabgrasses | Common ragweed |
| Crowfootgrass | Florida purslane |
| Giant foxtail | (Florida pusley,
<i>Richardia scabra</i>) |
| Green foxtail | Jimsonweed |
| Yellow foxtail | Lambsquarters |
| Goosegrass | Sesbania (Coffeebean,
Coffeeweed) |
| Seedling johnsongrass | Prostrate pigweed |
| Browntop millet | Redroot pigweed |
| Fall panicum | (Carlessweed) |
| Signalgrass | Smooth pigweed |
| (Brachiaria) | Prickly sida |
| | (Spiny sida, Teaweed) |
| | Redweed |
| | Sicklepod (Cassia) |
| | Pennsylvania smartweed |
| | Velvetleaf |

Cocklebur, morningglory and giant ragweed - control of these weeds may be erratic, ranging from poor to excellent depending on soil temperature, time of germination, depth of seed in soil, amount of moisture. Control may be improved with timely cultivation.

Broadcast Rates (Pounds per Acre)

Soil Texture	SENCOR 50 WP	SURFLAN 75 WP
Medium (loam, silt loam, silt, sandy clay, sandy clay loam).	2/3	1-1/3
Heavy (silty clay, silt clay loam, ^a /clay, clay loam)	1	2
Mississippi Delta Only	1 to 1-1/4	2

^a Silty clay loam soils are transitional soils and may be classified as medium textured soils in some regions of the U.S.

Do not use on sandy loam or loamy sand soils with less than two percent organic matter. Do not apply to sandy soils.

Directions For Use

Chop and thoroughly mix crop residues into soil to a depth of 4 to 6 inches by deep plowing, offset disking or tandem disking prior to the application of the herbicide combination. Use machinery that breaks large clods prior to application.

Fill the spray tank half full of clean water and add the recommended amount of SENCOR 50 WP and Surflan 75 WP. Mix thoroughly and add water to fill spray tank. Apply in 20 to 40 gallons of water per acre (broadcast basis) using any properly calibrated low pressure herbicide sprayer equipped with herbicide tips and screens no finer than 50 mesh in nozzle and in-line strainers. Agitate thoroughly before and during application with good bypass agitation.

Sprayers should be accurately calibrated before applying SENCOR 50 WP and Surflan 75 WP. Avoid boom overlaps that will increase rates above those recommended. Check the sprayer daily during the application to be sure it is working properly and delivering a uniform spray pattern. As the amount of water used (spray volume) decreases, the importance of accurate calibration and uniform application increases.

Seed must be planted at least 1½ inches below soil surface. Apply within 2 days after planting but before crop emergence. Do not spray over top of emerged soybeans.

A one-half inch rain or more or its equivalent in overhead irrigation is needed to activate the herbicide combination. If a one-half inch rain does not occur within 7 days after application, incorporate the herbicide combination into the top one inch of the soil with a ground-driven rotary hoe, rolling cultivator or similar implement that will not damage soybean plants. Cultivation will control emerging annual weeds and grasses and improve herbicidal effectiveness. If heavy rains pack the soil surface, rotary hoe to break the crust and to aid soybean emergence.

Special Precautions

Poor weed control and/or crop injury may result if directions are not followed. Over-application may result in crop injury and/or soil residue. Do not use treated vines for feed or forage. Do not contaminate any body of water nor apply to any area not specified on the label. Do not allow sprays to drift onto adjacent desirable plants.

For additional precautions, restrictions, limitations and sprayer clean-up information, refer to the appropriate section of this label and the Surflan 75 WP label.

Note

SENCOR must be applied in strict accordance with the "DIRECTIONS FOR USE" and "RECOMMENDED APPLICATIONS" previously outlined. Applications of SENCOR made at rates above those recommended may result in burning, yellowing, or stunting of the crop. If heavy rains occur soon after application, plant injury may result, especially in poorly drained areas where water may stand for several days. Do not plant any crop other than soybeans on fields treated with SENCOR within 4 months (120 days) after treatment.

Potatoes

SENCOR 50% Wettable Powder is recommended for use in ground or aircraft spray equipment as a preemergence and/or postemergence application to potatoes (Postemergence applications may be made only on white-skinned potato varieties that are not early maturing). For optimum control of weeds in potatoes apply SENCOR 50% Wettable Powder before weeds are 1-inch tall.

Ground Application:

SENCOR 50% Wettable Powder is recommended for use with ground spray equipment, except air blast sprayers, as a preemergence and/or postemergence application for control of certain grass and broadleaf weeds in potatoes. Apply as a broadcast spray after planting and before crop emergence or after drag-off if this operation is part of the normal cultural practice. (Postemergence applications may be made only on white-skinned potato varieties that are not early maturing).

Aerial Application:

SENCOR 50% Wettable Powder is recommended for use in aerial spray equipment as a preemergence and/or postemergence application for control of certain grass and broadleaf weeds in potatoes. Apply as a broadcast spray after planting and before crop emergence or after drag-off if this operation is part of the normal cultural practice. Postemergence applications should be applied after crop emergence. (Postemergence applications may be made only on white-skinned potato varieties that are not early maturing).

Preemergence

WEEDS CONTROLLED: SENCOR 50% Wettable Powder applied preemergence to potatoes, according to directions, will effectively control:

Grass Weeds:

Crabgrass
Giant foxtail
Green foxtail
Yellow foxtail
Seedling johnsongrass
Fall panicum
Signalgrass

Broadleaf Weeds:

Carpetweed
Cocklebur
Jimsonweed
Lambsquarters
Wild mustard
Prostrate pigweed
Redroot pigweed
Common ragweed
Shepherdspurse
Sicklepod (*Cassia*)
Pennsylvania smartweed

HARD TO CONTROL WEEDS: Although SENCOR 50% Wettable Powder applied preemergence may not provide commercially acceptable control in every instance, it will suppress growth of the following weeds and reduce their competition with potato plants

Grass Weeds
Barnyardgrass
Nutsedge

Broadleaf Weeds:
Common purslane

Postemergence

WEEDS CONTROLLED: SENCOR 50% Wettable Powder applied early post-emergence to potatoes, according to directions, will effectively control

Grass Weeds:
Giant foxtail
Green foxtail

Broadleaf Weeds:
Cocklebur
Lambsquarters
Redroot pigweed
Common ragweed
Pennsylvania smartweed
Russian thistle

HARD TO CONTROL WEEDS: Although SENCOR 50% Wettable Powder applied postemergence may not provide commercially acceptable control in every instance, it will suppress growth of the following weeds and reduce their competition with potato plants

Grass Weeds:
Barnyardgrass

Broadleaf Weeds:
Kochia
Sunflower

Recommended Applications

CROP	Pounds SENCOR 50% W.P.	REMARKS
Potatoes	1 to 2	PREEMERGENCE: Apply specified dosage per acre as a broadcast spray to the soil surface after planting and before crop emergence or after drag-off if this operation is part of normal cultural practice. * Distribute the chemical evenly over the area to be treated. Do not incorporate into soil. Use the lower rate on sandy soils. See "NOTE" below before using this product on potatoes.
Potatoes (white-skinned varieties that are not early maturing)	1/2 to 1	POSTEMERGENCE: Apply specified dosage per acre as a broadcast spray over the tops of potato plants. * Use rates of 1/2 to 1 lb. for control of redroot pigweed and lambsquarters only. Apply the 1 lb. rate for control of other weeds listed on this label. Do not apply within 60 days of harvest. See "NOTE" below before using this product on potatoes. SPLIT APPLICATIONS: This product may be applied once preemergence and once post-emergence as directed above. * Do not apply post-emergence within 60 days of harvest. See "NOTE" below before using this product on potatoes.

***IMPORTANT:** Do not apply more than 2 pounds total SENCOR 50% Wettable Powder per acre in a single crop season regardless of method of application.

NOTE

Do Not Use SENCOR 50% Wettable Powder on potatoes in Kern County, California.

Postemergence applications may cause some chlorosis or minor necrosis.

Do not use SENCOR for postemergence applications on early maturing, smooth-skinned, white- or red-skinned varieties of potatoes.

Do not plant treated area to crops other than potatoes for one year after treatment. All cover crops that are planted during this one year period must be plowed down and not grazed.

Do not apply to sweet potatoes or yams.

Sensitive Crops: Do not plant sensitive crops such as onions, lettuce, corn, beans and cucurbits during the next growing season following applications of SENCOR. Injury may result.

Restrictions

Do not use on other crops grown for food or forage. Use only according to label directions. Do not contaminate any body of water by cleaning of equipment or disposal of wastes, nor apply to any area not specified on the label.

Do not allow sprays to drift on to adjacent desirable plants.

Observe all cautions and limitations on labeling of all products used in mixtures.

Container Disposal

Containers should be disposed of by burying with wastes, or by burning (keep out of smoke).

Caution

May be harmful if swallowed or inhaled. Avoid contact with eyes, skin or clothing. Avoid breathing of dust or spray mist. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before re-use.

Do not contaminate feed or food. Keep out of reach of children.

Warranty and Limitation of Damages

Chemagro warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the risks referred to therein. CHEMAGRO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY, and no agent of Chemagro is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

RESPONSEability to you and nature



CHEMAGRO AGRICULTURAL DIVISION
MOBAY CHEMICAL CORPORATION
Box 4913 Kansas City Mo 64120

LASSO® is a Reg. T.M. of Monsanto
TREFLAN® is a Reg. T.M. of Elanco Products