

3125-280

7-25-74

1081

U.S. LABEL

Reason to Issue: To increase pre-harvest interval for cauliflower to 28 days, and for cabbage to 35 days.

Date of Draft: 5/28/74 (Pre-Reg.) (S)

EPA Reg. No. 3125-280

ACCEPTED

JUL 25 1974

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. _____ SUBJECT
TO ATTACHED COMMENTS.

(R)
MONITOR 4
Liquid Insecticide

AMENDMENT
To Previously Registered Labeling

3125-280

Revise as follows:

RECOMMENDED APPLICATIONS

CROP	INSECT	DOSAGE - Pints MONITOR 4	REMARKS
Broccoli Brussels Sprouts Cabbage Cauliflower	Cabbage looper Imported Cabbage- worm Diamondback moth larvae Aphids	1 to 2	<p><u>In a Preventive Program:</u> Beginning when the plants are small, use the 1 pint rate (0.5 pound active) per acre and apply at 7-day intervals.</p> <p><u>For Clean-up of Existing Populations:</u> Use 1-1/2 to 2 pints (0.75 to 1.0 pound active) per acre. Do not apply, to Brussels sprouts later than 14 days before the first harvest, or to cauliflower later than <u>28</u> days before the first harvest, or to cabbage later than <u>35</u> days before first harvest. For broccoli, the dosage rate of 1 pint per acre may be made up to 14 days before first harvest, but do not apply rates above 1 pint per acre within 21 days of first harvest.</p>

ALL COPY EXCEPT THAT WHICH IS DOUBLE UNDERSCORED HAS BEEN PREVIOUSLY REGISTERED.