

Reason to Issue: To revise claims
for use on tobacco. 2/21/85 Draft:
Revise "Remarks" for clarity.
6/19/85 Draft: Revise based on
internal comments.

Date of Draft: 6/19/85 (Pre-Reg) (H)
Supersedes Draft Dated: 2/21/85

EPA Reg. No. 3125-283

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their
direct supervision and only for those uses covered by the Certified
Applicator's certification.

® NEMACUR 3

Emulsifiable Systemic Insecticide-Nematicide

ACTIVE INGREDIENT:

Ethyl 3-methyl-4-(methylthio)phenyl
(1-methylethyl)phosphoramidate 35%

AMENDMENT TO PREVIOUSLY REGISTERED LABELING

REVISE TO READ AS FOLLOWS:

Tobacco

(Not for Use on Shade-Grown Tobacco)

ACCEPTED

SEP 18 1985

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 3125-283

COMPATIBILITY

NEMACUR 3 is physically compatible with many registered soil insecticides, herbicides and fungicides. When tank mixing, add any wettable powders, flowables, or dry flowables to 1/2 the required amount of water and agitate. Then add emulsifiable materials one at a time to this mixture, always adding NEMACUR 3 to the mixture last with the remaining water. Agitate continuously during and after mixing procedure until mix is applied to the soil.

To determine the physical compatibility of NEMACUR 3 with other products, the following procedure should be followed if local experience is unavailable: Pour the recommended proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least five (5) minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible. For further information contact your local Mobay representative.

RECOMMENDED APPLICATIONS

CROP	PEST	DOSAGE OF NEMACUR 3	REMARKS
FIELD CROPS			
Tobacco (Not for use on shade- grown tobacco)	Nematodes	1-1/3 to 2 gallons	Uniformly broadcast the specified dosage in a minimum of 20 gallons of water per acre over the soil surface. Incorporate to a depth of 2 to 4 inches by disking or tilling. Where a range in rates is recommended, use the high rate in fields with high populations of nematodes or in fields having a history of serious nematode damage. Plant crop in the usual manner.
NEMACUR 3 can be tank mixed with fungicides, herbicides and insecticides registered and recommended for use on tobacco. Observe all cautions and limitations on labeling of all products used in tank mixtures.			