

U.S. LABEL

Reason to Issue: To add claims for
preplant application to pine-
apple (Hawaii only).

Date of Draft: 6/19/85 (Pre-Reg) (H)

EPA Reg. No. 3125-283

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their
direct supervision and only for those uses covered by the Certified
Applicator's certification.

ACCEPTED**JUN 17 1986**

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under
EPA Reg. No. 3125-283

® NEMACUR 3

Emulsifiable Systemic Insecticide-Nematicide

ACTIVE INGREDIENT:

Ethyl 3-methyl-4-(methylthio)phenyl

(1-methylethyl)phosphoramidate 35%

AMENDMENT TO PREVIOUSLY REGISTERED LABELINGRevise to Read as Follows:**RECOMMENDED APPLICATIONS**

CROP	PEST	DOSAGE OF NEMACUR 3	REMARKS
Pineapple (Hawaii only)	Nematodes (<u>Rotylenchulus</u> and <u>Meloidogyne</u> species)	<u>Preplant</u> 6 2/3 gal	<p><u>PREPLANT APPLICATION:</u></p> <p>Apply the specified dosage per acre as a water emulsion spray over the entire area to be treated. Incorporate thoroughly to a depth of 4 to 6 inches to insure uniform distribution.</p> <p>Additional postplant applications may be made to the plant crop at 1 to 3 month intervals by broadcast spray or drip irrigation equipment as indicated below. However, do not apply more than a total of 12 4/3 gallons of NEMACUR 3 (40 pounds of active ingredient) per acre per plant crop.</p>

RECOMMENDED APPLICATIONS (Continued)

CROP	PEST	DOSAGE OF NEMACUR 3	REMARKS
Pineapple (Hawaii only)	Nematodes (<u>Rotylenchulus</u> and <u>Meloidogyne</u> species)	<u>Postplant</u> 1 1/3 to 8 pints	<p><u>POSTPLANT APPLICATION:</u></p> <p>- <u>PLANT CROP.</u> Apply specified dosage in 50 to 250 gallons of water per acre as a broadcast spray or through drip irrigation. Applications may begin immediately after planting. Make additional applications at intervals of 1 to 3 months as needed. Do not apply within 30 days before harvest.</p> <p>Do not apply more than a total of 13 1/3 gallons of NEMACUR 3 (40 pounds of active ingredient) per acre per plant crop.</p> <p>-<u>RATOON CROP.</u></p> <p>Apply specified dosage in 50 to 250 gallons of water per acre as a broadcast spray or through drip irrigation. The first application may be made immediately following crop harvest. Make additional applications at intervals of 1 to 3 months as needed. Do not apply within 30 days before harvest.</p> <p>Do not apply more than a total of 6 2/3 gallons of NEMACUR 3 (20 pounds of active ingredient) per acre to each ratoon crop.</p>

Do not use green forage or green fodder (cannery waste, such as cull fruit, fruit skin or shells, crowns, cores and basal ends of fruit may be fed) for animal feed.