

U.S. LABEL

Reason to Issue: To reduce pre-harvest
interval to 30 days for all citrus fruits.

Date of Draft: 10/28/86 (Pre-Reg) (H)

EPA Reg. No. 3125-283

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons
under their direct supervision and only for those uses covered
by the Certified Applicator's certification.



NEMACUR 3

EMULSIFIABLE SYSTEMIC INSECTICIDE - NEMATOCIDE

ACTIVE INGREDIENT:

Ethyl 3-methyl-4-(methylthio)phenyl
(1-methylethyl)phosphoramidate 35%

AMENDMENT TO PREVIOUSLY REGISTERED LABELING
REVISE TO READ AS FOLLOWS:

RECOMMENDED APPLICATIONS

CROP	PEST	DOSAGE OF NEMACUR 3	REMARKS
FRUIT			
Citrus (Except Arizona, California, and Texas)	Nematodes	3-1/3 to 6-2/3 gallons	<p>BROADCAST: Apply specified dosage in not less than 20 gallons of solution per acre to the soil with ground equipment and incorporate immediately.*</p> <p>BAND: Apply proportionally less product per acre based on band width** with ground equipment and incorporate immediately.* Center the treated band on the tree row using a minimum band width of "to the drip line."</p> <p>LOW PRESSURE IRRIGATION: Apply proportionally less product per acre per season based on the area wetted.** This rate per season may be applied in a single application or multiple applications.</p>
Citrus (Arizona, California and Texas only) (In California do not apply to Kumquat, Tangelo, or Citrus hybrids)	Nematodes	6-2/3 gallons	<p>BROADCAST: Apply specified dosage in not less than 20 gallons of solution per acre to the soil with ground equipment and incorporate immediately.*</p> <p>BAND: Apply proportionally less product per acre based on band width** with ground equipment and incorporate immediately.* Center the treated band on the tree row using a minimum band width of "to the drip line".</p> <p>LOW PRESSURE IRRIGATION: Apply proportionally less product per acre per season based on the area wetted**. This rate per season may be applied in a single application or multiple applications.</p>

RECOMMENDED APPLICATIONS (Cont'd)

*Incorporate the product into the soil or supply sufficient irrigation for incorporation. Control of nematodes is best obtained when this is adequate rainfall or irrigation after application. Under dry conditions, follow with 1/2 to 1 inch of irrigation.

**Example: To treat trees planted on 20 foot centers with a 10 foot band centered on the row or 50% of the surface area wetted by low pressure irrigation, apply one-half of the broadcast rate.

Do not apply within 30 days of harvest. Do not exceed 20 pounds active ingredient (6 2/3 gallons of NEMACUR 3) per acre per season (broadcast rate). Do not graze livestock in treated areas.

On non-bearing citrus make initial applications when trees have recovered from transplanting shock and have initiated new growth, provided that one or more irrigations have been applied and soil has settled around the roots. Phytotoxicity may occur if applications are made before soil has settled and plants are growing.

U.S. LABEL

Reason to Issue: To reduce pre-harvest interval to 30 days for all citrus fruits.

Date of Draft: 10/28/86 (Pre-Reg) (H)

EPA Reg. No. 3125-283

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

® NEMACUR 3

EMULSIFIABLE SYSTEMIC INSECTICIDE - NEMATOCIDE

ACCEPTED

MAY 12 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 3125-283

ACTIVE INGREDIENT:

Ethyl 3-methyl-4-(methylthio)phenyl

(1-methylethyl)phosphoramidate 35%

AMENDMENT TO PREVIOUSLY REGISTERED LABELING
REVISE TO READ AS FOLLOWS:

RECOMMENDED APPLICATIONS

CROP	PEST	DOSAGE OF NEMACUR 3	REMARKS
FRUIT			
Citrus (Except Arizona, California, and Texas)	Nematodes	3-1/3 to 6-2/3 gallons	<p>BROADCAST: Apply specified dosage in not less than 20 gallons of solution per acre to the soil with ground equipment and incorporate immediately.*</p> <p>BAND: Apply proportionally less product per acre based on band width** with ground equipment and incorporate immediately.* Center the treated band on the tree row using a minimum band width of "to the drip line."</p> <p>LOW PRESSURE IRRIGATION: Apply proportionally less product per acre per season based on the area wetted.** This rate per season may be applied in a single application or multiple applications.</p>
Citrus (Arizona, California and Texas only) (In California do not apply to Kumquat, Tangelo, or Citrus hybrids)	Nematodes	6-2/3 gallons	<p>BROADCAST: Apply specified dosage in not less than 20 gallons of solution per acre to the soil with ground equipment and incorporate immediately.*</p> <p>BAND: Apply proportionally less product per acre based on band width** with ground equipment and incorporate immediately*. Center the treated band on the tree row using a minimum band width of "to the drip line".</p> <p>LOW PRESSURE IRRIGATION: Apply proportionally less product per acre per season based on the area wetted**. This rate per season may be applied in a single application or multiple applications.</p>

RECOMMENDED APPLICATIONS (Cont'd)

*Incorporate the product into the soil or supply sufficient irrigation for incorporation. Control of nematodes is best obtained when this is adequate rainfall or irrigation after application. Under dry conditions, follow with 1/2 to 1 inch of irrigation.

**Example: To treat trees planted on 20 foot centers with a 10 foot band centered on the row or 50% of the surface area wetted by low pressure irrigation, apply one-half of the broadcast rate.

Do not apply within 30 days of harvest. Do not exceed 20 pounds active ingredient (6 2/3 gallons of NEMACUR 3) per acre per season (broadcast rate). Do not graze livestock in treated areas.

On non-bearing citrus make initial applications when trees have recovered from transplanting shock and have initiated new growth, provided that one or more irrigations have been applied and soil has settled around the roots. Phytotoxicity may occur if applications are made before soil has settled and plants are growing.

YLD12

Base Pre-Reg (6026)

U.S. LABEL

Reason to Issue: To propose use for
mole cricket control; clarify
irrigation statement; add nozzle
system restriction.

Date of Draft: 10/30/86 (Pre-Reg) (3)

EPA Reg. No. 3125-283

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or
persons under their direct supervision and only for those
uses covered by the Certified Applicator's certification

® NEMACUR 3
TURF NEMATOCIDE

ACTIVE INGREDIENTS:

Ethyl 3-methyl-4-(methylthio)phenyl
(1-methylethyl)phosphoramidate 35%

AMENDMENT
To Previously Registered Labeling

Revise as Shown:

RECOMMENDED APPLICATIONS

ACCEPTED
MAY 12 1987
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, this pesticide is registered under EPA Reg. No. 3125-283.

CROP	PEST	DOSAGE NEMACUR 3 TURF NEMATOCIDE	REMARKS
Turf Grasses FOR USE ON GOLF COURSES	Nematodes <u>Mole</u> <u>Crickets</u>	9 to 12 fl ozs per 1000 square feet or 3 to 4 gal/acre	Apply specified dosage in a minimum of 1/2 gallon of water per 1,000 sq. ft. (approximately 20 gallons per acre) to provide uniform distribution. Do not use Boomjet® nozzle or similar single nozzle system that dispenses spray into the air over a wide swath. Irrigate immediately after treatment applying a minimum of 1/2 inch of water within 6 hours of application. Do not treat newly seeded areas until the plants have developed secondary root systems. Do not apply more than twice per year. Do not use on residential lawns or public recreational areas other than golf courses. Not recommended for use on tees and greens.

Boomjet - Registered trademark of Spray System Company

All copy except that which is underlined has been previously registered.