SUPPLEMENTAL

SENCOR 70% Wettable Powder EPA Reg. No 3125-294

SUGARCANE (Louisiana and Texas Only)

Preemergence and postemergence applications of SENCOR 70% Wettable Powder with aerial recommended for control of the following weeds in sugarcane in Louisiana and Texas:

Grass Weeds

Crabgrass **Fox tails** Seedling johnsongrass Signalgrass (Brachiaria)

Broadleaf Weeds

Soinv amaranth Field bindweed Chickweed Curly dock Henbit Lambsquarters London rocket

Marestail Wild mustard Winter oats Pigweeds Common purslane Sowthistle

RECOMMENDED APPLICATIONS

Pounds SENCOR	REMARKS
70% WP	
2-% to 4-%	BROADCAST: Apply specified dosage per acre using 20 to 30 gallons of water with ground equipment or 5 gallons of water with aircraft spray equipment. Apply as a broadcast spray during the Fall after planting or to the stubble after harvest. Make a second application in early Spring prior to green-up. Use the higher rate on heavy clay soil and on soil with a high percentage of organic matter. If necessary a third application may be made in late Spring at layby. Do not apply within 60 days of harvest.
1-1/8 to 2-%	BAND:Apply specified dosage in 10 to 20 gallons of water per acre in a 30 to 36 inch band over the row during the Fall after planting or to the stubble after harvest. Make a second application in early Spring prior to green-up. Use the higher rate on heavy clay soil and on soil with a high percentage of organic matter. If necessary a third application may be made in late Spring at layby. Do not apply within 60 days of harvest.

Do not use treated foliage for feed or forage.

IMPORTANT: This supplemental labeling is not intended to provide full information for use of this product. Before using SENCCR 70% Wettable Powder, read and carefully observe caution statements, directions for use, restrictions, and disposal information on the label.

> MOBAY CHEMICAL CORPORATION CHEMAGRO AGRICULTURAL DIVISION Kansas City, Missouri 64120