



NET CONTENTS POUNDS

# DEXON D

## 72% WETTABLE POWDER

COTTONSEED TREATMENT FUNGICIDE

Contains 32% DEXON plus 40% DACONIL 2787<sup>†</sup>  
**FOR CONTROL OF PLANT DISEASES**

**STOP READ THE LABEL BEFORE USE**  
EPA Reg. No. 3125-298

DEXON is a Reg. TM of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen  
DACONIL 2787 is a registered trademark of Diamond Shamrock

ACTIVE INGREDIENTS	
Sodium [4-(dimethylamino)phenyl] diazinesulfonate †	32%
Chlorothalonil (Tetrachloro isophthalonitrile) ††	40%
INERT INGREDIENTS	28%
	<hr/> 100%
†U.S. Patent No. Re. 24,960	
††U.S. Patent No. 3,290,353 and No. 3,331,735	

### DIRECTIONS FOR USE

**IMPORTANT** Read these entire Directions and Conditions of Sale, including the Warranty and Limitation of Damages provision, before using DEXON D 72% Wettable Powder

**CONDITIONS OF SALE** These directions for Use reflect the opinion of experts based on tests of effectiveness, of toxicity to plants and laboratory animals and of residues on or in food and feed under normal conditions of use. However, because of the wide range of conditions under which this product may be used, even though label directions are followed, it is impossible to eliminate all risks associated with its use because of abnormal conditions beyond the control of the seller, Chemagro and the Seller offer, and the Buyer uses, this product subject to the understanding that the Buyer assumes all such risks.

DEXON D 72% Wettable Powder is recommended for treatment of cottonseed and in protection of germinating seeds and developing seedlings from disease

infecting organisms such as *Rhizoctonia* and *Pythium* species.

Use suitable equipment to properly apply the prescribed amount of DEXON D 72% Wettable Powder on cottonseed. Any screen or strainer in equipment should be no finer than 50 mesh to prevent clogging. Premix the required amount of DEXON D 72% Wettable Powder in a clean container with a small amount of water and add to the tank as it is being filled. Maintain agitation during filling and application. Do not allow diluted product to remain in equipment overnight. Rinse out equipment after use and properly dispose of wastes by burying in non-crop area away from water supplies.

**SEED TREATMENT** Mix DEXON D 72% Wettable Powder with an appropriate amount of water (normally about 1/2 to 1 percent of seed weight) to apply at rate of 2 ounces of DEXON D 72% Wettable Powder per 100 pounds of acid delinted cottonseed, or 3 ounces per 100 pounds of machine delinted cottonseed. See chart.

### RECOMMENDED APPLICATIONS

CROP	DISEASE	Ounces DEXON D 72% W.P.	REMARKS
Cotton	Pythium Rhizoctonia	2 to 3	Apply specified dosage to 100 pounds of seed as a slurry treatment in sufficient water to provide uniform coating of the seed

**NOTE** To meet U.S. Federal Seed Act requirements, all seed treated with DEXON D 72% Wettable Powder should be labeled "Treated with sodium [4-(dimethylamino)phenyl] diazinesulfonate (DEXON) and Chlorothalonil (Tetrachloroisophthalonitrile)"

### SEED TREATMENT CHART

Condition of Seed	Rate/lwt seed	Formulation gal of Slurry	Pounds Seed Treated per gal of Slurry
Acid delinted	2 to 3	1 to 1 1/2	1650
Machine delinted	3 to 4	1 to 1 1/2	820

Treated seed must not be used for food, food or feed purposes, or in any application other than that for which it is intended. Applications at rates above those shown may result in equipment clogging.

This product is toxic to fish and wildlife. Keep out of any body of water. Treated seeds, spilled slurry, or surface may be hazardous to birds and other wildlife. If spilled, use treated seeds by burying.

### CONTAINER DISPOSAL

Use equipment and dispose of waste by burying in non-crop area away from water supplies. Dispose of container by burying with water inside.

**DANGER KEEP OUT OF REACH OF CHILDREN**

Contains 32% DEXON plus 40% DACONIL 2787

FOR CONTROL OF PLANT DISEASES

STOP - READ THE LABEL BEFORE USE

EPA Reg. No. 3125-298

DEXON is a Reg. TM of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen.

DACONIL 2787 is a registered trademark of Diamond Shamrock.

DIRECTIONS FOR USE

**IMPORTANT:** Read these entire Directions and Conditions of Sale, including the Warranty and Limitation of Damages provision, before using DEXON D 72% Wettable Powder.

**CONDITIONS OF SALE:** These directions for Use reflect the opinion of experts based on tests of effectiveness, of toxicity to plants and laboratory animals and of residues on or in food and feed under normal conditions of use. However, because of the wide range of conditions under which this product may be used, even though label directions are followed, it is impossible to eliminate all risks associated with its use because of abnormal conditions beyond the control of the seller, Chemagro and the Seller offer, and the Buyer uses, this product subject to the understanding that the Buyer assumes all such risks.

DEXON D 72% Wettable Powder is recommended for treatment of cottonseed and in protection of germinating seeds and developing seedlings from disease

ACTIVE INGREDIENTS:

Sodium [4-(dimethylamino)phenyl]diazene-sulfonate 1 ..... 32%  
 Chlorothalonil (Tetrachloro-isophthalonitrile) 11 ..... 40%

INERT INGREDIENTS

..... 28%  
 -----  
 100%  
 U.S. Patent No. Re. 24,960  
 11 U.S. Patent No. 3,290,353 and No. 3,331,735

including organisms such as *Rhizoctonia* and *Pythium* species.

Use suitable equipment to properly apply the prescribed amount of DEXON D 72% Wettable Powder on cottonseed. Any screen or strainer in equipment should be no finer than 50 mesh to prevent clogging. Premix the required amount of DEXON D 72% Wettable Powder in a clean container with a small amount of water and add to the tank as it is being filled. Maintain agitation during filling and application. Do not allow diluted product to remain in equipment overnight. Rinse out equipment after use and properly dispose of wastes by burying in non-crop area away from water supplies.

**SEED TREATMENT:** Mix DEXON D 72% Wettable Powder with an appropriate amount of water (normally about 1/2 to 1 percent of seed weight) to apply at rate of 2 ounces of DEXON D 72% Wettable Powder per 100 pounds of acid delinted cottonseed, or 3 ounces per 100 pounds of machine delinted cottonseed. See chart.

RECOMMENDED APPLICATIONS

CROP	DISEASE	Ounces DEXON D 72% W.P.	REMARKS
Cotton	Pythium Rhizoctonia	2 to 3	Apply specified dosage to 100 pounds of seed as a slurry treatment in sufficient water to provide uniform coating of the seed

**NOTE:** To meet U.S. Federal Seed Act requirements, all seed treated with DEXON D 72% Wettable Powder should be labeled "Treated with Sodium [4-(dimethylamino)phenyl]diazene-sulfonate (DEXON) and Chlorothalonil (Tetrachloroisophthalonitrile)"

SEED TREATMENT CHART

Cottonseed	Rate (wt. seed)	Amount Formulation (gal. of Slurry)	Pounds Seed Treated per gal. of Slurry
Hand delinted	2 oz.	2 lb.	1650
Machine delinted	3 oz.	1 lb.	820

RESTRICTIONS

Treated seed must not be used for feed, food, or oil purposes. Use only according to label directions. Applications at rates above those shown may be illegal for residues.

This product is toxic to fish and wildlife. Keep out of any body of water. Treated seeds exposed on soil surface may be hazardous to birds and other wildlife. Dispose of excess treated seeds by burying.

CONTAINER AND WASTE DISPOSAL

Rinse equipment and dispose of waste by burying in non-crop lands away from water supplies. Dispose of container by burying with wastes or by

burning (keep out of smoke). Never reuse container.

**DANGER KEEP OUT OF REACH OF CHILDREN**

Causes severe eye damage. May be fatal if swallowed or inhaled. Do not breathe dust or spray mist.

Contaminated clothing with soap and hot water before reuse.

Do not get in eyes or on skin. In case of contact with eyes, flush with plenty of water for at least 15 minutes. Get immediate medical attention. Wear natural rubber gloves, a mask suitable for use against toxic dusts and sprays, and goggles when treating seed. Wash hands, arms and face thoroughly with soap and warm water after use and before eating or smoking. Wash all

**NOTE:** DEXON D 72% Wettable Powder may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician. Affected persons respond to treatment with anti-histamines or steroid creams and/or systemic steroids.

WARRANTY AND LIMITATION OF DAMAGES

Chemagro warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the risks referred to therein. CHEMAGRO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, GUARANTEE, OR OTHER EXPRESS OR IMPLIED WARRANTY.

**FITNESS OR MERCHANTABILITY,** and no agent of Chemagro is authorized to do so except in writing with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages, such as lost profits or values, et

**CHEMAGRO**

Division of Bayer Corporation  
 Box 4913, Kansas City, Mo. 64120