



SUPPLEMENTAL LABELING

Möbax Chemical Corporation
Agricultural Chemicals Division
Box 4913, Kansas City, MO 64120

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

DI-SYSTON 8
Emulsifiable Systemic Insecticide

EPA Reg. No. 3125-307

RECOMMENDED APPLICATIONS

CROP	INSECT	DOSAGE DI-SYSTON 8	REMARKS
Wheat (Fall)	Hessian fly Aphids (Oat bird cherry)	0.25 fl. ounces per 1000 ft. of row (see chart below) or up to 1 pint per acre	Apply as a soil injection in water emulsion or with liquid fertilizer at planting time in the fall.
		Ounces DI-SYSTON 8 per 1000 ft. of row based on 3 inch band	
		Row Spacing	Ounces/Acre
		7 or 8 inch rows	15.5 .25
		10 inch rows	13.25 .25
		12 inch rows	10.75 .25
		14 inch rows	9.5 .25
		16 inch rows	8.5 .25
		18 inch rows	7.0 .25

IMPORTANT: This supplemental labeling is not intended to provide full information for use of this product. Before using DI-SYSTON 8, read and carefully observe the pre-cautionary statements, directions for use, restrictions, storage and disposal statements, and other pertinent information on the label.

ACCEPTED

JUN - 7 1982

U.S. Department of Agriculture
Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 3125-307



SUPPLEMENTAL LABELING


Möbry Chemical Corporation
Agricultural Chemicals Division
Box 4913, Kansas City, MO 64120

July 13, 1981

DI-SYSTON 8
INSECTICIDE

EPA Reg. No. 3125-307

RECOMMENDED APPLICATIONS



CROP	INSECT	DOSAGE - DI-SYSTON 8	REMARKS
Wheat (Foliar Appli- cation Spring or Fall)	Aphids (Greenbugs) Mites (See NOTE in "REMARKS")	4 to 12 fluid ounces per acre	For application by aircraft or ground equipment: Apply specified dosage per acre in sufficient water, to obtain thorough coverage but not less than 1 gallon total volume per acre. Use the lower rates on plants up to tillering (stooling). Higher rates should be used after plants begin to tiller. For application in liquid fertilizer: Mix 6 to 12 fluid ounces of DI-SYSTON 8 in the amount of liquid fertilizer to be applied per acre. Apply this mixture as a top dressing with suitable ground equipment. Fall Applications: Two applications may be made in the fall. Allow 30 days between applications. Spring Applications: Application at green-up may be made following "fall applications." A second application may be made 30 days following the first spring application. Do not make any applications within 30 days of grain harvest. NOTE (1) Do not graze treated fields after any application of DI-SYSTON 8. (2) Due to greenbug resistance in certain areas, DI-SYSTON used alone may not provide satisfactory control. Consult your local agricultural advisor or Extension Service for recommendations.

IMPORTANT: This supplemental labeling is not intended to provide full information for use of this product. Before using DI-SYSTON 8 read and carefully observe caution statements, directions for use, restrictions and disposal information on the label.