

Pm 13

3125-307

Page 1/1 of 23



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

FEB 10 1994

Karen M. Pither
MILES, INC
Agricultural Division
8400 Hawthorn Road
Box 4913
Kansas City, MO 64120

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Subject: Label Amendment Submission of 7/1/93 in Response to PR Notice 93-7
REISSUANCE OF ACCEPTANCE LETTER
EPA Reg. No. 3125-307
DI-SYSTON 8

Dear Registrant:

This letter replaces the letter sent to you on January 31, 1993. The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments listed on the enclosed sheet.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.



Recycled/Recyclable
Printed with Soy/Canola Ink on paper that
contains at least 50% recycled fiber

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
 Office of Pesticide Programs (7505C)
 U.S. Environmental Protection Agency
 401 M Street, SW
 Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
 Office of Pesticide Programs
 Room 266A, Crystal Mall 2
 1921 Jefferson Davis Highway
 Arlington, VA 22202

Sincerely,


 Jim Tompkins, Deputy Chief
 Registration Support Branch
 Registration Division (7505W)

Attachment

MILES, INC
Agricultural Division
Kansas City, MO

EPA Reg. No. 3125-307
DI SYSTON 8

Comments:

Move the following statement to within the Agricultural Use Requirements Box: "Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas." Refer to the Worker Protection Standard Worksheet, Part 2, Section F, item 6 for the precise location.

User Safety Recommendations must either be placed in a box or printed on the label in a contrasting color from surrounding text.

The Agricultural Use Requirements section must either be placed in a box or printed on the label in a contrasting color from surrounding text.

Correct the typographical errors in the PPE statements under the heading "Applicators and other handlers must wear:". The words "exposure" should be "exposures".



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

JAN 31 1994

4 July 23
OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Karen M. Pither
MILES INC.
AGRICULTURE DIVISION
8400 HAWTHORN RD BOX 4913
KANSAS CITY, MO 64120

Subject: Label Amendment Submission of 07/01/93 in Response to PR Notice 93-7
EPA Reg. No. 3125-307
DI SYSTON 8

IMPORTANT NOTICE: This Letter Requires You to Submit a Notification to the Office of Compliance Monitoring within 20 Days.

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments listed on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

SERIOUS ERROR(S) ON SUBMITTED LABELING:

EPA has determined that one or more of the errors on the labeling you submitted to the Agency is categorized as "serious" as defined in PR Notice 93-11. In general, serious errors are those that may create a potential for harm to workers, handlers, other persons, or the environment; or those that prevent the achievement of the basic goals of the Worker Protection Standard (WPS) or FIFRA.

YOU MUST NOT SELL OR DISTRIBUTE (INCLUDING RELEASE FOR SHIPMENT) ANY PRODUCT BEARING THE SUBMITTED LABELING. Sale or distribution of any product with the incorrect labeling is a violation of FIFRA that could result in initiation of an action for civil penalties and/or cancellation of your product. This matter has been referred to the Office of Compliance Monitoring (OCM) for further action.

YOU MUST SUBMIT A NOTIFICATION TO THE OFFICE OF COMPLIANCE MONITORING WITHIN 20 DAYS OF THE DATE OF THIS LETTER.



Recycled/Recyclable
Printed with SoyCandis Ink on paper that
contains at least 50% recycled fiber

Each notification must include the following information:

1. Registrant's name and address.
2. EPA Registration number of the product for which the labeling was submitted.
3. Whether any product already has been sold or distributed (this includes released for shipment) bearing the submitted labeling.

In addition, the notification must include the following information, IF you have already sold or distributed any product bearing the submitted labeling:

1. Quantity of the product that has been sold or distributed.
2. EPA Producing Establishment number.
3. Location of the product, if known.
4. Proposed plan for assuring that product with the incorrect labeling in your control, as well as that in the hands of dealers/distributors/users, will be corrected. The plan must include your proposal for assuring that the labeling is corrected as soon as possible.

Submit the notification to:

Michael Wood, Director
Compliance Division, (7203)
Worker Protection Notification
Office of Compliance Monitoring
U.S. Environmental Protection Agency
401 M Street SW
Washington, DC 20460

Failure to submit a notification also may result in initiation of an action under FIFRA for civil penalties, if you sold or distributed (this includes released for shipment) product bearing the incorrect labeling.

IN ADDITION, YOU SHOULD CONTACT IMMEDIATELY THE PERSONS TO WHOM YOU SOLD OR DISTRIBUTED YOUR PRODUCT AND INFORM THEM NOT TO SELL OR DISTRIBUTE ANY PRODUCT WITH THE INCORRECT LABELING.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the WPS labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have neither been reviewed nor accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

Submit the required notification. Then by the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling as amended

AND

- WITHIN one year from date of this acceptance.


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
 Office of Pesticide Programs (7505C)
 U.S. Environmental Protection Agency
 401 M Street, SW
 Washington, D.C. 20460-0001

Hand or courier deliveries of final labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

cc: Office of Compliance Monitoring

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division

Karen M. Pither
MILES INC
AGRICULTURE DIVISION
8400 HAWTHORN RD BOX 4913
KANSAS CITY MO 64120

Subject: Label Amendment Submission of 07/01/93
to Comply with WPS Labeling Requirements
EPA Reg Nr. 3125-307
DI - SYSTON 8

The following specific comments pertain to your
WPS labeling submission concerning the product
cited above:

User Safety Recommendations must either be placed in a box
or printed on the label in a contrasting color from
surrounding text.

Add the following statement to the Agricultural Use
Requirements box on your labeling: "Notify workers of the
application by warning them orally and by posting warning
signs at entrances to treated areas." The exact location is
shown in Section F on Part 2 of the Product Worksheet in
Supplement Three-A to PR Notice 93-7.

There is a typographical error(s) in the PPE statements that
you added to the Precautionary Statements.

DI-SYSTON 8 Emulsifiable Systemic

307-7888.YLD

U.S. LABEL

9 7 23
Base Pre-Reg. (7888)

Reason to Issue: To revise in accordance to Worker Protection Standard, PR Notice 93-7.

Date of Draft: 06/07/93(Pre-Reg)(P)

RESTRICTED USE PESTICIDE
Due to acute human oral, dermal, and inhalation toxicity
For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

DI-SYSTON® 8

EMULSIFIABLE SYSTEMIC INSECTICIDE
8 POUND PER GALLON FORMULATION
LOW ODOR FORMULA

FORMULATED FOR EFFECTIVE SOIL TREATMENT

ACTIVE INGREDIENT:

Disulfoton, O,O-Diethyl S-[2-(ethylthio)ethyl] phosphorodithioate	85%
INERT INGREDIENTS:	15%
	100%

Contains 8 lbs. O,O-Diethyl S-[2-(ethylthio)ethyl] phosphorodithioate per gallon.

EPA Reg. No. 3125-307
EPA Est. 3125-MO-1

NET CONTENTS: ___ GALLONS

STOP - READ THE LABEL BEFORE USE

KEEP OUT OF REACH OF CHILDREN

DANGER PELIGRO



POISON

(See rear panel for Statements of Practical Treatment and Other Precautionary Statements).

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

~~PRECAUCION AL USUARIO: Si usted no puede leer o entender inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.~~

~~(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)~~

MILES

ACCEPTED
with COMMENTS
in EPA Letter Dated

JAN 31 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

3125-307

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER



POISON

Poisonous if swallowed, inhaled, or absorbed through the skin. Rapidly absorbed through the skin. Do not get in eyes, on skin, or on clothing. Do not breathe fumes or spray mist. ~~Wear protective clothing, natural rubber gloves, and goggles.~~

~~Wear a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR Part II.~~

~~Do not contaminate feed or food. Keep out of reach of children. Keep all unprotected persons out of the operating area or vicinity. Wash hands, arms, and face thoroughly with soap and warm water before eating or smoking.~~

~~**HANDLING OF CONTAMINATED ARTICLES:** Clothing and rags receiving undiluted DI-SYSTON 8 as a result of leaks, spills, or mishaps should be removed immediately and not reused. These contaminated articles should be buried in the same manner as the empty container; see container disposal section for directions. (Contaminated articles which should be discarded would include: hats, gloves, aprons, coats, boots, etc. not made of rubber or coated with rubber or other similar materials). Clothing receiving spray mist or droplets from the mixed, diluted DI-SYSTON 8 should be removed and decontaminated before reuse. This clothing can be decontaminated by machine washing separately from other items with soap or detergent and bleach in hot water twice before reuse. Protective clothing and equipment should be washed down with detergent or soap and bleach in water. Wash water from the cleaning of protective clothing or equipment should not be allowed to run off or otherwise enter water supplies.~~

~~**Personal Protective Equipment:** Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category F on an EPA chemical-resistance category selection chart.~~

~~**Applicators and other handlers must wear:**~~

- ~~• Coveralls over long-sleeved shirt and long pants~~
- ~~• Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or viton~~
- ~~• Chemical-resistant footwear plus socks~~

- ~~• Protective eyewear~~

- ~~• Chemical-resistant headgear for overhead exposure~~

- ~~• Chemical-resistant apron when cleaning equipment, mixing or loading~~

- ~~• For exposure in enclosed areas, a respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G)~~

- ~~• For exposure outdoors, dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C)~~

~~Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.~~

~~**Engineering controls statements:** When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.~~

~~**User Safety Recommendations:**~~

~~**User should:**~~

- ~~• Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.~~
- ~~• Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.~~
- ~~• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.~~

~~**SYMPTOMS OF POISONING:** A sense of "tightness" in the chest. Sweating. Contracted pupils. Stomach pains. Vomiting and diarrhea.~~

STATEMENTS OF PRACTICAL TREATMENT**Organophosphate**

In case of poisoning, call a physician immediately. Have patient lie down and keep quiet. If swallowed, vomiting should be induced. Administer water freely and induce vomiting by giving one dose (1/2 oz. or 15 ml.) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Never give anything by mouth to an unconscious person. Professional medical assistance should be secured immediately. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. Get medical attention immediately. If in eyes, wash with plenty of flowing water. Get medical attention immediately.

NOTE TO PHYSICIAN: ANTIDOTE - Administer atropine sulfate in large therapeutic doses. Repeat as necessary to the point of tolerance. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Compound inhibits cholinesterase resulting in stimulation of the central nervous system, the parasympathetic nervous system, and the somatic motor nerves. Do not give morphine. Watch for pulmonary edema, which may develop in serious cases of poisoning even after 12 hours. At first sign of pulmonary edema, the patient should be placed in an oxygen tent and treated symptomatically.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to fish in neighboring areas. Do not contaminate water by disposing of equipment washwaters. Cover or incorporate spills.

This pesticide is toxic to bees exposed to direct application. Applications should be timed to coincide with periods of minimum bee activity, usually between late evening and early morning.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

RESTRICTIONS

Not for home garden use. Do not use on other crops used for food or forage. Use only according to label directions.

Application at rates above those shown may result in illegal crop residues. Do not treat food crops grown in the greenhouse. Do not apply in any manner not specified on the label.

~~RE-ENTRY STATEMENTS FOR FARM WORKERS~~

~~Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not allow reentry into treated fields within 24 hours of application unless appropriate protective clothing is worn. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.~~

~~(The following may appear on supplemental labeling accompanying the product). Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings should include these statements-~~

~~found under the label heading "Precautionary Statements" and "Re-entry Statements" included on this label. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: DANGER, area treated with DI-SYSTON 8 on (date of application). Do not allow reentry into treated fields within 24 hours of application unless appropriate protective clothing is worn. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash with plenty of flowing water. Get medical attention. If swallowed, vomiting should be induced. (See Precautionary Statements on label).~~

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

The REI is 72 hours in outdoor areas where average rainfall is less than 25 inches a year.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or butyl rubber, or nitrile rubber or viton
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

IMPORTANT: Read these entire Directions and Conditions of Sale before using DI-SYSTON 8 emulsifiable systemic insecticide.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE THE DIRECTIONS FOR CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED FOR A RANGE OF WEATHER CONDITIONS SIMILAR TO THOSE WEATHER CONDITIONS THAT ARE ORDINARY AND CUSTOMARY IN THE GEOGRAPHIC AREA WHERE THE PRODUCT IS USED. INSUFFICIENT CONTROL OF PESTS AND/OR INJURY TO THE CROP TO WHICH THE PRODUCT IS APPLIED MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL WEATHER, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO OTHER CROPS, ANIMALS, MAN, OR THE ENVIRONMENT. MILES OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL WEATHER, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF MILES AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

DI-SYSTON systemic insecticide enters the plant by absorption through the roots. The insecticide placed near the roots is translocated to all parts of the plant and kills sucking insects feeding on the plant.

The formulation of DI-SYSTON 8 is highly versatile and offers many forms of application. It can be sprayed or injected into the furrow at planting time or soil injected as a side-dress treatment at planting time or after plants become established. Applications may be made in water emulsion, with liquid fertilizer, or undiluted in specially designed equipment for low volume soil injection but never on crops for which less than 1 quart DI-SYSTON 8 per acre is the recommended dosage. When mixing with certain liquid high phosphorous fertilizers, the use of COMPLEX® compatibility agent or other equivalent agents at the rate of 1 quart per 100 gal. of mixture may be needed to assure proper mixing. Time and method of application and limits for use are discussed in detail for each specific crop under "Remarks" below.

RECOMMENDED APPLICATIONS

IMPORTANT: When used on cotton, dry beans, tomatoes, or lettuce at the maximum recommended rates under adverse conditions such as extremely cool or wet or extremely dry weather, DI-SYSTON may cause some delay in emergence, stunting of seedlings, or reduction of stand. Such damage may be more pronounced in light sandy soils. Plant injury may also occur or may be enhanced when herbicides are used in conjunction with DI-SYSTON. Refer to herbicide labels for special precautions.

DIRECTIONS FOR USE (Continued)**USE IN CHEMIGATION SYSTEMS**

DI-SYSTON 8 may be applied through irrigation systems only on those crops listed under Recommended Applications where application through irrigation systems is recommended.

Types of Irrigation Systems: Apply DI-SYSTON 8 only through sprinkler, including center pivot, lateral move, side roll, or overhead solid set irrigation systems. Do not apply DI-SYSTON 8 through any other type of irrigation system.

GENERAL DIRECTIONS FOR ALL RECOMMENDED TYPES OF IRRIGATION SYSTEMS

Uniform Water Distribution and System Calibration: The irrigation system must provide uniform distribution of treated water. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Chemigation Monitoring: A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Drift: Do not apply when wind speed favors drift beyond the area intended for treatment.

Required System Safety Devices: The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Using Water from Public Water Systems: DO NOT APPLY DI-SYSTON 8 THROUGH ANY IRRIGATION SYSTEM PHYSICALLY CONNECTED TO A PUBLIC WATER SYSTEM. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

DI-SYSTON 8 may be applied through sprinkler irrigation systems which may be supplied by a public water system only if the water from the public water system is discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and to top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. Before beginning chemigation, always make sure that the air gap exists and that there is no blockage of the overflow of the reservoir tank.

Any irrigation system using water supplied from a public water system must also meet the following requirements.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump, when the water pressure decreases to the point where the pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

DIRECTIONS FOR USE (Continued)

Posting Requirements: This sign is in addition to any sign posted to comply with the Worker Protection Standard. Posting of areas to be chemigated is required when 1) any part of a treated area is within 300 feet of sensitive areas such as residential areas, labor camps, businesses, day care centers, hospitals, in-patient clinics, nursing homes or any public areas such as schools, parks, playgrounds, or other public facilities not including public roads, or 2) when the chemigated area is open to the public such as golf courses or retail greenhouses.

Posting must conform to the follow requirements: Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive areas. When there are no usual points of entry, signs must be posted in the corners of the treated areas and in any other location affording maximum visibility to sensitive areas. The printed side of the sign should face away from the treated area towards the sensitive area. The signs shall be printed in English. Signs must be posted prior to application and must remain posted until foliage has dried and soil surface water has disappeared. Signs may remain in place indefinitely as long as they are composed of materials to prevent deterioration and maintain legibility for the duration of the posting period.

All words shall consist of letters at least 2 1/2 inches tall, and all letters and the symbol shall be a color which sharply contrasts with their immediate background. At the top of the sign shall be the words KEEP OUT, followed by an octagonal stop sign symbol at least 8 inches in diameter containing the word STOP. Below the symbol shall be the words PESTICIDES IN IRRIGATION WATER.

Posting required for chemigation does not replace other posting and reentry interval requirements for farm worker safety.

Agitation: For application of DI-SYSTON 8 alone, a chemical supply tank is not necessary for premixing since DI-SYSTON 8 mixes well with water in the irrigation line. If a chemical supply tank is used, constant strong mechanical or hydraulic agitation must be maintained in the chemical supply tank during the entire period of application.

Chemical Supply Tank Dilution: If a chemical supply tank is used, you must determine the required amounts of DI-SYSTON 8 and water to mix in the tank.

The amount of DI-SYSTON 8 needed equals the number of pints of DI-SYSTON 8 to be applied per acre multiplied by the number of acres to be chemigated.

The amount of emulsion needed equals the gallons of emulsion delivered per hour by the injection pump multiplied by the number of hours chemigation will take place.

The amount of water needed equals the amount of emulsion needed minus the amount of DI-SYSTON 8 needed.

For example, for center-pivot or automatic-move linear systems, if you want to apply 3 pints of DI-SYSTON 8 per acre to 130 acres in 20 hours and your injection pump delivers 15 gallons per hour, you need: 3 pints of DI-SYSTON 8 per acre X 130 acres = 390 pints or 48.75 gallons of DI-SYSTON 8. And, you need: 15 gallons per hour X 20 hours = 300 gallons of emulsion, minus 48.75 gallons of DI-SYSTON 8 = 251.25 gallons of water.

Another example, for solid set and manually controlled linear systems, is if you want to apply 3 pints of DI-SYSTON 8 per acre during the last 60 minutes of a set irrigating 6 acres and your injection pump delivers 20 gallons per hour, you need 3 pints of DI-SYSTON 8 per acre X 6 acres = 18 pints or 2.25 gallons of DI-SYSTON 8 for the 6 acre set. And, you need 20 gallons per hour X 1 hour = 20 gallons of emulsion, minus 2.25 gallons of DI-SYSTON 8 = 17.25 gallons of water for the 6 acre set.

Cleaning the Chemical Injection System: In order to accurately apply pesticides, the chemical injection system must be kept clean; free of chemical or fertilizer residues and sediments. Refer to your owners manual or ask your equipment supplier for the cleaning procedure for your injection system.

Flushing the Irrigation System: At the end of the application period, allow time for all lines to flush the pesticide through all nozzles or emitters before turning off irrigation water. To ensure the lines are flushed and free of pesticides, a dye indicator may be injected into the lines to mark the end of the application period.

EQUIPMENT AREA CONTAMINATION PREVENTION

It is recommended that nozzles in the immediate area of control panels, chemical supply tanks, pumps and system safety devices be plugged to prevent chemical contamination of these areas.

Center-Pivot and Automatic-Move Linear Systems: Inject the specified dosage per acre continuously for one complete revolution or move of the system. DO NOT USE END GUNS. The system should be run at maximum speed.

Solid Set and Manually Controlled Linear Systems: Injection should be during the last 30 to 60 minutes of regular irrigation period or as a separate 30 to 60 minute application not associated with a regular irrigation. DO NOT USE END GUNS.

RECOMMENDED APPLICATIONS			
CROP	INSECT	DOSAGE DI-SYSTON 8	REMARKS
Barley	Aphids Mites	1 pint per acre	Apply as a soil injection in a water emulsion or with liquid fertilizer at planting time. Do not apply within 60 days of harvest. Do not graze or cut for forage within 30 days of treatment. Do not graze treated fields after any foliar application of DI-SYSTON 8.
Barley (Foliar Application Spring or Fall)	Aphids (Including Greenbugs) (See NOTE in Remarks)	1/2 to 1 pint per acre	For application by aircraft or ground equipment: Apply specified dosage per acre in sufficient water to obtain thorough coverage but not less than 1 gallon total volume per acre. Use the lower rates on plants up to tillering (stooling). Higher rates should be used after plants begin to tiller. For application in liquid fertilizer: Mix 1/2 to 1 pint of DI-SYSTON 8 in the amount of liquid fertilizer to be applied per acre. Apply this mixture as a top dressing with suitable ground equipment. Foliar applications may be made, following an at-planting application, however, regardless of method of application, do not apply more than a total of 2 pints per acre of DI-SYSTON 8 per crop season. Do not make any applications within 30 days of grain harvest. NOTE: (1) Do not graze treated fields after any application of DI-SYSTON 8. (2) Due to greenbug resistance in certain areas, DI-SYSTON used alone may not provide satisfactory control. Consult your local agricultural advisor or Extension Service for recommendations.
Corn Field Popcorn	Aphids Leafhoppers Mites Thrips	1.2 fl oz per 1000 feet of row for any row spacing* (for 40 inch row spacing use 1 pint per acre)	TREATMENT AT PLANTING TIME: As a water emulsion, spray in 4- to 6-inch band directly behind the planter shoe in front of the press wheel or apply with liquid fertilizer in split bands on each side of the corn row. Do not apply directly to the seed. Do not forage or cut treated corn for food, feed or ensilage within 28 days of treatment. POST-PLANT SIDE-DRESS TREATMENT: An alternative treatment for early season control of aphids, leafhoppers, mites, and thrips, may be made in a water emulsion or with liquid fertilizer as a soil injection on each side of the furrow after emergence. Do not forage or cut treated corn for food, feed, or ensilage within 28 days of treatment.
	Aphids Mites	8 to 16 fl oz NOTE: Apply only one soil treatment and one foliar treatment per crop season.	FOLIAR TREATMENT: Apply specified dosage per acre as a foliar spray in sufficient water for thorough coverage but not less than 1 gallon total volume per acre. Do not forage or cut treated corn for food, feed, or ensilage within 28 days of treatment.

RECOMMENDED APPLICATIONS									
CROP	INSECT	DOSAGE DI-SYSTON 8	REMARKS						
FIELD CROPS (Continued) Cotton	Aphids Mites (See Note) Thrips (Except Western flower thrips)	.75 to 1.2 fl oz per 1000 feet of row (for any row spacing*) or 9 to 16 fl oz per acre (40 inch row spacing)	<p>Preplant: As a preplant soil incorporated treatment, DI-SYSTON 8 may be applied in approximately a 16-inch band as a water emulsion.</p> <p>- OR -</p> <p>Planting: Apply as a water emulsion or with liquid fertilizer. Spray/apply in the furrow or soil inject on each side of the furrow. Read "IMPORTANT" statement under "RECOMMENDED APPLICATIONS" for additional information before use on this crop.</p> <p style="text-align: center;">RATE CHART FOR PREPLANT OR PLANTING TIME APPLICATION</p> <table style="width: 100%; border: none;"> <tr> <td style="text-align: center; border: none;"><u>Heavy Soil</u></td> <td style="text-align: center; border: none;"><u>Medium Soil</u></td> <td style="text-align: center; border: none;"><u>Light Soil</u></td> </tr> <tr> <td style="text-align: center; border: none;">1.2 fl oz/1000 ft of row or 1 pt/acre/acre</td> <td style="text-align: center; border: none;">1 fl oz/1000 ft of row or 12 fl oz/acre</td> <td style="text-align: center; border: none;">.75 fl oz/1000 ft of row or 9 fl oz/acre</td> </tr> </table> <p>The preplant or planting application may be followed by the post-plant application described below or by one or two side-dress or broadcast over the foliage soil applications of DI-SYSTON 15% Granular. Refer to the DI-SYSTON 15% Granular label for directions. Do not make more than 3 soil applications of DI-SYSTON per crop season regardless of method of application or formulation used.</p> <p>Post-Plant: If an additional treatment is needed, apply in a water emulsion or with liquid fertilizer as a soil injection on each side of the row. A single post-plant application at rates up to 2 pints/acre may be applied up to first squaring (90 days before harvest) or a single post-plant application at rates up to 1 pint/acre may be applied up to 28 days of harvest. Do not apply more than twice per season. Allow a minimum of 21 days between applications. Read "IMPORTANT" statement under "RECOMMENDED APPLICATIONS" for additional information before use on this crop.</p> <p>Do not graze treated fields after any application of DI-SYSTON 8.</p> <p>If preplant, planting or post-plant applications are made, no foliar application may be made within the same crop season.</p> <p>NOTE: West of the Rocky Mountains, some mite populations have developed resistant to organo-phosphate insecticides, including DI-SYSTON. Consult your local Cooperative Extension Service, agricultural advisor or Miles representative for details.</p>	<u>Heavy Soil</u>	<u>Medium Soil</u>	<u>Light Soil</u>	1.2 fl oz/1000 ft of row or 1 pt/acre/acre	1 fl oz/1000 ft of row or 12 fl oz/acre	.75 fl oz/1000 ft of row or 9 fl oz/acre
<u>Heavy Soil</u>	<u>Medium Soil</u>	<u>Light Soil</u>							
1.2 fl oz/1000 ft of row or 1 pt/acre/acre	1 fl oz/1000 ft of row or 12 fl oz/acre	.75 fl oz/1000 ft of row or 9 fl oz/acre							

RECOMMENDED APPLICATIONS				
CROP	INSECT	DOSAGE DI-SYSTON 8	REMARKS	
FIELD CROPS (Continued) Cotton	Aphids Thrips (Except Western flower thrips)	3 to 9 fl oz	FOLIAR - PRIOR TO BLOOM: Apply specified dosage per acre as a foliar spray, using aircraft or ground equipment, in sufficient water for thorough coverage but not less than 1 gallon total volume per acre. Repeat as necessary but do not apply more than 3 times per season with the last application no later than bloom.	
		NOTE: If foliar applications are made, do not apply soil treatments within the same crop year.		
Sorghum	Aphids (Greenbugs) (See NOTE Below)	.9 to 1.2 fl oz per 1000 ft. of row (for any row spacing*) (for 40 inch row spacing use 12 to 16 fl oz per acre)	AT PLANTING APPLICATION: FURROW Apply as a water emulsion or in liquid fertilizer in the seed furrow approximately 1/2 inch above the seed. Do not apply directly on the seed.	
		1.2 fl oz per 1000 ft of row (for any row spacing*) (for 40 inch row spacing use 1 pint per acre)	AT PLANTING APPLICATION: BAND Apply as a water emulsion or in liquid fertilizer in a 4 or 6 inch band directly behind the planter shoe in front of the press wheel. Do not apply directly on the seed.	
		NOTICE: The following side-dress and/or foliar applications may be made in addition to granular or liquid at planting applications. A total of 5 applications may be made to a single sorghum crop (2 soil applications, granular or liquid, plus 3 foliar applications of rates 8 fluid ounces or less/acre).		
	1.2 fl oz per 1000 ft of row (for any row spacing*) (for 40 inch row spacing use 1 pint per acre)	POST PLANTING APPLICATION: SIDE-DRESS Apply as a water emulsion or with liquid fertilizer injected into the row on both side of the furrow at any time up to boot stage, but not within 45 days of use for forage or fodder.		
	Aphids (Greenbugs) (See Note Below)	4 to 8 fl oz per acre	POST PLANTING APPLICATION: FOLIAR SPRAY - A maximum of 3 foliar applications may be made of rates 8 fluid ounces or less/acre. Applied specified dosage per acre in sufficient water to obtain thorough coverage. For aerial application use not less than 1 gallon of water per acre. Use not less than 5 gallons of water per acre with ground equipment. For Midge: make one application when 50% of the seed heads have emerged from the boot. Make a second application 3 to 5 days later.	
	Sorghum Midge (Arkansas, Louisiana, Oklahoma, Texas)		Days From Last Application to Harvest for Foliar Rates of 4 to 8 Fluid Oz/Acre	
Banks grass mites (in Texas, excluding Trans-Pecos area, and in Oklahoma)	8 fl oz per acre	APPLICATION SCHEDULE		
		FOR GRAIN	FOR FORAGE OR FODDER	
		Any soil plus any foliar	34	45
		1 or 2 foliar	7	45
		3 foliar	34	60
NOTE: Due to Greenbug resistance in certain areas, DI-SYSTON used alone may not provide satisfactory control. Consult your local agricultural advisor or Extension Service for recommendations.				

RECOMMENDED APPLICATIONS			
CROP	INSECT	DOSAGE DI-SYSTON 8	REMARKS
FIELD CROPS (Continued) Tobacco	Aphids Flea beetles	4 pints (1/2 gallon) per acre	Apply as a water emulsion broadcast spray in sufficient water for thorough coverage. Disk or till the soil to a depth of 2 to 3 inches before setting plants. Do not make more than one field application per crop season regardless of the formulation of DI-SYSTON used.
Wheat (Foliar Application Spring or Fall)	Aphids (Greenbugs) Mites (See NOTE in "REMARKS")	4 to 12 fl oz per acre	<p>For application by aircraft or ground equipment: Apply specified dosage per acre in sufficient water, to obtain thorough coverage but not less than 1 gallon total volume per acre. Use the lower rates on plants up to tillering (stooling). Higher rates should be used after plants begin to tiller.</p> <p>For application in liquid fertilizer: Mix 6 to 12 fluid ounces of DI-SYSTON 8 in the amount of liquid fertilizer to be applied per acre. Apply this mixture as a top dressing with suitable ground equipment.</p> <p>Fall Application: Two applications may be made in the fall. Allow 30 days between applications.</p> <p>Spring Applications: Application at green-up may be made following "fall application". A second application may be made 30 days following the first spring application. Do not make any applications within 30 days of grain harvest.</p> <p>NOTE (1) Do not graze treated fields or cut for forage after any application of DI-SYSTON 8. (2) Due to Greenbug resistance in certain areas, DI-SYSTON used alone may not provide satisfactory control. Consult your local agricultural advisor or Extension Service for recommendations.</p>
Wheat (Fall)	Hessian fly Aphids (Greenbugs and Oat bird cherry Except California)	0.25 fl oz per 1000 ft of row* (see chart below) - OR - up to 1 pint per acre	<p>Apply as a soil injection in water emulsion or with liquid fertilizer at planting time in the fall.</p> <p>NOTE (1) Do not graze treated fields or cut for forage after any application of DI-SYSTON 8. (2) Due to Greenbug resistance in certain areas, DI-SYSTON used alone may not provide satisfactory control. Consult your local agricultural advisor or Extension Service for recommendations.</p>
		Ounces DI-SYSTON 8 per 1000 ft. of row based on	
		Row Spacing	Ounces/Acre on 3 inch band
		7 or 8 inch rows	15.5 .25
		10 inch rows	13.25 .25
		12 inch rows	10.75 .25
		14 inch rows	9.5 .25
		16 inch rows	8.5 .25
		18 inch rows	7.0 .25

RECOMMENDED APPLICATIONS			
CROP	INSECT	DOSAGE DI-SYSTON 8	REMARKS
NON-BEARING FRUIT Strawberries (Propagating Plants Only)	Aphids Mites (except cyclamen mite)	2.8 to 5 fl oz per 1000 feet of row (for any row spacing*) - OR - 2 to 4 pints per acre (42 inch row spacing)	Apply in a water emulsion or with liquid fertilizer as a soil injection on each side of the row as needed. Do not use fruit from treated plants for food purposes.
NUTS Pecans (South Central and South Western States Only)	Aphids	1.5 to 3 pints per acre	Apply by spraying 6-foot bands of soil on both sides of the trees. The bands should be located under the main drip area of the trees. Work the treated soil to a depth of 2 to 3 inches. On irrigated pecans, apply water as soon as possible after treatment. Make only 1 application per season. Early spring treatment is recommended. Do not harvest nuts within 80 days of treatment. Do not graze grass under treated trees. Do not tank mix this product with Phosalone.
Pecans	Aphids	4 to 6 fl oz	FOLIAR APPLICATIONS: Apply specified dosage in 100 gallons of water as a full coverage spray using not more than 400 gallons of finished spray per acre. Do not apply more than 3 times per season nor within 30 days of harvest. Do not graze grass under treated area. Do not tank mix this product with Phosalone.
		3/4 to 1 pint per acre	APPLICATION BY AIRCRAFT: Apply specified dosage per acre in sufficient water to obtain thorough coverage but not less than 5 gallons total volume per acre. Do not apply more than 3 times per season nor within 30 days of harvest. Do not graze grass under treated trees. Do not tank mix this product with Phosalone.

RECOMMENDED APPLICATIONS			
CROP	INSECT	DOSAGE DI-SYSTON 8	REMARKS
VEGETABLES Beans	Aphid Leafhopper Mexican bean beetle (larvae) Mites Thrips	.9 to 1.9 fl oz per 1000 feet of row (for any row spacing*) -OR- 1 to 2 pints per acre (30 inch row spacing) (For extremely narrow row spacing, do not exceed the 2 pints per acre rate)	For snap or green Lima beans, apply in a water emulsion or with liquid fertilizer as a soil injection on each side of the seed furrow at planting time only. For dry beans, apply as above at planting time or inject as a side-dressing on each side of the row after plants become established but not within 60 days of harvest. Do not apply directly on the seed. Do not apply more than once per season. Do not feed vines or hay. Read "IMPORTANT" statement under "Recommended Applications" for additional information before use on this crop.
Brussels sprouts	Aphids Flea beetles Leafhopper Mites Thrips	1.1 fl oz per 1000 feet of row (for any row spacing*) -OR- 1 pint per acre (36 inch row spacing)	Transplant Seed Beds: Apply evenly over bed area in a water emulsion as a broadcast spray prior to seeding at the rate of 1 pint per acre. Work into the top 2 to 3 inches of soil and plant seed in normal manner. Field Application: Apply specified dosage in a water emulsion or with liquid fertilizer as an injection on each side of the seed furrow or transplanted row at planting and where necessary, injected as a side-dressing after plants are established. Do not apply directly on the seed. Do not apply more than two field applications per season nor within 30 days of harvest for Brussels sprouts, or 40 days of harvest for cauliflower. Allow a minimum of 21 days between applications.
Cauliflower	Aphids Flea beetles Leafhopper		
Broccoli	Aphids Flea beetles Leafhopper	1.1 fl oz per 1000 ft of row (for any row spacing*) - OR - 1 pint per acre (36 inch row spacing*)	Transplant Seed Beds: Apply evenly over bed area in a water emulsion as a broadcast spray prior to seeding at the rate of 1 pint per acre. Work into the top 2 to 3 inches of soil and plant seed in normal manner.
Cabbage (Including tight-heading varieties of Chinese cabbage)	Aphids Flea beetles		Field Applications: Apply specified dosage in a water emulsion or with liquid fertilizer as an injection on each side of the seed furrow or transplanted row at planting or injected as a side-dressing after plants are established. Do not apply directly on the seed. Do not apply more than one field application per season. Do not use within 14 days of harvest for broccoli or 42 days of harvest for cabbage.
	Root aphids	1.7 fl oz per 1000 ft of row (for any row spacing*) - OR - 1 to 2 pints per acre (20 inch row spacing)	

RECOMMENDED APPLICATIONS			
CROP	INSECT	DOSAGE DI-SYSTON 8	REMARKS
VEGETABLES (Continued) Lettuce	Aphids Leafhoppers Mites Thrips	.6 to 1.2 fl oz per 1000 feet of row (for any row spacing*) -OR- 1 to 2 pints per acre (20 inch row spacing)	<p>Spring or Summer Seeded Lettuce: Apply in a water emulsion or with liquid fertilizer as an injection on each side of the seed furrow at planting time only. Use the higher rate on heavy organic soils. Do not apply directly on the seed. Do not apply to transplanted lettuce.</p> <p>Fall or Winter Seeded Lettuce: Apply specific dosage in a water emulsion or as a soil injected side-dressing with liquid fertilizer at thinning time. Do not apply within 60 days of harvest. Read "IMPORTANT" statement under "RECOMMENDED APPLICATIONS" for additional information before use on this crop.</p>
Peas Lentils	Aphids Leafhoppers	1 to 2.5 pints per acre	Apply in a water emulsion or with liquid fertilizer as an in-furrow spray or as an injection on each side of the seed furrow at planting or injected as a side-dressing after plants are established. Do not apply more than once per season nor within 50 days of harvest. Do not feed vines or hay.
Potatoes	Aphids	3 to 4 pints per acre	SOIL APPLICATION -- PRE-PLANT BROADCAST: Apply specified dosage per acre as a preplant broadcast application. Incorporate to a depth of 2 to 3 inches. Plant crop in usual manner. Do not make more than 2 soil applications of DI-SYSTON 8 per crop season regardless of method of application or formulation used. Do not tank mix this product with Phosalone.
	Aphids Flea beetles (except tuber beetles) Leafhoppers Potato psyllid Southern potato wireworm Colorado potato beetle (early season reduction)	2.25 to 3.5 fl oz per 1000 feet of row (for any row spacing*) (for 38 inch row spacing use 2 to 3 pints per acre)	<p>SOIL APPLICATION - FURROW: Apply undiluted in specifically designed equipment for low volume soil injection, or apply in a water emulsion, or with liquid fertilizer as an in-furrow spray at planting. An additional side-dress application may be made as a water emulsion injection after plants become established. Do not make more than 2 soil applications of DI-SYSTON per crop season regardless of method of application or formulation used. Do not apply second soil application within 75 days of harvest. Higher rate to be used in organic soils. Do not tank mix this product with Phosalone.</p> <p>NOTE: Colorado potato beetle populations in the northeast states (except Maine), as well as Virginia and North Carolina, have developed resistance to insecticides, including DI-SYSTON. Consult your local Extension Service or Miles sales representative for details.</p>
	Green peach aphids (East of the Rocky Mountains only)	6 to 16 fl oz per acre	FOLIAR APPLICATIONS: Using aircraft or ground equipment, apply specified dosage per acre in sufficient water to obtain thorough coverage, but not less than 1 gallon total volume per acre. Begin applications when pest first appears. Use the higher rates for heavy infestations. Repeat as necessary, but not more than 3 times per season nor within 30 days of harvest. Do not tank mix this product with Phosalone.
NOTE: Foliar applications may be made following at-planting soil treatment as recommended above or in the label for DI-SYSTON 15% Granular.			

* ON ROW CROPS WHERE ROW SPACING IS EXTREMELY NARROW, DO NOT PLACE TREATED ZONES (BAND OR FURROW) CLOSER TOGETHER THAN 6 INCHES.

RECOMMENDED APPLICATIONS			
CROP	INSECT	DOSAGE DI-SYSTON 8	REMARKS
VEGETABLES (Continued) Potatoes	Green peach aphid Colorado potato beetle	3 pints per acre	SPRINKLER IRRIGATION APPLICATION - ONLY IN OREGON, WASHINGTON, IDAHO AND UTAH: A single application at the rate of 3 pints per acre may be made for control of Green peach aphid and Colorado potato beetle. Do not apply within 60 days of harvest. Application may be made following at-planting soil application as recommended on this label or the label for DI-SYSTON 15% Granular. Follow all directions given under the Chemigation section of this label.
Tomatoes	Aphids Flea beetles Leafhoppers Leaf miners Mites Colorado potato beetle (early season reduction)	1.2 to 3.5 fl oz per 1000 feet of row (for any row spacing*) -OR- 1 to 3 pints per acre (38 inch row spacing)	<p>Transplant Seed Beds: Apply evenly over bed area in a water emulsion as a broadcast spray prior to seeding at the rate of 3 pints per acre. Work into the top 2 to 3 inches of soil and plant seed in normal manner.</p> <p>Field Applications: Apply specified dosage in a water emulsion or with liquid fertilizer as an injection on each side of the seed furrow or transplanted row at planting time. Do not apply directly on the seed. Where 2 field applications are required, use only 1.2 to 2.4 fluid ounces per 1000 feet of row, or 1 to 2 pints per acre, injected on each side of the seed furrow or transplanted row at planting and as a side-dressing after plants are established. Allow a minimum of 21 days between applications. Do not apply within 30 days of harvest.</p> <p>(NOTE: <u>Florida Only</u> - Rates up to 10.5 fluid ounces per 1000 feet of row, with a 72-inch row spacing, may be applied to highly alkaline soils as an injection on each side of the seed furrow or transplanted row at planting time only; or where 2 field applications are required, up to 5.25 fluid ounces per 1000 square feet of row with a 72-inch row spacing, at planting time and up to 5.25 fluid ounces per 1000 feet of row as a side-dressing after plants become established. Do not apply within 30 days of harvest.)</p> <p>Read "IMPORTANT" statement under "RECOMMENDED APPLICATIONS" for additional information before use on this crop.</p>

* ON ROW CROPS WHERE ROW SPACING IS EXTREMELY NARROW, DO NOT PLACE TREATED ZONES (BAND OR FURROW) CLOSER TOGETHER THAN 6 INCHES.

STORAGE AND DISPOSAL

Pesticide Disposal: Do not contaminate water, food or feed by storage or disposal.

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking invert to prevent leakage. If the container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides above. In spill or leak incidents, keep unauthorized people away. You may contact the Miles Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Miles Kansas City Emergency Response Telephone No. is 816-242-2582, or contact Chemtrec at 800-424-9300.

Miles, Inc.
Crop Protection Products
Box 4913, Kansas City, MO 64120-0013