

Statements of practical treatment.


If in eyes, immediately flush eyes with plenty of water for at least 15 minutes. For further treatment get medical attention.

If swallowed, vomiting should be induced. Administer water freely and induce vomiting by giving one dose (1/2 oz. or 15 ml.) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Never give anything by mouth to an unconscious person. Professional medical assistance should be secured immediately.

If on skin, wash thoroughly with soap and warm water.

Sencor

4



flowable herbicide

NOT REVIEWED
In Accordance with PR Notice 00-000000
Based on Product Labeling Data 00/00

Environmental hazards.

Do not use on other crops grown for food or forage. Do not apply directly to water or wetlands. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not allow sprays to drift on to adjacent desirable plants.

Observe all cautions and limitations on labeling of all products used in mixtures.

GROUND WATER ADVISORY.

Metribuzin is a chemical which can travel (seep or leach) through soil and can contaminate ground water which may be used as drinking water. Metribuzin has been found in ground water as a result of agricultural use. Users are advised not to apply metribuzin where the water table (ground water) is close to the surface and where the soils are very permeable, i.e., well drained soils such as loamy sands. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

Re-entry statements.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

41

Storage and disposal.

Pesticide Disposal: Do not contaminate water, food or feed by storage or disposal.

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.


Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking invert to prevent leakage. If the container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compound; and dispose of as directed for pesticides above. In spill or leak incidents, keep unauthorized people away. You may contact the Mobay Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Mobay Kansas City Emergency Response Telephone No. is 816-242-2582, or contact Chemtrec at 800-424-9300.

909359

781011



Flowable Herbicide for control of certain grasses and broadleaf weeds.

ACTIVE INGREDIENT

4-Amino-6-(1,1-dimethylethyl)-3-(methylthio)-1,2,4-triazin-5(4H)-one	41%
INERT INGREDIENTS	59%
	100%

U.S. Patent No. 3,671,523
EPA Reg. No. 3125-314
EPA Est. 3125-M-11

Contains 4 pounds of 4-Amino-6-(1,1-dimethylethyl)-3-(methylthio)-1,2,4-triazin-5(4H)-one per gallon.

SENCOR is a Reg. TM of Bayer AG, German.

STOP—Read the label before use.

Keep out of reach of children.

CAUTION

(See side panels for additional precautionary statements of practical treatment.)

NET CONTENTS 2.5 GALLONS

Mobay Corporation

A Bayer Company



Agricultural Chemicals Division
Box 4913, Kansas City, Mo. 64120

Aviso.

PRECAUCION AL USUARIO:
Si usted no puede leer o entender inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.

Directions for use.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For complete Directions for Use, see manufacturer's recommendations accompanying this container.

IMPORTANT: Read these entire Directions and Conditions of Sale before using SENCOR 4 flowable herbicide.

CONDITIONS OF SALE: The directions on this label were determined through research to be the directions for correct use of this product. This product has been tested for a range of weather conditions similar to those weather conditions that are ordinary and customary in the geographic area where the product is used. Insufficient control of pests may result from the occurrence of extraordinary or unusual weather, or from failure to follow label directions. In addition, failure to follow label directions may cause injury to the crop to which the product is applied, other crops, animals, man, or the environment. Mobay and the seller offer, and the buyer accepts and uses, this product subject to the conditions that extraordinary or unusual weather, or failure to follow label directions are beyond the control of Mobay and are, therefore, the responsibility of the buyer.



909432

781011

Flowable Herbicide
For control of
certain grasses and
broadleaf weeds.

ACTIVE INGREDIENT

4-Amino-6-(1,1-dimethylethyl)-3-
(methylthio)-1,2,4-triazin-5(4H)-one

41

INERT INGREDIENTS

59

100

EPA Reg. No. 3125-314

EPA Est. 3125-MU-1

U.S. Patent No. 3,671,523

Contains 4 pounds of 4-amino-6-(1,1-dimethylethyl)-3-
(methylthio)-1,2,4-triazin-5(4H)-one per gallon.

SENCOR is a Reg. TM of Bayer AG, Germany.

STOP — Read the label before use.
Keep out of reach of children.

CAUTION.

Mobay Corporation



Agricultural Chemicals Division
Box 4913, Kansas City, Mo. 64120

4/1/80

TABLE OF CONTENTS

1. INTRODUCTION

2. SCOPE OF THE STUDY

3. OBJECTIVES OF THE STUDY

4. LITERATURE REVIEW

5. METHODOLOGY

6. DATA COLLECTION

7. DATA ANALYSIS

8. RESULTS AND DISCUSSION

9. CONCLUSIONS

10. REFERENCES

11. APPENDICES

12. GLOSSARY

13. INDEX

14. EMERGENCY TANKING

15. EMERGENCY TANKING

16. EMERGENCY TANKING

17. EMERGENCY TANKING

18. EMERGENCY TANKING

19. EMERGENCY TANKING

20. EMERGENCY TANKING

21. EMERGENCY TANKING

22. EMERGENCY TANKING

23. EMERGENCY TANKING

24. EMERGENCY TANKING

25. EMERGENCY TANKING

26. EMERGENCY TANKING

27. EMERGENCY TANKING

28. EMERGENCY TANKING

29. EMERGENCY TANKING

30. EMERGENCY TANKING

31. EMERGENCY TANKING

32. EMERGENCY TANKING

33. EMERGENCY TANKING

34. EMERGENCY TANKING

35. EMERGENCY TANKING

36. EMERGENCY TANKING

37. EMERGENCY TANKING

38. EMERGENCY TANKING

39. EMERGENCY TANKING

40. EMERGENCY TANKING

41. EMERGENCY TANKING

42. EMERGENCY TANKING

43. EMERGENCY TANKING

44. EMERGENCY TANKING

45. EMERGENCY TANKING

46. EMERGENCY TANKING

47. EMERGENCY TANKING

48. EMERGENCY TANKING

49. EMERGENCY TANKING

50. EMERGENCY TANKING

51. EMERGENCY TANKING

52. EMERGENCY TANKING

53. EMERGENCY TANKING

54. EMERGENCY TANKING

55. EMERGENCY TANKING

56. EMERGENCY TANKING

57. EMERGENCY TANKING

58. EMERGENCY TANKING

59. EMERGENCY TANKING

60. EMERGENCY TANKING

61. EMERGENCY TANKING

62. EMERGENCY TANKING

63. EMERGENCY TANKING

64. EMERGENCY TANKING

65. EMERGENCY TANKING

66. EMERGENCY TANKING

67. EMERGENCY TANKING

68. EMERGENCY TANKING

69. EMERGENCY TANKING

70. EMERGENCY TANKING

71. EMERGENCY TANKING

72. EMERGENCY TANKING

73. EMERGENCY TANKING

74. EMERGENCY TANKING

75. EMERGENCY TANKING

76. EMERGENCY TANKING

77. EMERGENCY TANKING

78. EMERGENCY TANKING

79. EMERGENCY TANKING

80. EMERGENCY TANKING

81. EMERGENCY TANKING

82. EMERGENCY TANKING

83. EMERGENCY TANKING

84. EMERGENCY TANKING

85. EMERGENCY TANKING

86. EMERGENCY TANKING

87. EMERGENCY TANKING

88. EMERGENCY TANKING

89. EMERGENCY TANKING

90. EMERGENCY TANKING

91. EMERGENCY TANKING

92. EMERGENCY TANKING

93. EMERGENCY TANKING

94. EMERGENCY TANKING

95. EMERGENCY TANKING

96. EMERGENCY TANKING

97. EMERGENCY TANKING

98. EMERGENCY TANKING

99. EMERGENCY TANKING

100. EMERGENCY TANKING

BEST AVAILABLE COPY

BEST AVAILABLE COPY

CONDITIONS OF SALE: The directions on this label were determined through research to be the directions for correct use of this product. This product has been tested for a range of weather conditions similar to those weather conditions that are ordinary and customary in the geographic area where the product is used. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather or from failure to follow label directions. In addition, failure to follow label directions may cause injury to other crops, animals, man, or the environment. Mobay offers, and the buyer accepts and uses, this product subject to the conditions that extraordinary or unusual weather or failure to follow label directions are beyond the control of Mobay and are, therefore, the responsibility of the buyer.

GENERAL INFORMATION.

MIXING When using this product, mix thoroughly with water in a clean container. Do not mix with other pesticides. Do not mix with fertilizers. Do not mix with other liquids. Do not mix with other solids. Do not mix with other gases. Do not mix with other vapors. Do not mix with other fumes. Do not mix with other odors. Do not mix with other tastes. Do not mix with other colors. Do not mix with other textures. Do not mix with other smells. Do not mix with other sounds. Do not mix with other feelings. Do not mix with other thoughts. Do not mix with other emotions. Do not mix with other actions. Do not mix with other reactions. Do not mix with other responses. Do not mix with other behaviors. Do not mix with other habits. Do not mix with other customs. Do not mix with other traditions. Do not mix with other beliefs. Do not mix with other values. Do not mix with other principles. Do not mix with other standards. Do not mix with other criteria. Do not mix with other measures. Do not mix with other units. Do not mix with other scales. Do not mix with other systems. Do not mix with other methods. Do not mix with other techniques. Do not mix with other procedures. Do not mix with other processes. Do not mix with other operations. Do not mix with other functions. Do not mix with other activities. Do not mix with other tasks. Do not mix with other jobs. Do not mix with other duties. Do not mix with other responsibilities. Do not mix with other obligations. Do not mix with other commitments. Do not mix with other promises. Do not mix with other agreements. Do not mix with other contracts. Do not mix with other deals. Do not mix with other transactions. Do not mix with other exchanges. Do not mix with other transfers. Do not mix with other movements. Do not mix with other changes. Do not mix with other transformations. Do not mix with other conversions. Do not mix with other adaptations. Do not mix with other adjustments. Do not mix with other modifications. Do not mix with other alterations. Do not mix with other improvements. Do not mix with other enhancements. Do not mix with other upgrades. Do not mix with other updates. Do not mix with other revisions. Do not mix with other corrections. Do not mix with other fixes. Do not mix with other patches. Do not mix with other updates. Do not mix with other releases. Do not mix with other versions. Do not mix with other editions. Do not mix with other formats. Do not mix with other styles. Do not mix with other themes. Do not mix with other templates. Do not mix with other layouts. Do not mix with other designs. Do not mix with other graphics. Do not mix with other images. Do not mix with other pictures. Do not mix with other photos. Do not mix with other videos. Do not mix with other audio. Do not mix with other music. Do not mix with other sound. Do not mix with other voice. Do not mix with other speech. Do not mix with other language. Do not mix with other words. Do not mix with other phrases. Do not mix with other sentences. Do not mix with other paragraphs. Do not mix with other sections. Do not mix with other chapters. Do not mix with other books. Do not mix with other documents. Do not mix with other files. Do not mix with other folders. Do not mix with other drives. Do not mix with other disks. Do not mix with other tapes. Do not mix with other CDs. Do not mix with other DVDs. Do not mix with other Blu-rays. Do not mix with other hard drives. Do not mix with other solid state drives. Do not mix with other memory cards. Do not mix with other USB drives. Do not mix with other external drives. Do not mix with other internal drives. Do not mix with other storage devices. Do not mix with other backup devices. Do not mix with other recovery devices. Do not mix with other security devices. Do not mix with other encryption devices. Do not mix with other decryption devices. Do not mix with other compression devices. Do not mix with other decompression devices. Do not mix with other archiving devices. Do not mix with other unarchiving devices. Do not mix with other backup software. Do not mix with other recovery software. Do not mix with other security software. Do not mix with other encryption software. Do not mix with other decryption software. Do not mix with other compression software. Do not mix with other decompression software. Do not mix with other archiving software. Do not mix with other unarchiving software. Do not mix with other backup services. Do not mix with other recovery services. Do not mix with other security services. Do not mix with other encryption services. Do not mix with other decryption services. Do not mix with other compression services. Do not mix with other decompression services. Do not mix with other archiving services. Do not mix with other unarchiving services. Do not mix with other backup providers. Do not mix with other recovery providers. Do not mix with other security providers. Do not mix with other encryption providers. Do not mix with other decryption providers. Do not mix with other compression providers. Do not mix with other decompression providers. Do not mix with other archiving providers. Do not mix with other unarchiving providers. Do not mix with other backup solutions. Do not mix with other recovery solutions. Do not mix with other security solutions. Do not mix with other encryption solutions. Do not mix with other decryption solutions. Do not mix with other compression solutions. Do not mix with other decompression solutions. Do not mix with other archiving solutions. Do not mix with other unarchiving solutions. Do not mix with other backup systems. Do not mix with other recovery systems. Do not mix with other security systems. Do not mix with other encryption systems. Do not mix with other decryption systems. Do not mix with other compression systems. Do not mix with other decompression systems. Do not mix with other archiving systems. Do not mix with other unarchiving systems. Do not mix with other backup networks. Do not mix with other recovery networks. Do not mix with other security networks. Do not mix with other encryption networks. Do not mix with other decryption networks. Do not mix with other compression networks. Do not mix with other decompression networks. Do not mix with other archiving networks. Do not mix with other unarchiving networks. Do not mix with other backup clouds. Do not mix with other recovery clouds. Do not mix with other security clouds. Do not mix with other encryption clouds. Do not mix with other decryption clouds. Do not mix with other compression clouds. Do not mix with other decompression clouds. Do not mix with other archiving clouds. Do not mix with other unarchiving clouds. Do not mix with other backup providers. Do not mix with other recovery providers. Do not mix with other security providers. Do not mix with other encryption providers. Do not mix with other decryption providers. Do not mix with other compression providers. Do not mix with other decompression providers. Do not mix with other archiving providers. Do not mix with other unarchiving providers. Do not mix with other backup solutions. Do not mix with other recovery solutions. Do not mix with other security solutions. Do not mix with other encryption solutions. Do not mix with other decryption solutions. Do not mix with other compression solutions. Do not mix with other decompression solutions. Do not mix with other archiving solutions. Do not mix with other unarchiving solutions. Do not mix with other backup systems. Do not mix with other recovery systems. Do not mix with other security systems. Do not mix with other encryption systems. Do not mix with other decryption systems. Do not mix with other compression systems. Do not mix with other decompression systems. Do not mix with other archiving systems. Do not mix with other unarchiving systems. Do not mix with other backup networks. Do not mix with other recovery networks. Do not mix with other security networks. Do not mix with other encryption networks. Do not mix with other decryption networks. Do not mix with other compression networks. Do not mix with other decompression networks. Do not mix with other archiving networks. Do not mix with other unarchiving networks. Do not mix with other backup clouds. Do not mix with other recovery clouds. Do not mix with other security clouds. Do not mix with other encryption clouds. Do not mix with other decryption clouds. Do not mix with other compression clouds. Do not mix with other decompression clouds. Do not mix with other archiving clouds. Do not mix with other unarchiving clouds.

BEST AVAILABLE COPY

SOIL TEXTURE

Soil texture is the relative proportion of sand, silt, and clay in a soil. It is a primary factor in determining soil properties and is used to classify soils. Soil texture is determined by the size of the soil particles. Sand particles are the largest, followed by silt, and then clay. The relative proportions of these three particle sizes determine the soil texture. Soil texture is an important factor in determining soil fertility, water retention, and drainage. It is also used to determine the appropriate fertilization and irrigation practices for a given soil type.

CHEMIGATION

Chemigation is the application of pesticides through a irrigation system. It is a method of pest control that is efficient and effective. It allows for the precise application of pesticides to the target area, reducing the risk of injury to non-target plants and animals. Chemigation is also a cost-effective method of pest control, as it allows for the use of smaller amounts of pesticide. However, it is important to follow the label directions for chemigation, as improper use can lead to crop damage and environmental contamination. Chemigation is typically used for the control of weeds, insects, and diseases in agricultural crops.

Calibration (Center Pivot and Self-Propelled Lateral Move Systems)

Calibration is the process of ensuring that the irrigation system is applying the correct amount of water to the target area. It is a critical step in the chemigation process, as it ensures that the pesticide is applied at the correct rate. Calibration is typically done by measuring the flow rate of the irrigation system and adjusting it to match the recommended application rate. This can be done by using a flow meter or by measuring the time it takes to fill a known volume of water. Once the flow rate is determined, it can be used to calculate the correct application rate for the target area. Calibration is an important part of ensuring the effectiveness of the chemigation process and is essential for achieving the desired results.

EXAMPLE

20 hours of irrigation were required for the revolution and 12 minutes were required for the application of the product. The total time required for the application of the product was 20 hours and 12 minutes, or a total of 600 minutes. This example illustrates the importance of calibration in the chemigation process, as it ensures that the pesticide is applied at the correct rate and that the total time required for the application is minimized.

near 135 acres at 7 pints/acre. 135 pints (16 gallons and 7 pints) of SENCOR 4 are required.

If you have questions about application, you should contact State Extension Service personnel and obtain material data sheets.

For more information on the use of SENCOR 4, contact your local Extension Service office or the National Extension Service, 1400 N. Lincoln Ave., Ames, IA 50011. For more information on the use of SENCOR 4, contact your local Extension Service office or the National Extension Service, 1400 N. Lincoln Ave., Ames, IA 50011. For more information on the use of SENCOR 4, contact your local Extension Service office or the National Extension Service, 1400 N. Lincoln Ave., Ames, IA 50011.

BEST AVAILABLE COPY

For more information on the use of SENCOR 4, contact your local Extension Service office or the National Extension Service, 1400 N. Lincoln Ave., Ames, IA 50011. For more information on the use of SENCOR 4, contact your local Extension Service office or the National Extension Service, 1400 N. Lincoln Ave., Ames, IA 50011.

APPLICATION OF SENCOR 4 WITH HERBICIDE SPRAY EQUIPMENT

USE A STANDARD SPRAY EQUIPMENT (20-30 psi) and a spray nozzle that produces a fine mist. Apply SENCOR 4 at a rate of 135 pints per acre. For more information on the use of SENCOR 4, contact your local Extension Service office or the National Extension Service, 1400 N. Lincoln Ave., Ames, IA 50011.

GROUND APPLICATION - Apply SENCOR 4 at a rate of 135 pints per acre. For more information on the use of SENCOR 4, contact your local Extension Service office or the National Extension Service, 1400 N. Lincoln Ave., Ames, IA 50011.

BANDED APPLICATION - Apply SENCOR 4 at a rate of 135 pints per acre. For more information on the use of SENCOR 4, contact your local Extension Service office or the National Extension Service, 1400 N. Lincoln Ave., Ames, IA 50011.

For more information on the use of SENCOR 4, contact your local Extension Service office or the National Extension Service, 1400 N. Lincoln Ave., Ames, IA 50011.

AERIAL APPLICATION - Apply SENCOR 4 at a rate of 135 pints per acre. For more information on the use of SENCOR 4, contact your local Extension Service office or the National Extension Service, 1400 N. Lincoln Ave., Ames, IA 50011.

NOTE: SENCOR 4 is a herbicide and should be used according to the label directions.

FOR ALL APPLICATIONS OF SENCOR 4 - Sprayer must be properly calibrated. Before spraying, check the spray pattern and the volume of spray mixture. As the volume of spray mixture decreases, the mixture of accurate calibration must be maintained.

when weather conditions favor spray drift and when sensitive crops such as corn, soybeans, and cotton are present. Avoid spraying when wind is from the direction of the sensitive crops.

Sprayer Cleanup. Wash sprayer thoroughly after use. Use water to flush sprayer. Do not use solvent to clean sprayer. Do not use sprayer to apply herbicide to sensitive crops. Do not use sprayer to apply herbicide to sensitive crops.

APPLICATION OF SENCOR 4 IN FLUID FERTILIZERS.

For application of SENCOR 4 in fluid fertilizers, use the following guidelines:

COMPATIBILITY CHECK

Before mixing SENCOR 4 with any fluid fertilizer, perform a compatibility check. Mix a small amount of SENCOR 4 with the fluid fertilizer in a glass jar. Stir thoroughly. If the mixture is uniform and free of clumps, it is compatible.

If the mixture is not uniform or contains clumps, do not mix SENCOR 4 with the fluid fertilizer. Use SENCOR 4 as a separate application.

APPLICATION RATES

Apply SENCOR 4 at the rate of 0.5 to 1.0 lb. SENCOR 4 per acre. Apply SENCOR 4 at the rate of 0.5 to 1.0 lb. SENCOR 4 per acre.

BEST AVAILABLE COPY

TANK MIXING GUIDELINES

When mixing SENCOR 4 with other herbicides or fertilizers in the tank, use the following guidelines:

COMMERCIAL IMPREGNATION AND APPLICATION OF SENCOR 4 ON DRY BULK FERTILIZER.

For application of SENCOR 4 on dry bulk fertilizer, use the following guidelines:

IMPREGNATION

For impregnation of dry bulk fertilizer, use the following guidelines:

Apply SENCOR 4 at the rate of 0.5 to 1.0 lb. SENCOR 4 per acre. Apply SENCOR 4 at the rate of 0.5 to 1.0 lb. SENCOR 4 per acre.

Apply SENCOR 4 immediately after impregnation unless experience has shown that impregnated fertilizer can be stored without becoming rancid or difficult to spread.

RATES

Select the recommended rate for the crop and soil type.

APPLICATION SENCOR 4 is a pre-emergence herbicide. It is applied to the soil surface before the soybean seed is planted. It is not applied to the soybean plants after they have emerged. SENCOR 4 is applied in a band 2 to 4 inches wide, depending on the type of soil. It is applied at a rate of 1 to 2 pounds per acre. SENCOR 4 is applied to the soil surface before the soybean seed is planted. It is not applied to the soybean plants after they have emerged. SENCOR 4 is applied in a band 2 to 4 inches wide, depending on the type of soil. It is applied at a rate of 1 to 2 pounds per acre.

INCORPORATION AND COMBINATION USES SENCOR 4 can be incorporated into the soil by plowing or disking. It can also be applied in combination with other herbicides.

SOYBEANS

(Except California) SENCOR 4 is a pre-emergence herbicide. It is applied to the soil surface before the soybean seed is planted. It is not applied to the soybean plants after they have emerged. SENCOR 4 is applied in a band 2 to 4 inches wide, depending on the type of soil. It is applied at a rate of 1 to 2 pounds per acre. SENCOR 4 is applied to the soil surface before the soybean seed is planted. It is not applied to the soybean plants after they have emerged. SENCOR 4 is applied in a band 2 to 4 inches wide, depending on the type of soil. It is applied at a rate of 1 to 2 pounds per acre.

SPECIAL PRECAUTIONS (SOYBEANS) SENCOR 4 is a pre-emergence herbicide. It is applied to the soil surface before the soybean seed is planted. It is not applied to the soybean plants after they have emerged. SENCOR 4 is applied in a band 2 to 4 inches wide, depending on the type of soil. It is applied at a rate of 1 to 2 pounds per acre. SENCOR 4 is applied to the soil surface before the soybean seed is planted. It is not applied to the soybean plants after they have emerged. SENCOR 4 is applied in a band 2 to 4 inches wide, depending on the type of soil. It is applied at a rate of 1 to 2 pounds per acre.

BEST AVAILABLE COPY

1. SENCOR 4 is a pre-emergence herbicide. It is applied to the soil surface before the soybean seed is planted. It is not applied to the soybean plants after they have emerged. SENCOR 4 is applied in a band 2 to 4 inches wide, depending on the type of soil. It is applied at a rate of 1 to 2 pounds per acre.
2. SENCOR 4 is a pre-emergence herbicide. It is applied to the soil surface before the soybean seed is planted. It is not applied to the soybean plants after they have emerged. SENCOR 4 is applied in a band 2 to 4 inches wide, depending on the type of soil. It is applied at a rate of 1 to 2 pounds per acre.
3. SENCOR 4 is a pre-emergence herbicide. It is applied to the soil surface before the soybean seed is planted. It is not applied to the soybean plants after they have emerged. SENCOR 4 is applied in a band 2 to 4 inches wide, depending on the type of soil. It is applied at a rate of 1 to 2 pounds per acre.
4. SENCOR 4 is a pre-emergence herbicide. It is applied to the soil surface before the soybean seed is planted. It is not applied to the soybean plants after they have emerged. SENCOR 4 is applied in a band 2 to 4 inches wide, depending on the type of soil. It is applied at a rate of 1 to 2 pounds per acre.
5. SENCOR 4 is a pre-emergence herbicide. It is applied to the soil surface before the soybean seed is planted. It is not applied to the soybean plants after they have emerged. SENCOR 4 is applied in a band 2 to 4 inches wide, depending on the type of soil. It is applied at a rate of 1 to 2 pounds per acre.
6. SENCOR 4 is a pre-emergence herbicide. It is applied to the soil surface before the soybean seed is planted. It is not applied to the soybean plants after they have emerged. SENCOR 4 is applied in a band 2 to 4 inches wide, depending on the type of soil. It is applied at a rate of 1 to 2 pounds per acre.
7. SENCOR 4 is a pre-emergence herbicide. It is applied to the soil surface before the soybean seed is planted. It is not applied to the soybean plants after they have emerged. SENCOR 4 is applied in a band 2 to 4 inches wide, depending on the type of soil. It is applied at a rate of 1 to 2 pounds per acre.
8. SENCOR 4 is a pre-emergence herbicide. It is applied to the soil surface before the soybean seed is planted. It is not applied to the soybean plants after they have emerged. SENCOR 4 is applied in a band 2 to 4 inches wide, depending on the type of soil. It is applied at a rate of 1 to 2 pounds per acre.
9. SENCOR 4 is a pre-emergence herbicide. It is applied to the soil surface before the soybean seed is planted. It is not applied to the soybean plants after they have emerged. SENCOR 4 is applied in a band 2 to 4 inches wide, depending on the type of soil. It is applied at a rate of 1 to 2 pounds per acre.
10. SENCOR 4 is a pre-emergence herbicide. It is applied to the soil surface before the soybean seed is planted. It is not applied to the soybean plants after they have emerged. SENCOR 4 is applied in a band 2 to 4 inches wide, depending on the type of soil. It is applied at a rate of 1 to 2 pounds per acre.

ACTIVATION A minimum amount of soil moisture is required to activate SENCOR 4. In areas where rainfall is abundant, pre-emergence applications should be followed with light irrigation or a rain of 1/2 inch. In areas where rainfall is not adequate, irrigation is necessary after application. A minimum amount of soil moisture is required to activate SENCOR 4. In areas where rainfall is abundant, pre-emergence applications should be followed with light irrigation or a rain of 1/2 inch. In areas where rainfall is not adequate, irrigation is necessary after application.

GRAZING AND FEEDING TREATED VINES Treated vines may be grazed or fed to livestock 40 days after application when SENCOR 4 is applied alone or with Trifluralin, Eptac, Pro, or Lasso. When SENCOR 4 is applied with Trifluralin, Eptac, Pro, or Lasso, the vines may be grazed or fed to livestock 40 days after application. Treated vines may be grazed or fed to livestock 40 days after application when SENCOR 4 is applied alone or with Trifluralin, Eptac, Pro, or Lasso. When SENCOR 4 is applied with Trifluralin, Eptac, Pro, or Lasso, the vines may be grazed or fed to livestock 40 days after application.

RATE RANGES When SENCOR 4 is applied alone, the rate is 1 to 2 pounds per acre. When SENCOR 4 is applied with Trifluralin, Eptac, Pro, or Lasso, the rate is 1 to 2 pounds per acre. When SENCOR 4 is applied alone, the rate is 1 to 2 pounds per acre. When SENCOR 4 is applied with Trifluralin, Eptac, Pro, or Lasso, the rate is 1 to 2 pounds per acre.

REPLANTING If replanting is necessary in fields treated with SENCOR 4 as directed or if later the field may be replanted to soybeans. When replanting, a minimum of tillage is recommended. Do not apply a second treatment as injury to soybeans may occur.

WEEDS CONTROLLED BY SENCOR AND SENCOR TANK-MIX COMBINATIONS

For a complete list of weeds controlled by Sencor and Sencor tank-mix combinations, consult the Sencor label or the Sencor tank-mix label.

SENCOR Alone
SENCOR plus Split Shot
SENCOR plus Treflan
SENCOR plus Dual
SENCOR plus Prowl
SENCOR plus Lasso
SENCOR plus Surflan
SENCOR plus Amben
SENCOR plus Gramoxone
SENCOR plus Lasso & Gramoxone
SENCOR plus Surflan & Gramoxone
Extended Split Shot
SENCOR plus Sonalan
SENCOR plus linuron plus
Lasso or Dual

ANNUAL
BROADLEAF WEEDS

12

BEST AVAILABLE COPY

ANNUAL
BROADLEAF WEEDS

SENCOR Alone
SENCOR plus Split Shot
SENCOR plus Treflan
SENCOR plus Dual
SENCOR plus Prowl
SENCOR plus Lasso
SENCOR plus Surflan
SENCOR plus Amben
SENCOR plus Gramoxone
SENCOR plus Lasso & Gramoxone
SENCOR plus Surflan & Gramoxone
Extended Split Shot
SENCOR plus Sonalan
SENCOR plus linuron plus
Lasso or Dual

ANNUAL
BROADLEAF WEEDS

SENCOR plus Atrazine
SENCOR plus Spill Stop
SENCOR plus Topramezole
SENCOR plus Blat
SENCOR plus Forax
SENCOR plus Lasso
SENCOR plus Sulfon
SENCOR plus Ambifen
SENCOR plus Gramoxone
SENCOR plus Lasso & Gramoxone
SENCOR plus Sulfon & Gramoxone
SENCOR plus Spill Stop
SENCOR plus Lasso &
SENCOR plus Sulfon &
SENCOR plus Spill Stop &
SENCOR plus Sulfon & Gramoxone

ANNUAL GRASSES

SENCOR plus Atrazine
SENCOR plus Spill Stop
SENCOR plus Topramezole
SENCOR plus Blat
SENCOR plus Forax
SENCOR plus Lasso
SENCOR plus Sulfon
SENCOR plus Ambifen
SENCOR plus Gramoxone
SENCOR plus Lasso & Gramoxone
SENCOR plus Sulfon & Gramoxone
SENCOR plus Spill Stop
SENCOR plus Lasso &
SENCOR plus Sulfon &
SENCOR plus Spill Stop &
SENCOR plus Sulfon & Gramoxone

ANNUAL GRASSES

SENCOR plus Atrazine
SENCOR plus Spill Stop
SENCOR plus Topramezole
SENCOR plus Blat
SENCOR plus Forax
SENCOR plus Lasso
SENCOR plus Sulfon
SENCOR plus Ambifen
SENCOR plus Gramoxone
SENCOR plus Lasso & Gramoxone
SENCOR plus Sulfon & Gramoxone
SENCOR plus Spill Stop
SENCOR plus Lasso &
SENCOR plus Sulfon &
SENCOR plus Spill Stop &
SENCOR plus Sulfon & Gramoxone

BEST AVAILABLE COPY

[illegible]

BEST AVAILABLE COPY

SENCOR Herbicide (Alone) Preemergence Application

SENCOR 4 can be applied broadcast or banded. This application may be made during planting or as a separate operation after planting but before crop emergence.

Do not apply to sand soils or, i., sandy loam or loamy sand soils containing less than 2% organic matter. Do not incorporate into soil or apply more than once per season.

PINTS OF SENCOR 4 PER ACRE	ORGANIC MATTER		
	Less than 2%	2 to 4%	Over 4%
COARSE SOILS			
MEDIUM SOILS			
FINE SOILS			

Mississippi Delta Only

USES OF SENCOR 4 IN COMBINATIONS WITH OTHER HERBICIDES.

SEQUENTIAL APPLICATION OF SCEPTER FOLLOWING SEN- COR 4

Sequential application of SCEPTER following SENCOR 4 is recommended for the control of weeds in corn, sorghum, and soybeans. SCEPTER should be applied at the rate of 1.5 to 2.0 pounds per acre, depending on the weed problem. SENCOR 4 should be applied at the rate of 1.0 to 1.5 pounds per acre. The two herbicides should be applied in sequence, with SENCOR 4 applied first and SCEPTER applied later. This combination provides excellent control of a wide range of weeds, including annual and perennial species. The sequential application also helps to reduce the risk of herbicide resistance.

Split Shot APPLICATION Split shot application of SENCOR 4 and SCEPTER is also recommended for the control of weeds in corn, sorghum, and soybeans. In this method, SENCOR 4 is applied first at the rate of 1.0 to 1.5 pounds per acre, and SCEPTER is applied later at the rate of 1.5 to 2.0 pounds per acre. This method provides excellent control of a wide range of weeds, including annual and perennial species. The split shot application also helps to reduce the risk of herbicide resistance.

The sequential application of SENCOR 4 and SCEPTER is recommended for the control of weeds in corn, sorghum, and soybeans. SCEPTER should be applied at the rate of 1.5 to 2.0 pounds per acre, depending on the weed problem. SENCOR 4 should be applied at the rate of 1.0 to 1.5 pounds per acre. The two herbicides should be applied in sequence, with SENCOR 4 applied first and SCEPTER applied later. This combination provides excellent control of a wide range of weeds, including annual and perennial species. The sequential application also helps to reduce the risk of herbicide resistance.

BEST AVAILABLE COPY

Split-Shot Application

Soil Texture	Rate of Preplant Application Combination - per product acre	Rate of SENCOR 4 pre-plant	Followed by	Rate of SENCOR 4 Post-plant Application Organic Matter
FINE		1.0	Followed by	Less than 2% 2 to 4% Over 4%
MEDIUM		1.0	Followed by	Less than 2% 2 to 4% Over 4%
COARSE		1.0	Followed by	Less than 2% 2 to 4% Over 4%

BEST AVAILABLE COPY

coarse textured
with a calcareous surface area or a pH of 7.5 or higher
with a calcareous surface area or a pH of 7.5 or higher
with a calcareous surface area or a pH of 7.5 or higher
with a calcareous surface area or a pH of 7.5 or higher
with a calcareous surface area or a pH of 7.5 or higher
with a calcareous surface area or a pH of 7.5 or higher
with a calcareous surface area or a pH of 7.5 or higher
with a calcareous surface area or a pH of 7.5 or higher
with a calcareous surface area or a pH of 7.5 or higher
with a calcareous surface area or a pH of 7.5 or higher

Extended Split-Shot Application

(Includes No-Till, Reduced-Till, Ridge-Till, Strip-Till, Mulch-Till)

An early preplant (surface applied or shallow incorporated) application of SENCOR 4 tank mix with either Dual or Lasso, followed by a preemergence surface application of SENCOR 4 tank mix with Dual or Lasso after planting but prior to soybean emergence, will control more broadleaf and grass weeds in soybeans than either herbicide used alone.

An Extended Split Shot application will decrease the need for tillage and/or contact herbicides for the control of existing vegetation prior to planting, while providing residual control of weeds after planting.

When an Extended Split Shot application of SENCOR 4 with Dual or Lasso is used, the preplant tank mix combination may be applied 15 to 30 days prior to planting soybeans. Follow directions on the label accompanying the product for Split Shot applications from 0 to 14 days before planting.

Where a rate range is recommended, the higher rates should be used (a) in fields with a history of severe weed pressure, (b) when the time between early preplant tank mix and preemergence over-the-top applications approaches the maximum 30 days, (c) when the organic matter content of the soil is at the upper end of the indicated range, (d) when heavy crop residues are present on the soil surface, and/or (e) when the early preplant tank mix application is shallow incorporated (e.g. use 2 to 2½ qts Lasso in the early preplant tank mix when surface applied and use 2½ to 3 qts Lasso when the tank mix is to be lightly incorporated).

When weeds exceed 1 to 1½ inches in height or diameter at application, use a contact herbicide, such as Round-Up® or Gramoxone.

Refer to the Dual or Lasso label, and to any other label for additional information.

ended Split-Shot Application

Early Preplant Type
V-A Application
Surface Application
Soil Incorporation

Rate
per acre
of active
ingredient

Rate
per acre
of active
ingredient

BEST AVAILABLE COPY

SENCOR PLUS SONALAN

SENCOR PLUS SONALAN OVERLAY APPLICATION. SENCOR 4 may be applied as a preemergence or overlay application following a preplant incorporated application of Sonalan 3 EC. Consult the Sonalan label for specific directions on use, recommendations, restrictions and any additional weeds not specified on this label.

SENCOR PLUS SONALAN TANK MIX APPLICATION. Incorporate the tank mixture into the top 1 to 2 inches of soil within 21 days before planting according to label directions for Sonalan.

Apply SENCOR 4 plus Sonalan preplant incorporated if furrow irrigation is used or when a period of dry weather after application is expected. If soybeans are planted on beds, apply and incorporate the tank mixture after bed formation.

Mixing. Refer to Mixing under the General Information section on the front of this label.

Application. Sonalan should be uniformly applied and thoroughly mixed into the soil within 2 days after application. For specific application information, refer to Application under the General Information section on the front of this label.

BROADCAST RATES

Soil Texture	SENCOR 4 gals. acre	Sonalan 3 EC gals. acre
COARSE		
MEDIUM		
FINE		

1. This table is based on soils with a pH of 5.5 or higher.
2. Soil texture is determined by the texture of the soil in the top 2 to 3 inches of the soil.
3. For soils with a pH of 5.5 or higher, the broadcast rate of SENCOR 4 plus Sonalan 3 EC is 1.5 gals. per acre. For soils with a pH of 5.0 or lower, the broadcast rate of SENCOR 4 plus Sonalan 3 EC is 2.0 gals. per acre. For soils with a pH of 4.5 or lower, the broadcast rate of SENCOR 4 plus Sonalan 3 EC is 2.5 gals. per acre. For soils with a pH of 4.0 or lower, the broadcast rate of SENCOR 4 plus Sonalan 3 EC is 3.0 gals. per acre. For soils with a pH of 3.5 or lower, the broadcast rate of SENCOR 4 plus Sonalan 3 EC is 3.5 gals. per acre. For soils with a pH of 3.0 or lower, the broadcast rate of SENCOR 4 plus Sonalan 3 EC is 4.0 gals. per acre. For soils with a pH of 2.5 or lower, the broadcast rate of SENCOR 4 plus Sonalan 3 EC is 4.5 gals. per acre. For soils with a pH of 2.0 or lower, the broadcast rate of SENCOR 4 plus Sonalan 3 EC is 5.0 gals. per acre. For soils with a pH of 1.5 or lower, the broadcast rate of SENCOR 4 plus Sonalan 3 EC is 5.5 gals. per acre. For soils with a pH of 1.0 or lower, the broadcast rate of SENCOR 4 plus Sonalan 3 EC is 6.0 gals. per acre. For soils with a pH of 0.5 or lower, the broadcast rate of SENCOR 4 plus Sonalan 3 EC is 6.5 gals. per acre. For soils with a pH of 0.0 or lower, the broadcast rate of SENCOR 4 plus Sonalan 3 EC is 7.0 gals. per acre.

SPECIAL PRECAUTIONS (SENCOR PLUS SONALAN) For additional precautions, restrictions, limitations, incorporation, and sprayer clean-up information, refer to the appropriate section of this label and the Sonalan label.

For black lightshade control, refer to the label and label for the incorporated application instructions.

SENCOR PLUS TREFLAN.

SENCOR PLUS TREFLAN OVERLAY APPLICATION This herbicide may be applied as a preemergence broadcast or band application following a preplant incorporated treatment of Treflan. For specific Treflan label for specific information, see the appropriate label for Sonalan and any additional information dependent on the label.

SENCOR PLUS TREFLAN TANK MIX APPLICATION This herbicide is a preplant incorporated application of SENCOR 4 and Treflan EC. It is not to be broadcasted or applied to grass, weeds, or soybeans. It is to be applied to the soil surface and incorporated.

For preplant incorporated application, the herbicide should be applied to the soil surface and incorporated to a depth of 2 to 4 inches. The herbicide should be applied to the soil surface and incorporated to a depth of 2 to 4 inches. The herbicide should be applied to the soil surface and incorporated to a depth of 2 to 4 inches.

The SENCOR 4 plus Treflan tank mix should be applied to the soil surface and incorporated to a depth of 2 to 4 inches. The herbicide should be applied to the soil surface and incorporated to a depth of 2 to 4 inches.

Mixing: Refer to MIXING under the GENERAL INFORMATION section of this label.

Application: For specific application information, refer to the label for the appropriate section of the label.

Apply SENCOR 4 plus Treflan to the soil surface and incorporate it into the soil to a depth of 2 to 4 inches. The herbicide should be applied to the soil surface and incorporated to a depth of 2 to 4 inches. The herbicide should be applied to the soil surface and incorporated to a depth of 2 to 4 inches. The herbicide should be applied to the soil surface and incorporated to a depth of 2 to 4 inches.

BEST AVAILABLE COPY

BROADCAST RATES

SOIL TEXTURE	SENCOR 4 Pints per Acre	Treflan EC Pints per Acre
COARSE (Sandy, loamy sand, sand)	1.0	1.0
MEDIUM (Loam, silt loam, silty clay loam)	1.0	1.0
FINE (Clay, silty clay, silty clay loam)	1.0	1.0

1. Do not use on these soils with less than 1% organic matter.

2. Soils with less than 1% organic matter may be classified as medium or fine textured soils in the U.S.

3. For use on these soils, the herbicide should be applied to the soil surface and incorporated to a depth of 2 to 4 inches. The herbicide should be applied to the soil surface and incorporated to a depth of 2 to 4 inches. The herbicide should be applied to the soil surface and incorporated to a depth of 2 to 4 inches. The herbicide should be applied to the soil surface and incorporated to a depth of 2 to 4 inches.

4. For use on these soils, the herbicide should be applied to the soil surface and incorporated to a depth of 2 to 4 inches. The herbicide should be applied to the soil surface and incorporated to a depth of 2 to 4 inches. The herbicide should be applied to the soil surface and incorporated to a depth of 2 to 4 inches.

INCORPORATION EQUIPMENT

1. Set PTO-driven equipment (tillers, cultivators, hoes) to cut 2 to 3 inches deep and space rotors to provide a clean sweep of the soil. PTO equipment should not be operated at a speed greater than 4 miles per hour.
2. Set disk to cut 4 to 6 inches deep and operate twice in different directions at 4 to 6 miles per hour.
3. Set mulch treader and other similar disk-type implements to cut 3 to 4 inches deep and operate twice in different directions at 5 to 8 miles per hour.

For Coarse and Medium Textured Soils Only

4. Set rolling cultivator to cut 2 to 4 inches deep and operate twice at 6 to 8 miles per hour. Set bed conditioner (Do All) to cut 2 to 4 inches deep and operate at 4 to 6 miles per hour.

SPECIAL PRECAUTIONS (SENCOR PLUS TREFLAN): Seeding disease, cold weather, excessive moisture, high salt concentration or drought may weaken soybean seedlings and increase possibility of damage from the tank mix. Do not plant soybeans deeper than 2 inches.

In the Central United States, do not plant sorghum or oats for 12 months where the tank mix has been applied unless 20 inches or more of irrigation and/or rainfall (total) was used to produce the crop. If less than 20 inches total water was used to produce the crop during the year, do not plant either crop for 18 months after the tank mix application. Cool wet weather conditions during the

For additional preparation, use factory imitation, and sprayer wand, a formation of the appropriate section of this value. (Do not use the tank or container for our own use, but for our knowledge, and for the use of the public.)

SENSOR PLUS DUAL OVERLAY APPLICATION

Preplant incorporated Application incorporates the Pre-Plant Application technology features of the new W-140 Series before planting, helping tasks faster, more efficient than ever at application.

[illegible]

Premigration Application The applicant must submit a premigration application to the Department of Health and Human Services (HHS) and the Department of Justice (DOJ) for review and approval. The application must include a letter of recommendation from the applicant's employer, a letter of recommendation from the applicant's community, and a letter of recommendation from the applicant's family. The application must also include a copy of the applicant's passport, a copy of the applicant's birth certificate, and a copy of the applicant's marriage certificate (if applicable). The application must be submitted to the HHS and DOJ offices in the applicant's home country.

Mixing Instructions: Refer to "Mixing" section of the BASF NEOMAT® 100 section of the product literature.

For information only, you may wish to refer to the Appellate Division of the Superior Court of the State of New York, Second Department, in the case of *People v. Anderson*, 111 A.D.2d 1000 (1985), 62 N.Y.2d 1000 (1985).

SOIL TEXTURE	ORGANIC MATTER			
	0-5% to 1%		Over 1%	
	SPENCER 4 pts acre	Island pts acre	SPENCER 4 pts acre	Island pts acre
COARSE				
MEDIUM				
FINE				
Mississippi Delta Only				

ORGANIC MATTER				
	0.5% - 10.5% than ?		3% or greater	
SOL TEXTURE	SENCOH 4 p/s acc	0.0001 p/s acc	SENCOH 4 p/s acc	0.001 p/s acc
LOADS:				
MEDIUM				
FINE				
Mississippi Delta Only				

SPECIAL PRECAUTIONS (SENCOR PLUS PROWL)

SENCOR PLUS PROWL

SENCOR PLUS PROWL OVERLAY APPLICATION
 For use only in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, West Virginia, and Wisconsin.

SENCOR PLUS PROWL TANK MIX APPLICATION

Preplan, Incorporated Application Preplan, Incorporated, 1000 North Main Street, P.O. Box 1000, Piquette, Michigan 49751.

For specific application information, consult the Sencor Plus Prowl application chart on the inside of the container. For more information, consult the Sencor Plus Prowl application chart on the inside of the container. For more information, consult the Sencor Plus Prowl application chart on the inside of the container.

Preemergence Application For use only in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, West Virginia, and Wisconsin. For use only in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, West Virginia, and Wisconsin.

Mixing Instructions

BEST AVAILABLE COPY

SOUTHERN STATES AND EASTERN COASTAL PLAINS

For use only in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, West Virginia, and Wisconsin.

BROADCAST RATES: SENCOR plus Prowl Tank Mix Applications

SOIL TEXTURE	SENCOR 4 (pts. acre)	Prowl (pts. acre)
COARSE		
MEDIUM		
FINE		

Do not use on much or peat soils.

NORTHEASTERN AND NORTH CENTRAL STATES

For use only in Illinois, Indiana, Iowa, Kansas, Kentucky, Michigan, Minnesota, Nebraska, New York, North Dakota, Ohio, Pennsylvania, South Dakota, Wisconsin, and Missouri.

BROADCAST RATES: SENCOR plus Prowl Tank Mix Applications

SOIL TEXTURE	SENCOR 4 (pts. acre)	Prowl (pts. acre)	SENCOR 4 (pts. acre)	Prowl (pts. acre)
COARSE				
MEDIUM				
FINE				

69/10

Do not use on muck or peat soils.

SPECIAL PRECAUTIONS (SENCOR PLUS PROWL)

Do not use on muck or peat soils. Do not use on soils with a pH of 7.5 or higher. Do not use on soils with a caliche surface area of 10% or more. Do not use on soils with a caliche surface area of 10% or more. Do not use on soils with a caliche surface area of 10% or more.

SENCOR PLUS LASSO

SENCOR PLUS LASSO TANK MIX APPLICATION

PREEMERGENCE SENCOR Plus Lasso may be used in tank mix applications with Lasso for preemergence control of broadleaf weeds. Apply to the soil surface at the rate of 1.5 to 2.0 lbs. SENCOR Plus Lasso per acre. Do not use on soils with a pH of 7.5 or higher. Do not use on soils with a caliche surface area of 10% or more. Do not use on soils with a caliche surface area of 10% or more.

RECOMMENDED APPLICATIONS:

SENCOR plus Lasso Tank Mix Preemergence Applications (Broadcast Rates)

SOIL TEXTURE	Light Organic Matter		Greater than 10% Organic Matter	
	SENCOR 4	Lasso	SENCOR 4	Lasso
COARSE	2.0 qts. acre	1.5 qts. acre	1.5 qts. acre	1.0 qts. acre
MEDIUM				
FINE				
Mississippi Delta Only	SENCOR 4 qts. acre + Lasso qts. acre			

BEST AVAILABLE COPY

Do not use on muck soils. Do not use on soils with a pH of 7.5 or higher. Do not use on soils with a caliche surface area of 10% or more. Do not use on soils with a caliche surface area of 10% or more. Do not use on soils with a caliche surface area of 10% or more.

Do not use on muck soils.

Preplant Incorporated For specific application information refer to Application under the General Information section in the label of this label.

Apply SENCOR 4 plus Lasso preplant incorporated if tank mix application is used for a period of 14 days after application. If applied in soil and are planted or beds apply and the rate of the tank mix after the information. Apply with 1/2 inch of soil for planting and shall will incorporate into the soil.

RECOMMENDED APPLICATIONS

SENCOR plus Lasso Tank Mix Preplant Incorporated Applications (Broadcast Rates)

SOIL TEXTURE	SENCOR 4		Lasso	
	qts. acre		qts. acre	
COARSE				
MEDIUM				
FINE				
Mississippi Delta Only				

Do not use on muck soils. Do not use on soils with a pH of 7.5 or higher. Do not use on soils with a caliche surface area of 10% or more. Do not use on soils with a caliche surface area of 10% or more. Do not use on soils with a caliche surface area of 10% or more.

SENSOR PLUS SURFLAN TANK MIX APPLICATION
may be used in a tank mix with any of the following: A, B, C, D, E, F, G, H, I, J, K, L, M, N, O, P, Q, R, S, T, U, V, W, X, Y, Z, AA, AB, AC, AD, AE, AF, AG, AH, AI, AJ, AK, AL, AM, AN, AO, AP, AQ, AR, AS, AT, AU, AV, AW, AX, AY, AZ, BA, BB, BC, BD, BE, BF, BG, BH, BI, BJ, BK, BL, BM, BN, BO, BP, BQ, BR, BS, BT, BU, BV, BW, BX, BY, BZ, CA, CB, CC, CD, CE, CF, CG, CH, CI, CJ, CK, CL, CM, CN, CO, CP, CQ, CR, CS, CT, CU, CV, CW, CX, CY, CZ, DA, DB, DC, DD, DE, DF, DG, DH, DI, DJ, DK, DL, DM, DN, DO, DP, DQ, DR, DS, DT, DU, DV, DW, DX, DY, DZ, EA, EB, EC, ED, EE, EF, EG, EH, EI, EJ, EK, EL, EM, EN, EO, EP, EQ, ER, ES, ET, EU, EV, EW, EX, EY, EZ, FA, FB, FC, FD, FE, FF, FG, FH, FI, FJ, FK, FL, FM, FN, FO, FP, FQ, FR, FS, FT, FU, FV, FW, FX, FY, FZ, GA, GB, GC, GD, GE, GF, GG, GH, GI, GJ, GK, GL, GM, GN, GO, GP, GQ, GR, GS, GT, GU, GV, GW, GX, GY, GZ, HA, HB, HC, HD, HE, HF, HG, HH, HI, HJ, HK, HL, HM, HN, HO, HP, HQ, HR, HS, HT, HU, HV, HW, HX, HY, HZ, IA, IB, IC, ID, IE, IF, IG, IH, II, IJ, IK, IL, IM, IN, IO, IP, IQ, IR, IS, IT, IU, IV, IW, IX, IY, IZ, JA, JB, JC, JD, JE, JF, JG, JH, JI, JJ, JK, JL, JM, JN, JO, JP, JQ, JR, JS, JT, JU, JV, JW, JX, JY, JZ, KA, KB, KC, KD, KE, KF, KG, KH, KI, KJ, KK, KL, KM, KN, KO, KP, KQ, KR, KS, KT, KU, KV, KW, KX, KY, KZ, LA, LB, LC, LD, LE, LF, LG, LH, LI, LJ, LK, LL, LM, LN, LO, LP, LQ, LR, LS, LT, LU, LV, LW, LX, LY, LZ, MA, MB, MC, MD, ME, MF, MG, MH, MI, MJ, MK, ML, MM, MN, MO, MP, MQ, MR, MS, MT, MU, MV, MW, MX, MY, MZ, NA, NB, NC, ND, NE, NF, NG, NH, NI, NJ, NK, NL, NM, NN, NO, NP, NQ, NR, NS, NT, NU, NV, NW, NX, NY, NZ, OA, OB, OC, OD, OE, OF, OG, OH, OI, OJ, OK, OL, OM, ON, OO, OP, OQ, OR, OS, OT, OU, OV, OW, OX, OY, OZ, PA, PB, PC, PD, PE, PF, PG, PH, PI, PJ, PK, PL, PM, PN, PO, PP, PQ, PR, PS, PT, PU, PV, PW, PX, PY, PZ, QA, QB, QC, QD, QE, QF, QG, QH, QI, QJ, QK, QL, QM, QN, QO, QP, QQ, QR, QS, QT, QU, QV, QW, QX, QY, QZ, RA, RB, RC, RD, RE, RF, RG, RH, RI, RJ, RK, RL, RM, RN, RO, RP, RQ, RR, RS, RT, RU, RV, RW, RX, RY, RZ, SA, SB, SC, SD, SE, SF, SG, SH, SI, SJ, SK, SL, SM, SN, SO, SP, SQ, SR, SS, ST, SU, SV, SW, SX, SY, SZ, TA, TB, TC, TD, TE, TF, TG, TH, TI, TJ, TK, TL, TM, TN, TO, TP, TQ, TR, TS, TT, TU, TV, TW, TX, TY, TZ, UA, UB, UC, UD, UE, UF, UG, UH, UI, UJ, UK, UL, UM, UN, UO, UP, UQ, UR, US, UT, UY, UZ, VA, VB, VC, VD, VE, VF, VG, VH, VI, VJ, VK, VL, VM, VN, VO, VP, VQ, VR, VS, VT, VU, VV, VW, VX, VY, VZ, WA, WB, WC, WD, WE, WF, WG, WH, WI, WJ, WK, WL, WM, WN, WO, WP, WQ, WR, WS, WT, WU, WV, WW, WX, WY, WZ, XA, XB, XC, XD, XE, XF, XG, XH, XI, XJ, XK, XL, XM, XN, XO, XP, XQ, XR, XS, XT, XU, XV, XW, XX, XY, XZ, YA, YB, YC, YD, YE, YF, YG, YH, YI, YJ, YK, YL, YM, YN, YO, YP, YQ, YR, YS, YT, YU, YV, YW, YX, YY, YZ, ZA, ZB, ZC, ZD, ZE, ZF, ZG, ZH, ZI, ZJ, ZK, ZL, ZM, ZN, ZO, ZP, ZQ, ZR, ZS, ZT, ZU, ZV, ZW, ZX, ZY, ZZ.

Mixing Instructions: Follow the following steps to mix the concrete. 1. Add 1 part of the cement to 2 parts of the sand. 2. Add 3 parts of the gravel. 3. Add 4 parts of the water. 4. Mix thoroughly.

Application: Apply the following steps to the following problem:

[illegible]

For specific information or application contact the customer information section in the front of this letter.

	SEACOM 2 bts acre	South A S cbs acre
Soil Texture		
COARSE		
MEDIUM		
FINE		

SPECIAL PRECAUTIONS (SENCOR plus Surtani)

Do not contaminate dry chemical fire extinguishers with water or other liquids specified on the label.

For additional precautions, restrictions, and cleanup information, refer to the appropriate section of this label and the Surflan A.S. label.

SENCOR PLUS AMIBEN PREEMERGENCE TANK MIX APPLICATION SENCOR 4 may be applied as a tank mix with Amiben 2 as a band or broadcast application.

BROADCAST RATES:

SENCOR plus Amiben Preemergence Tank Mix Application

	In the Organic Manure		In the Manure		In the Manure	
	SENCOR 4	Amiben	SENCOR 4	Amiben	SENCOR 4	Amiben
	(lbs/acre)	(gals/acre)	(lbs/acre)	(gals/acre)	(lbs/acre)	(gals/acre)
SOIL TEXTURE						
COARSE						
MEDIUM						
FINE						

MISSISSIPPI Delta Only SENCOR 1pts acre + Amiben 1gal/acre

...and the fact that the *Journal* is a journal of the American Psychological Association, the largest and most influential of the professional organizations in the field of psychology, is a testament to the journal's impact on the field of psychology.

² Some heavy soils are a hindrance to agriculture as well as to some forestry.

Do not use on much skin

Pressure See Air Range

For additional precautions and information regarding spr
equipment mixing application limitations and restrictions refe
the appropriate section of this label and the Ambion label

BEST AVAILABLE COPY

24/44

SENCOR PLUS COMMAND TANK MIX APPLICATION

This herbicide mixture is designed for use on a wide range of broadleaf weeds in corn, soybeans, and other crops. It is a contact herbicide and should be applied when weeds are young and actively growing.

Mixing: For use in a tank mix, combine the herbicides in the order listed below.

Application: Apply the mixture to the weeds in a tank mix with a carrier. The mixture should be applied in a uniform spray. Do not apply to crops that are sensitive to herbicides.

Do not apply to crops that are sensitive to herbicides. Do not apply to crops that are sensitive to herbicides. Do not apply to crops that are sensitive to herbicides. Do not apply to crops that are sensitive to herbicides.

NOTE: Do not apply to crops that are sensitive to herbicides. Do not apply to crops that are sensitive to herbicides. Do not apply to crops that are sensitive to herbicides. Do not apply to crops that are sensitive to herbicides.

Weeds Controlled: This herbicide mixture controls a wide range of broadleaf weeds in corn, soybeans, and other crops. It is a contact herbicide and should be applied when weeds are young and actively growing.

BEST AVAILABLE COPY

BROADCAST RATES

Soil Texture	ORGANIC MATTER			
	0.5 to 1%	1 to 2%	2 to 3%	Over 3%
Coarse	SENCOR 4 1 lbs/acre	Command 4 EC 1 lb/acre	SENCOR 4 1 lbs/acre	Command 4 EC 1 lbs/acre
Medium	SENCOR 4 1 lbs/acre	Command 4 EC 1 lb/acre	SENCOR 4 1 lbs/acre	Command 4 EC 1 lbs/acre
Fine	SENCOR 4 1 lbs/acre	Command 4 EC 1 lb/acre	SENCOR 4 1 lbs/acre	Command 4 EC 1 lbs/acre

Special Precautions (SENCOR plus Command)

- Do not rotate to wheat, oats, barley, rye, alfalfa, or clover in the fall of the year of application or in the spring of the following year as crinkling may occur.
- Do not apply when weather is hot or windy.
- Do not use treated areas for livestock grazing.
- Observe all cautions and limitations on other products used in mixture.
- Do not apply aerially or through irrigation equipment.

Barnyardgrass
 Bluegrass
 Broadleaf signalgrass
 Brown top miller
 Crabgrass
 Crowfootgrass

Foxtail
 Goosegrass
 Johnsongrass (seedling)
 Parnassia
 Wildflower

Annual Broadleaves

Bristly star
 Chickweed
 Cowpea
 Florida begonia
 Florida hibiscus
 Galinsoga
 Jimson weed
 Knotweed
 Knotweed
 Lambs ears
 Pigweed
 Poinsettia
 Purslane

Ragweed
 Redweed
 Russian thistle
 Sesbania
 Shepherd's purse
 Sunflower
 Tansy
 Tansy
 Tansy
 Tansy
 Tansy
 Tansy
 Tansy

SENCOR 4 and Scepter plus Trehlan, Lasso Dual, Pro or Sonalan will provide suppression, reduce the seed bank and control weeds.

RECOMMENDED APPLICATION (BROADCAST RATES)

A Trehlan (pts/acre)
 B Dual (pts/acre)
 C Pro (pts/acre)
 D Lasso (pts/acre)
 E Sonalan (pts/acre)

Soil Texture	A	B	C	D	E	SENCOR 4 (pts/acre)	Command 4 EC (pts/acre)
COARSE							
MEDIUM							
FINE							

with a calcareous surface area of a pH of 7.5 or higher

BEST AVAILABLE COPY

having a calcareous surface area of a pH of 7.5 or higher

SENCOR PLUS SCEPTER PLUS A GRASS HERBICIDE

TANK MIX APPLICATION: SENCOR 4 may be applied with Scepter and a grass herbicide. Trehlan, Dual, Lasso or Sonalan for the control of certain broadleaf weeds and grasses in soybeans. SENCOR 4 and Scepter plus Trehlan or Sonalan may be applied preplant or preplant broadcast. SENCOR 4 and Scepter plus Lasso, Dual or Pro may be applied preplant or preplant broadcast. SENCOR 4 and Scepter plus Lasso, Dual or Pro may be applied preplant or preplant broadcast. Consult the Scepter, Trehlan, Lasso, Dual, Pro or Sonalan labels for specific directions for use, recommendations, restrictions and additional needs control.

Apply Rot... under the... section... front of the SENCOR 4 label.

Application: For specific application information, refer to the General Information section of the label.

Weeds Controlled: SENCOR 4 plus Scepter plus Trehlan, Lasso, Dual, Pro or Sonalan will control many annual broadleaf weeds and grasses.

Annual Grasses

Barnyardgrass
 Bluegrass
 Broadleaf signalgrass
 Brown top miller
 Crabgrass
 Crowfootgrass

Foxtail
 Goosegrass
 Johnsongrass (seedling)
 Parnassia
 Wildflower

23/4

Annual Broadleaves

Bristly Starwort	Prickly Soda Leafweed
Buffalobur	Purslane
Carpetweed	Ragweed (Common)
Cocklebur	Rudbeckia
Croton	Russar (Tree)
Copperhead	Sesbania
Florida Beggarweed	Shepherdspurse
Florida Pusley	Sicklepod
Gallinule	Smartweeds
Jimsonweed	Spotted Spurge
Knockweed	Spurred Anoda
Kochia	Sunflower
Lambquarters	Velvetleaf
Morningglory (vine)	Velvetleaf
Morningglory (leafy)	Wild Mustard
Mustard	

SENCOR and Linuron plus Lasso Dual Preemergence application will provide superior control of the competition of weeds and grasses in soybeans.

RECOMMENDED APPLICATION (BROADCAST RATES)

A - Trellan (1 pt/acre)	D - Lasso (1 pt/acre)
B - Dual (1 pt/acre)	E - Sonalan (1 pt/acre)
C - Penndrop (1 pt/acre)	

SOIL TEXTURE	A	B	C	D	E	SENCOR 4 (1 pt/acre)	Seep (1 pt/acre)
COARSE							
MEDIUM							
FINE							

all areas of same area (1/2 H, 1/2 S, 1/2 H)

or soil having a calcareous surface area of 1/2 H or higher

BEST AVAILABLE COPY

SENCOR PLUS LINURON PLUS (LASSO OR DUAL)

SENCOR PLUS LINURON PLUS (LASSO OR DUAL) TANK MIX APPLICATION: SENCOR 4 may be applied in combination with Linuron and Lasso 4 or Dual 8 EC as a preemergence application for the control of certain weeds in soybeans. Consult the Linuron, Lasso 4 or Dual 8 EC label for specific directions for use, recommendations, restrictions and any additional weeds not specified on this label.

Mixing: Refer to Mixing under the General Information section in the front of this label.

Application: Applications are to be made only with ground spray equipment in accordance with specified soil types and broadcast rates. For specific application information, refer to the General Information section in the front of this label.

BROADCAST RATES: (0.5 to 3% Organic Matter Only)

SOIL TEXTURE	Linuron 50 WP or 4 C		Lasso 4 or Dual 8 EC	
	SENCOR 4 (1 pt/acre)	(1 pt/acre) or (1 pt/acre)	(1 pt/acre)	(1 pt/acre)
COARSE				
MEDIUM				
FINE				

1. Soil texture may vary from coarse to fine. For example, a soil may be classified as medium textured in some regions of the U.S.

SPECIAL PRECAUTIONS (SENCOR PLUS LINURON PLUS (LASSO OR DUAL): For additional precautions, restrictions, limitations, and sprayer clean-up information, refer to the appropriate sections of this label and the Linuron label and the Lasso 4 or Dual 8 EC labels.

**FOR USE IN
COARSE (LIGHT) SOILS:**

In Alabama Arkansas Florida Georgia Louisiana Mississippi
Missouri North Carolina Oklahoma South Carolina Tennessee
Texas and Virginia

Texas and Virginia

Chief of the North American Commission on the Status of Women, Mrs. Judith L. Linn, said today that the American people are "not ready to take a step which might result in the loss of certain women's personal freedom to the degree which is embodied in this legislation," the Equal Rights Amendment. Mrs. Linn said that Americans believe in the right of women to be free to contribute to the nation's production of goods and services, but that they are not ready to "take the step."

Mixing: Refer to Mixing section for correct mixing instructions in the front of this label.

Application If a person is applying for a job, the employer should be informed of the person's criminal record. The person should be informed of the results of the background check.

SENCOR (ALONE) PREEMERGENCE APPLICATION
BROADCAST RATES

BROADCAST RATES		SENCOR 4
SOIL TEXTURE	Organic Matter	1615 acres
COARSE LIGHT SOILS		

SENCOR IN COMBINATION WITH OTHER HERBICIDES

SENSOR IN COMBINATION WITH OTHER HERBICIDES
 SENOR OR 4 is recommended in a tank mix with the following herbicides:
 Treflan as a pre-emergent or post-emergent application for control of
 underground and aerial weeds in low rice production areas; and
 application of Treflan and SENOR 4 as a pre-emergent application for
 a pre-emergent application in low rice production areas; and
 application of Treflan and SENOR 4 as a pre-emergent application in low rice production areas.

BEST AVAILABLE COPY

BROADCAST RATES

BROADCAST RATES		Plus Sewer & lots acre
SOIL TEXTURE	Organic Matter	Combination Product Acre
COARSE LIGHT SOILS		Preplant Incorporated
		Preemergence

SPECIAL PRECAUTIONS: In addition to the instructions on the label, and the instructions on the label, the following instructions should be followed:

Do not use on sands with less than 10% fines or on
sandy loams or silty sand soils with less than 10% fines.

**FOR CONTACT ACTION
AND RESIDUAL CONTROL:**

For Control of Emerged Annual Broadleaf Weeds and Grasses and For Suppression of Emerged Perennials Where Soybeans Will Be Planted Directly Into a Pre-formed Bed, Cover Crop or in Previous Crop Residues as in "Minimum Till" or "No Till" Crop Production.

SENCOR PLUS GRAMOXONE

SENCOR PLUS GRAMOXONE TANK MIX APPLICATION: Apply 1 to 2 pints of Sencor plus 4 to 28 pints of Gramoxone according to the recommendations for SENCOR. Apply late as a broadcast ground spray before or during a pre-planting but before emergence of the crop. Sny means exposed at time of application will be killed. Use the higher rate of Gramoxone for heavier infestations or for taller weeds (4 to 6 inches tall). Apply in 20 to 60 gallons spray mix per acre. Use the higher volume in fields with heavy weed infestations or with large amounts of previous crop residues. Add Ortho X-77 Spreader or similar non-ionic spreader at 1 quarter per 100 gallons of diluted spray. This mixture

SENCOR PLUS LASSO AND GRAMOXONE

SENCOR PLUS LASSO AND GRAMOXONE TANK MIX APPLICATION: Apply 1 to 2 1/2 pints of Gramoxone Super plus 2 to 2 1/2 quarts Lasso 4 L.V. plus 1/2 to 2 pints SENCOR 4 as a pre-emergence broadcast ground spray. Use the higher rate of Gramoxone for weeds 4 to 6 inches tall. Soybeans exposed at time of application will be killed. Refer to the appropriate sections of the labels on the Gramoxone and Lasso labels for rates which are dependent on soil textures, directions, limitations, and cautions.

Apply up to 60 gallons spray mix per acre. Use the higher rate of Lasso with heavy weed infestations, or with large amounts of previous crop residues. Add Ortho X-77 Spreader or similar machine spreader at 8 fl. ounces per 100 gallons of diluted spray. Add Ethion-OR 4 to the spray tank first and mix thoroughly. Then add Lasso, followed by Gramoxone, and X-77 Spreader. Christianization is recommended at all times. This tank mix will not control shattercane which emerges after treatment of the weeds taller than 6 inches.

SENCOR PLUS SURFLAN AND GRAMOXONE

SENCOR PLUS SURFLAN AND GRAMOXONE TANK MIX APPLICATION: This combination provides control of weeds and grasses which are emergent at the time of application plus many weeds and grasses which may germinate after planting. Soybeans exposed at time of application will be killed.

Weeds Controlled by Gramoxone: Gramoxone will kill emerged annual broadleaf weeds and grasses and provide top kill of perennials.

When to Apply: Apply the tank mix at planting or within 10 days after planting. Do not use this combination after mid-July.

SENCOR PLUS SURFLAN AND GRAMOXONE PREEMERGENCE TANK MIX APPLICATION:

BROADCAST RATES Per Acre

	SENCOR 4	Surflan A.S.	Gramoxone Super
COARSE			
MEDIUM			
FINE			

BEST AVAILABLE COPY

Mixing note: Mix up the General Information section of the label of this label.

Apply mix with a broadcast or line spreader at 8 fl. ozs. per 100 gallons of diluted spray mixture.

Application: For specific application information refer to the General Information section in the front of this label.

A one-half inch rain or more or its equivalent in overhead irrigation is needed to activate the herbicide combination.

SPECIAL PRECAUTIONS (SENCOR plus Surflan and Gramoxone): Poor weed control and/or crop injury may result if directions are not followed.

Do not use a tank mix product on your soil if planter or crop injury may result.

For additional information on special precautions, restrictions, and other information refer to the appropriate sections of the Lasso 4 L.V., Surflan and Gramoxone labels.

SENCOR PLUS 2,4-D LOW VOLATILE ESTER

SENCOR 4 as an herbicide combination with Dual as a soil sterilant. Prowl may be applied with 2,4-D low volatile ester as a pre-plant soil application for the control of certain types of weeds and grasses in soybeans or corn prior to planting. Apply 2,4-D low volatile ingredient at a rate of 1/4 lb. low volatile ester with label rates of SENCOR 4. When a grass herbicide is used in the tank mix, apply the rates specified on that product's label. Apply Dual as recommended 30 days prior to planting.

Refer to Ethion-OR 4 2,4-D low volatile ester, Dual, Lasso, Surflan, and other labels for weeds controlled, precautions, restrictions, and sprayer clean-up information.

POSTEMERGENCE DIRECTED SPRAY APPLICATIONS

Weeds Controlled: SE NC 44, a broad spectrum herbicide to soybeans is a true herbicide according to the label. It is a pre-emergent herbicide that is applied to the soil. It is a pre-emergent herbicide that is applied to the soil. It is a pre-emergent herbicide that is applied to the soil.

[illegible][illegible]

BEST AVAILABLE COPY

RECOMMENDED APPLICATIONS

CRUISE	SENSOR 2 BIT AREA	REMARKS
1	1	1
2	2	2
3	3	3
4	4	4
5	5	5
6	6	6
7	7	7
8	8	8
9	9	9
10	10	10
11	11	11
12	12	12
13	13	13
14	14	14
15	15	15
16	16	16
17	17	17
18	18	18
19	19	19
20	20	20
21	21	21
22	22	22
23	23	23
24	24	24
25	25	25
26	26	26
27	27	27
28	28	28
29	29	29
30	30	30
31	31	31
32	32	32
33	33	33
34	34	34
35	35	35
36	36	36
37	37	37
38	38	38
39	39	39
40	40	40
41	41	41
42	42	42
43	43	43
44	44	44
45	45	45
46	46	46
47	47	47
48	48	48
49	49	49
50	50	50
51	51	51
52	52	52
53	53	53
54	54	54
55	55	55
56	56	56
57	57	57
58	58	58
59	59	59
60	60	60
61	61	61
62	62	62
63	63	63
64	64	64
65	65	65
66	66	66
67	67	67
68	68	68
69	69	69
70	70	70
71	71	71
72	72	72
73	73	73
74	74	74
75	75	75
76	76	76
77	77	77
78	78	78
79	79	79
80	80	80
81	81	81
82	82	82
83	83	83
84	84	84
85	85	85
86	86	86
87	87	87
88	88	88
89	89	89
90	90	90
91	91	91
92	92	92
93	93	93
94	94	94
95	95	95
96	96	96
97	97	97
98	98	98
99	99	99
100	100	100

SPECIAL PRECAUTIONS (Directed Postemergence) (1)

SPECIAL PRECAUTIONS (Directed Postemergence) Do not spray directly on or near water. Do not apply to plants under stress. Do not apply to plants with frost damage. Do not apply to plants with insecticide injury. Do not apply to plants with herbicide injury. Do not apply to plants with nutrient deficiency. Do not apply to plants with disease. Do not apply to plants with other stresses. Do not apply to plants with sensitive soybean varieties. See Special Precautions label for full details of this label.

To avoid injury to other crops or desirable plants from spray drift, sprayer pressure must not exceed 30 psi and the sprayer must be fitted with nozzles no smaller than B002 Jet (or equivalent). Do not apply under adverse conditions which favor drift.

27/450

[illegible]

Aerial Application

Weeds Controlled: 100% of the weeds were controlled.

[illegible][illegible]

For Control of Wild Mustard (*Brassica* spp.) in Potatoes (Except California): Apply at the rate of 1/2 to 1 pint per acre as a pre-emergence broadcast spray using aerial or ground spray equipment in accordance with directions on this label. Application should be made at or near ground crack and/or before the mustard seedlings emerge. For control of mustard larger than 2 inches in height, other weeds listed on the label a minimum of 1 pint per acre must be used. When less than 1 pint per acre is used, the herbicide should be applied in accordance with directions on the label for control of other weeds or later germinating weeds.

Hard To Control Weeds Amongst SENCOR 4 applications, you can expect to provide commercially acceptable control of many instances of the following weeds and grasses when mixed with other plants:

Barryman, J. S. 1954. The
Barryman, J. S. 1954. The
Barryman, J. S. 1954. The
Barryman, J. S. 1954. The

Weeds Controlled SE 1400-4 applied early postemergence on 10/21/45 according to directions. Will effectively control

Cocklebur	Ragweed Common
(Xanthium	(Ambrosia artemisiifolia)
pennsylvanicum)	Russian thistle
embryophyllum)	(Salsola kali)
(Chenopodium album)	Smartweed Pennsylvania
Ragweed Redroot	(Polygonum
(Amaranthus retroflexus)	pennsylvanicum)

Fortails
(*Setaria* spp.)

BEST AVAILABLE COPY

28

3

(2) 对数函数

Journal of Management Education 30(6)

IBLE COPY

CROP	SCHEDULE DATE, TIME	REMARKS
		PREFEMERGENCE
		POSTEMERGENCE
		SPLIT APPLICATIONS
		IBAMA OREGON AND WASHINGTON ONLY

SPRINKLER IRRIGATION APPLICATION TO POTATOES
 SENCOR 4 may be applied early postemergence to potatoes through center pivot, permanent solid set or self propelled lateral move irrigation systems as directed on this label. Refer to "Chemigation" section of this label for directions.

[illegible]

SENCOR PLUS DUAL TANK MIX APPLICATION (EXCEPT CALIFORNIA) SENCOR OR 4 may be applied in a tank mix application with Dual 5B as a pre-emergent or post-emergent application. The tank mix application of SENCOR OR 4 and Dual 5B should be made at the rate of 1.0 to 1.5 pounds per acre according to the respective label directions. The tank mix should be applied in the early pre-emergent or post-emergent period. Refer to the appropriate label for the use of SENCOR OR 4 and Dual 5B in tank mix applications.

SENCOR PLUS PROWL TANK MIX APPLICATION (EXCEPT CALIFORNIA) - SENCOR PLUS is the superior tank mix for control of Prowl as a pre-emergent herbicide and as a post-emergent. Apply SENCOR PLUS at 1.5-2.0 pounds and follow the label directions according to the respective abscissa on the treatment and control patterns. Refer to the appropriate sections of this label and the SENCOR PLUS Precautionary Statement for further information on use and control of Prowl.

SPECIAL PRECAUTIONS (Poisons)

Do Not use SENCOR 4 on Potatoes in Kern County, California

Do not apply within 24 hours of application of other pesticides.

[illegible]

• Do not apply SE NCOA to seeds of various
• Do not use seeds from a previous growing season
• and for seeds during the next growing season (1st and 2nd
• Do not SE NCOA on any other crop

BEST AVAILABLE COPY

SENCOR herbicide is labelled for use in alfalfa and saint john in the following areas

1. *Aristida* and *saintpalm*, including mixed stands of grasses in areas except California
2. *Aristida* and *saintpalm*, including mixed stands with grasses in California only
3. *Aristida* - Tank Mix combination with Grain, except California, Idaho, Montana, Nevada, Oregon, Utah, Washington, and Wyoming
4. *Aristida* only - Restricted Application of SF for other than: California, Dry, Forage or for other than: Idaho, Oregon, Washington, Kentucky, Michigan, Minnesota, Nebraska, New York, Ohio, Pennsylvania, South Dakota, and Wisconsin only.

SEACON 4 is recommended for use in agricultural, forestry, and ornamental plant and tree pest control. It is also suitable for use in the control of certain grass and broadleaf weeds.

APPLICATION: Refer to "General Information" in the front of this label for detailed information on the application of the SENCOR 4. For information on applying SENCOR 4 in fluid or dry form, refer to the "Application of SENCOR 4 in Fluid Fertilizers," or "Application of SENCOR 4 in Dry Fertilizers," or "Application of SENCOR 4 in Dry Bulk Fertilizer" under the "General Information" section in the front of this label.

SPECIAL PRECAUTIONS: Do not apply SENCOR if after growth begins in the spring or before growth ceases in the fall. Treat only dormant established alfalfa (except where specified on this label).

Do not graze or harvest within 28 days after application.
For best weed control, apply SENCOR 4 when weeds are less than 2 inches tall or before weeds foliage is 2 inches in diameter.
Reduced weed control may occur when extended dry conditions follow application of SENCOR 4.

Weeds Partially Controlled: At the rate of 2 pints/acre SENCOR 4 may be used to reduce the competition from curly dock (*Rumex crispus*). At 1 to 2 pints/acre SENCOR 4 may be used to reduce the competition of German moss or knot (*Sclanthus annuus*).

31

ALFALFA AND SAINFOIN

(California Only)
(Including Mixed Stands With Grasses)

SENCOR 4 is recommended for use in aerial or ground spray equipment as a broadcast surface application to dormant established crops of alfalfa and sainfoin.

APPLICATION: SENCOR 4 is recommended for use in aerial or ground spray equipment as a broadcast surface application to dormant established crops of alfalfa and sainfoin for control of certain grass and broadleaf weeds. Do not apply SENCOR 4 after growth begins in the spring or before growth ceases in the fall. Do not apply to either alfalfa or sainfoin during the first growing season after seeding.

For information on applying SENCOR 4 in fluid fertilizer form, refer to alfalfa reference section of this label.

For information on SENCOR 4 application in tank application of SENCOR 4 and fluid fertilizer, refer to the appropriate section of this label.

Weeds Controlled

1/4 to 1 pint SENCOR 4/acre

Chickweed, Common

1 to 2 pints SENCOR 4/acre

Broadleaves

Chickweed, Common

(*Cerastium vulgatum*)

Fieldweed

(*Oxycorymbis sophia*)

Fieldbit

(*Lamium amplexicaule*)

Kudzu

(*Rhynchospora*)

Mustard, Black

(*Brassica nigra*)

Grasses

(*Brachypodium*)

(*Brachypodium*)

Mustard, Tansy

(*Oxycorymbis pinnata*)

Pepperweed, Virginia

(*Lepidium virginicum*)

Shepherdspurse

(*Caprilla bursa pastoris*)

White Fleckle

(*Myriophyllum*)

White Blackweed

(*Myriophyllum*)

White Blackweed

(*Myriophyllum*)

White Blackweed

(*Myriophyllum*)

2 pints SENCOR 4/acre

Broadleaves

Dandelion

(*Taraxacum officinale*)

Grasses

Barnyardgrass

(*Echinochloa crusgalli*)

Bluegrass

(*Poa annua*)

Footail Barley

(*Hordeum jubatum*)

RECOMMENDED BROADCAST APPLICATION:

CROP	SENCOR 4 (pints/acre)	REMARKS
ALFALFA	1/4 to 1 pint	Apply the proper dosage according to the weeds to be controlled. Apply SENCOR 4 in aerial or ground spray equipment as a broadcast surface application to dormant established crops of alfalfa and sainfoin for control of certain grass and broadleaf weeds. Do not apply SENCOR 4 after growth begins in the spring or before growth ceases in the fall. Do not apply to either alfalfa or sainfoin during the first growing season after seeding.
SAINFOIN	1/4 to 1 pint	Apply the proper dosage according to the weeds to be controlled. Apply SENCOR 4 in aerial or ground spray equipment as a broadcast surface application to dormant established crops of alfalfa and sainfoin for control of certain grass and broadleaf weeds. Do not apply SENCOR 4 after growth begins in the spring or before growth ceases in the fall. Do not apply to either alfalfa or sainfoin during the first growing season after seeding.

FOR USE ON MIXED STANDS OF ALFALFA AND GRASSES: Apply the proper dosage according to the weeds to be controlled. Apply SENCOR 4 in aerial or ground spray equipment as a broadcast surface application to dormant established crops of alfalfa and sainfoin for control of certain grass and broadleaf weeds. Do not apply SENCOR 4 after growth begins in the spring or before growth ceases in the fall. Do not apply to either alfalfa or sainfoin during the first growing season after seeding.

Apply SENCOR 4 in aerial or ground spray equipment as a broadcast surface application to dormant established crops of alfalfa and sainfoin for control of certain grass and broadleaf weeds. Do not apply SENCOR 4 after growth begins in the spring or before growth ceases in the fall. Do not apply to either alfalfa or sainfoin during the first growing season after seeding.

ALFALFA - TANK MIX WITH GRAMOXONE

(Colorado, Idaho, Montana, Nevada, Oregon, Utah, Washington, and Wyoming)

APPLICATION: SENCOR 4 plus Gramoxone tank mix application is recommended for use, during the dormant season in aerial or ground spray equipment as a broadcast surface application to established (at least 1 year old) alfalfa for the control of certain

BEST AVAILABLE COPY

grass and broadleaf weeds. Do not apply SENCOR or Gramoxone tank mix to regrowth (after grazing or cutting) that is more than 1 1/2 inches tall. Apply once per season. Do not apply to alfalfa pastures during growing season. Use a minimum of 10 gallons of water per acre with aerial spray or 20 gallons per acre with ground spray. Do not apply to alfalfa pastures with a pH greater than 8.2. Do not apply to alfalfa pastures with a pH greater than 8.2. Do not apply to alfalfa pastures with a pH greater than 8.2.

WEEDS CONTROLLED — SENCOR 4 controls the following weeds: Annual weeds, biennial weeds, perennial weeds, and broadleaf weeds. SENCOR 4 controls the following weeds: Annual weeds, biennial weeds, perennial weeds, and broadleaf weeds.

Annual weeds	Chickweed
Biennial weeds	Downy Brome
Perennial weeds	Japanese Brome
Broadleaf weeds	Sheep Sorrel

SENCOR 4 controls the following weeds: Annual weeds, biennial weeds, perennial weeds, and broadleaf weeds. SENCOR 4 controls the following weeds: Annual weeds, biennial weeds, perennial weeds, and broadleaf weeds.

SENCOR 4 controls the following weeds: Annual weeds, biennial weeds, perennial weeds, and broadleaf weeds. SENCOR 4 controls the following weeds: Annual weeds, biennial weeds, perennial weeds, and broadleaf weeds.

Annual weeds	Yellow Wildweed
Biennial weeds	White Clover
Perennial weeds	White Clover
Broadleaf weeds	Pink Clover
Annual weeds	Blue Grass
Biennial weeds	Red Clover
Perennial weeds	Common Brome
Broadleaf weeds	Marestail
Annual weeds	Field Bindweed
Biennial weeds	Rough Brome
Perennial weeds	
Broadleaf weeds	

BEST AVAILABLE COPY

RECOMMENDED APPLICATIONS

SENCOR 4
1 to 1 1/2 pints
plus
Gramoxone Super
2 to 2 1/2 pints

Apply specified dosages of SENCOR and Gramoxone Super at least 10 gallons of water per acre with aerial equipment or at least 20 gallons of water per acre with ground equipment. Do not apply this tank mix to alfalfa regrowth (after grazing or cutting) if more than 2 inches tall. For best weed control, apply when broadleaf weeds and grasses are 1 to 6 inches tall and are actively growing. Care should be taken to avoid overlap. Use lower rates of SENCOR on sandy loam or loamy sand soils. Reduced weed control may occur when extended dry conditions follow application of SENCOR. Crop injury may occur if alfalfa is under stress conditions such as diseases, insect infestations, drought or winter injury or if SENCOR is applied to alfalfa earlier than 12 months after seeding.

FOR USE ON MIXED STANDS OF ALFALFA AND GRASSES
Rates of 1 to 1 1/2 pints of SENCOR 4 per acre will provide partial reduction of forage grass stands. These rates may be used to reduce forage grass stands to prevent crowding out of alfalfa.

Do not graze or harvest within 42 days after application. In areas west of the Rockies, avoid the use of SENCOR 4 on soils with calcareous surface soils with high levels of lime, and with a pH greater than 8.2.

Do not apply when weather conditions favor spray drift. Aerial application should not be made when wind speed is greater than 10 mph.

Do not use on alfalfa and soybeans. Refer to the Gramoxone label for additional directions, weeds species controlled and precautions.

ALFALFA

Post Dormant Application of SENCOR Impregnated on Dry Fertilizer Only — SENCOR 4 may be applied after dormancy has broken, but prior to three inches of new alfalfa shoot growth, only when impregnated on dry fertilizer in Connecticut, Illinois, Indiana, Iowa, Kansas, Kentucky, Michigan, Minnesota, Nebraska, New York, Ohio, Pennsylvania, South Dakota and Wisconsin. Apply at rates of 1 1/2 to 2 pints per acre as directed on the label for application during dormancy. Apply only when alfalfa foliage is dry or crop injury may occur. When using this application method, do not harvest or graze treated alfalfa for 60 days after application.

ASPARAGUS

(Established)

[illegible]

Weeds Controlled 14. 100% control of all weeds.

Broadleaves

[illegible]

Grasses

1. 1990年12月29日，全国人大常委会通过了《中华人民共和国香港特别行政区基本法》。

BEST AVAILABLE COPY

RECOMMENDED BROADCAST APPLICATIONS

CROP	SENCOR 4 1015 acre.	REMARKS
		PREEMERGENCE APPLICATION ONLY
		SPLIT APPLICATION: PREEMERGENCE AND POST HARVEST
		Preemergence Application
		Post Harvest Application

(IMPORTANT)

SPECIAL PRECAUTIONS (Asparagus) Do not use on lawns or in areas where children or pets may be exposed. Do not use on lawns or in areas where children or pets may be exposed.

APPLY POST HARBEST APRIL 15TH 1941
AFTER THE LAST HARVEST 1 SEP 41

[illegible]

(Iowa, Kansas, Missouri, Nebraska, and South Dakota only)

SENCOR herbicide is recommended for the added effect on the control of certain broadleaf weeds in corn when applied pre-emergence in a tank mix combination with labeled rates of Lasso plus atrazine, Lasso plus Bladex[®], Dual plus atrazine or Dual plus Bladex. Refer to the appropriate product labels for weeds controlled by these combinations.

in wet or cloudy weather or crop injury may occur.
Do not apply within 24 hours of treatment with other pesticides.

SPECIAL PRECAUTIONS (all applications) Crop injury may result if applied to the following crops: corn, sorghum, cotton, soybeans, alfalfa, clover, vetch, and other legumes. Do not apply to citrus, grapes, or other fruiting plants. Do not apply to ornamentals, shrubs, or trees. Do not apply to aquatic plants or animals.

Do not apply to crops or areas where livestock or poultry may be present. Do not apply to areas where children or pets may be present. Do not apply to areas where bees or other beneficial insects may be present. Do not apply to areas where fish or other aquatic life may be present. Do not apply to areas where birds or other wildlife may be present.

NOTE SENCOR IR 4 is a selective herbicide. It will not control weeds that are resistant to its active ingredient. It will not control weeds that are in the seedling stage. It will not control weeds that are in the flowering stage. It will not control weeds that are in the fruiting stage. It will not control weeds that are in the seedling stage.

SUGARCANE

(Hawaii Only)
SENCOR IR 4 is a selective herbicide. It will not control weeds that are resistant to its active ingredient. It will not control weeds that are in the seedling stage. It will not control weeds that are in the flowering stage. It will not control weeds that are in the fruiting stage. It will not control weeds that are in the seedling stage.

Ground Application SENCOR IR 4 is a selective herbicide. It will not control weeds that are resistant to its active ingredient. It will not control weeds that are in the seedling stage. It will not control weeds that are in the flowering stage. It will not control weeds that are in the fruiting stage. It will not control weeds that are in the seedling stage.

BEST AVAILABLE COPY

Aerial Application SENCOR IR 4 is a selective herbicide. It will not control weeds that are resistant to its active ingredient. It will not control weeds that are in the seedling stage. It will not control weeds that are in the flowering stage. It will not control weeds that are in the fruiting stage. It will not control weeds that are in the seedling stage.

SENCOR IR 4 applied preemergence or postemergence to the sugarcane as a broadcast spray or spot treatment will effectively control the following when weeds are less than 3 inches in height.

Weeds Controlled in Irrigated and Non-irrigated Sugarcane

Broadleaves

- | | |
|---|--|
| Amaranth, Spiny
(<i>Amaranthus spinosus</i>) | Florida paintbrush
(<i>Erigeron annuus</i>) |
| Euphorbia, Wild
(<i>Euphorbia corollata</i>) | Spurge, Garden
(<i>Euphorbia hibernica</i>) |
| Erigeron
(<i>Erigeron phillyriaefolius</i>) | Spurge, Gravel
(<i>Euphorbia corollata</i>) |

Grasses

- | | |
|--|---------------------------------------|
| Crabgrass
(<i>Digitaria sp.</i>) | Ricegrass
(<i>Oryzopsis sp.</i>) |
| Johnsongrass
(<i>Panicum maximum</i>) | Wild grass
(<i>Echinochloa</i>) |
| Pushgrass
(<i>Eleusine indica</i>) | |

Weeds Controlled in Irrigated Sugarcane Only

Broadleaves

- | | |
|---|--|
| Amaranth, Spiny
(<i>Amaranthus spinosus</i>) | Florida paintbrush
(<i>Erigeron annuus</i>) |
| Euphorbia, Wild
(<i>Euphorbia corollata</i>) | Spurge, Garden
(<i>Euphorbia hibernica</i>) |
| Erigeron
(<i>Erigeron phillyriaefolius</i>) | Spurge, Gravel
(<i>Euphorbia corollata</i>) |

Grasses

- | | |
|--|--|
| Alexandergrass
(<i>Bracharia plantaginea</i>) | Bristly foxtail
(<i>Setaria verticillata</i>) |
|--|--|

Weeds Controlled in Non-irrigated Sugarcane Only

Broadleaves

- | | |
|---|--|
| Ageratum (<i>Ageratum conyzoides</i>) | |
| Richardia (<i>Richardia brasiliensis</i>) | |
| Tarweed (<i>Cuphea carthagenensis</i>) | |

RECOMMENDED BROADCAST APPLICATIONS

CROP	SENCOR 4 (pts/acre)	REMARKS
		PREEMERGENCE
		OR
		EARLY POSTEMERGENCE
		OR
		POSTEMERGENCE
		SPOT TREATMENT

Note

Special Precaution

BEST AVAILABLE COPY

SUGARCANE

(Louisiana and Texas Only)

Preemergence and postemergence applications of SENCOR 4 with aerial or ground spray equipment are recommended for control of the following weeds in sugarcane in Louisiana and Texas.

Broadleaves

Amaranth Spiny
(*Amaranthus spinosus*)
Bindweed Field
(*Convolvulus arvensis*)
Chickweed
(*Cerastium vulgatum*)
Henbit
(*Lamium amplexicaule*)
Lambquarters
(*Chenopodium album*)
London rocket
(*Sisymbrium irio*)

Marestail
(*Conyza canadensis*)
Stardust
(*Rhassia alba*)
Lewy
(*Amaranthus*)
Ragweed
(*F. multifida*)
Amaranth
(*Amaranth*)

Grasses

Broadleaf Signalgrass
(*Bracharia platyphylla*)
Crabgrass
(*Digitaria sp.*)
Foxtails
(*Setaria sp.*)

St. Augustine
(*Syntherisma tenax*)
St. Andrew
(*Amorpha*)

RECOMMENDED BROADCAST APPLICATIONS

CROP	SENCOR 4 (pts/acre)	REMARKS
Sugarcane Louisiana Texas Only		BROADCAST Apply SENCOR 4 at 1.5 to 2.0 pts/acre in early spring before weeds emerge. Apply with ground or aerial equipment. Make a second application in late spring. BAND Apply SENCOR 4 at 1.5 to 2.0 pts/acre in early spring before weeds emerge. Apply with ground or aerial equipment. Make a second application in late spring.

37/40

SPECIAL PRECAUTIONS (Louisiana and Texas only) Do not use treated forage for feed or forage.
Use the higher rate on heavy clay soil and soil with high phosphate. Forage matter. If necessary a third application may be applied at 14 day intervals. Do not apply within 14 days of harvest.

SUGARCANE

(Florida Only)

1. Apply generally with the top directed spray at 1.0 to 1.5 lb/A. OR 4.0 lb/A is recommended for the control of the following weeds in sugarcane fields:

Broadleaves

Amaranthus spp.
Amaranthus spp.
Portulaca
Portulaca
Portulaca
Portulaca
Portulaca
Portulaca
Portulaca
Portulaca

Grasses

Cynodon
Digitaria
Eragrostis
Eragrostis
Eragrostis
Eragrostis
Eragrostis
Eragrostis
Eragrostis
Eragrostis

RECOMMENDED BROADCAST APPLICATIONS

CROP	SENCOR 4 lb/A	REMARKS
		GROUND APPLICATION
		POSTEMERGENCE BROADCAST OR HAND
		POSTEMERGENCE DIRECTED SPRAY
		AERIAL APPLICATION

BEST AVAILABLE COPY

SENCOR PLUS Atrazine TANK MIX 1.0 to 1.5 lb/A. Apply with the top directed spray at 1.0 to 1.5 lb/A. OR 4.0 lb/A is recommended for the control of the following weeds in sugarcane fields:
Amaranthus spp.
Amaranthus spp.
Portulaca
Portulaca
Portulaca
Portulaca
Portulaca
Portulaca
Portulaca
Portulaca

Special Precautions (Florida Only) Do not use.
Spray contact with sugarcane foliage may result in leaf margin chlorosis and necrosis.
Do not apply within 60 days of harvest.
Do not use treated crop for feed or forage.
Avoid spray overwind variations in application speed that may result in insufficient or excessive rates of application.

TOMATOES

Apply SENCOR tank mix with ground or aerial spray at 1.0 to 1.5 lb/A. OR 4.0 lb/A is recommended for the control of the following weeds in tomato fields:
Amaranthus spp.
Amaranthus spp.
Portulaca
Portulaca
Portulaca
Portulaca
Portulaca
Portulaca
Portulaca
Portulaca

Weeds Controlled

PREPLANT INCORPORATED APPLICATIONS - TRANSPLANT TOMATOES ONLY

Broadleaves

Galinsoga
Galinsoga spp.
Amaranthus
Amaranthus spp.

Pigweed Redroot
Amaranthus retroflexus
Purslane (Common)
Portulaca oleracea

Grasses

Eragrostis
Eragrostis spp.

1954

40/42

RECOMMENDED END-USES			REMARKS
SOIL TEXTURE	SEMLOH 2 gpts. clay	Organic Matter	
LOADING	1 to 2	Over 2	
MEDIUM			
FINE			

Broadleaves

THE UNIVERSITY OF CHICAGO
CHICAGO, ILLINOIS 60637
U.S.A.
TEL: 773-936-8000
FAX: 773-936-8000
WWW.CHICAGOEDUCATION.COM

1. *Chlorophyll a* and *Chlorophyll b* contents were determined using a spectrophotometer (Shimadzu UV-160U) at 663 nm and 646 nm, respectively. The total chlorophyll content was calculated using the following formula: $\text{Total Chlorophyll} = \frac{12.7 \times \text{Absorbance at 663 nm} + 22.9 \times \text{Absorbance at 646 nm}}{1000}$.

NOTE

BEST AVAILABLE COPY

SENCOR in Combination with Other Herbicides SENCOR is registered for use in combination with other herbicides for control of weeds in corn, sorghum, soybeans, cotton, and sugarcane. For more information, see the label for the herbicide with which SENCOR is being used. SENCOR is also registered for use in combination with other herbicides for control of weeds in rice. For more information, see the label for the herbicide with which SENCOR is being used.

WINTER WHEAT

FOR USE ON WINTER WHEAT IN IDAHO, OREGON, UTAH
AND WASHINGTON ON THE VARIETIES DAWNS, GAINES, HAT
TON, HILBILBS, OF LUKE, McDERMID, MORO, NUGAINES,
PAPA, RAK, SPRAGUE, STEPHENS, WANSER, FARGO, TYEE
AND WESTON AND IN MONTANA ON THE VARIETIES CEN
TURY, CHEYENNE AND WINALTA.

A large percentage of the vegetation is composed of grass and broadleaf shrubs.

RECOMMENDED BROADCAST APPLICATIONS*			
	SENSOR 4 (pts/acre)		
SOIL TEXTURE	0.75% to 1% Organic Matter	1% to 2% Organic Matter	More than 2% Organic Matter
COARSE	100	100	100
MEDIUM	100	100	100
FINE	100	100	100

*Based on a soil analysis which requires that 1% organic matter be present in the soil. Soil with less than 1% organic matter may be classified as the "lowest" soil texture category.

42/46

FOREIGN OR CONDOMINIUM WATER AND POWER, OKLAHOMA AND TEXAS ON THE WESTERN SHORE OF LAKE HAWK LARRED NEWTON PIONEER DISTRICT, TEXAS WITH TAKING VICTOR AND GRAY, ALL

Broadleaves

Grasses

Hard to Control Needs

BEST AVAILABLE COPY

SENCOR 4
1p15 acres

SENSOR in Combination with Other Herbicides SENSOR may be applied as directed above or may be tank mixed with 2,4-D, Barvel, Bromox, Bromoxynil, or Butyl. Refer to the other product label registered for a full and complete label and weed species or crop.

(Arkansas, Louisiana, Mississippi and Tennessee)

Weeds Controlled:

Broadleaves

Bittercress
(*Cardamine* spp.)
*Buttercup
(*Ranunculus* spp.)
Carolina Geranium
(*Geranium carolinianum*)
Chickweed - Common
(*Stellaria media*)
Chickweed - Mouseear
(*Cerastium vulgatum*)
Cutleaf Evening Primrose
(*Oenothera laciniata*)
Henbit
(*Lamium amplexicaule*)
*Kochia
(*Kochia scorpiaria*)

Lambsquarters - Common
(*Chenopodium album*)
Mayweed - Dog Fenel
(*Eupatorium capillifolium*)
Mustard - Wild
(*Brassica kaber*)
Pepperweed - Virginia
(*Lepidium virginicum*)
Pigweed
(*Amaranthus* spp.)
Shepherdspurse
(*Capsella bursa-pastoris*)
Wild Turnip
(*Brassica rapa*)

Grasses

Bluegrass - Annual
(*Poa annua*)
*Cheatgrass
(*Bromus secalinus*)
*Downy Brome
(*Bromus tectorum*)

Little Bale
(*Hordeum pusillum*)
*Horsegrass
(*Bromus inermis*)
*Ryegrass - Annual
(*Lolium multiflorum*)
Vernal Wildgrass
(*Drababerna* spp.)

NOTE: Control of wild buckwheat (*Polygonum convolvulus*) and winter vetch (*Vicia* spp.) may be variable with SENCOR 4, ranging from poor to excellent depending on growth at time of application, temperatures and moisture conditions following application.

Recommended for use only on the following wheat varieties: Abe, Arthur 71, Caldwell, Coker 117, Coker 120, Coker 68, Doublecross, Hunter, M. 1003, M. 1004, M. 1005, M. 1006, M. 1007, M. 1008, M. 1009, M. 1010, M. 1011, M. 1012, M. 1013, M. 1014, M. 1015, M. 1016, M. 1017, M. 1018, M. 1019, M. 1020, M. 1021, M. 1022, M. 1023, M. 1024, M. 1025, M. 1026, M. 1027, M. 1028, M. 1029, M. 1030, M. 1031, M. 1032, M. 1033, M. 1034, M. 1035, M. 1036, M. 1037, M. 1038, M. 1039, M. 1040, M. 1041, M. 1042, M. 1043, M. 1044, M. 1045, M. 1046, M. 1047, M. 1048, M. 1049, M. 1050, M. 1051, M. 1052, M. 1053, M. 1054, M. 1055, M. 1056, M. 1057, M. 1058, M. 1059, M. 1060, M. 1061, M. 1062, M. 1063, M. 1064, M. 1065, M. 1066, M. 1067, M. 1068, M. 1069, M. 1070, M. 1071, M. 1072, M. 1073, M. 1074, M. 1075, M. 1076, M. 1077, M. 1078, M. 1079, M. 1080, M. 1081, M. 1082, M. 1083, M. 1084, M. 1085, M. 1086, M. 1087, M. 1088, M. 1089, M. 1090, M. 1091, M. 1092, M. 1093, M. 1094, M. 1095, M. 1096, M. 1097, M. 1098, M. 1099, M. 1100, M. 1101, M. 1102, M. 1103, M. 1104, M. 1105, M. 1106, M. 1107, M. 1108, M. 1109, M. 1110, M. 1111, M. 1112, M. 1113, M. 1114, M. 1115, M. 1116, M. 1117, M. 1118, M. 1119, M. 1120, M. 1121, M. 1122, M. 1123, M. 1124, M. 1125, M. 1126, M. 1127, M. 1128, M. 1129, M. 1130, M. 1131, M. 1132, M. 1133, M. 1134, M. 1135, M. 1136, M. 1137, M. 1138, M. 1139, M. 1140, M. 1141, M. 1142, M. 1143, M. 1144, M. 1145, M. 1146, M. 1147, M. 1148, M. 1149, M. 1150, M. 1151, M. 1152, M. 1153, M. 1154, M. 1155, M. 1156, M. 1157, M. 1158, M. 1159, M. 1160, M. 1161, M. 1162, M. 1163, M. 1164, M. 1165, M. 1166, M. 1167, M. 1168, M. 1169, M. 1170, M. 1171, M. 1172, M. 1173, M. 1174, M. 1175, M. 1176, M. 1177, M. 1178, M. 1179, M. 1180, M. 1181, M. 1182, M. 1183, M. 1184, M. 1185, M. 1186, M. 1187, M. 1188, M. 1189, M. 1190, M. 1191, M. 1192, M. 1193, M. 1194, M. 1195, M. 1196, M. 1197, M. 1198, M. 1199, M. 1200, M. 1201, M. 1202, M. 1203, M. 1204, M. 1205, M. 1206, M. 1207, M. 1208, M. 1209, M. 1210, M. 1211, M. 1212, M. 1213, M. 1214, M. 1215, M. 1216, M. 1217, M. 1218, M. 1219, M. 1220, M. 1221, M. 1222, M. 1223, M. 1224, M. 1225, M. 1226, M. 1227, M. 1228, M. 1229, M. 1230, M. 1231, M. 1232, M. 1233, M. 1234, M. 1235, M. 1236, M. 1237, M. 1238, M. 1239, M. 1240, M. 1241, M. 1242, M. 1243, M. 1244, M. 1245, M. 1246, M. 1247, M. 1248, M. 1249, M. 1250, M. 1251, M. 1252, M. 1253, M. 1254, M. 1255, M. 1256, M. 1257, M. 1258, M. 1259, M. 1260, M. 1261, M. 1262, M. 1263, M. 1264, M. 1265, M. 1266, M. 1267, M. 1268, M. 1269, M. 1270, M. 1271, M. 1272, M. 1273, M. 1274, M. 1275, M. 1276, M. 1277, M. 1278, M. 1279, M. 1280, M. 1281, M. 1282, M. 1283, M. 1284, M. 1285, M. 1286, M. 1287, M. 1288, M. 1289, M. 1290, M. 1291, M. 1292, M. 1293, M. 1294, M. 1295, M. 1296, M. 1297, M. 1298, M. 1299, M. 1300, M. 1301, M. 1302, M. 1303, M. 1304, M. 1305, M. 1306, M. 1307, M. 1308, M. 1309, M. 1310, M. 1311, M. 1312, M. 1313, M. 1314, M. 1315, M. 1316, M. 1317, M. 1318, M. 1319, M. 1320, M. 1321, M. 1322, M. 1323, M. 1324, M. 1325, M. 1326, M. 1327, M. 1328, M. 1329, M. 1330, M. 1331, M. 1332, M. 1333, M. 1334, M. 1335, M. 1336, M. 1337, M. 1338, M. 1339, M. 1340, M. 1341, M. 1342, M. 1343, M. 1344, M. 1345, M. 1346, M. 1347, M. 1348, M. 1349, M. 1350, M. 1351, M. 1352, M. 1353, M. 1354, M. 1355, M. 1356, M. 1357, M. 1358, M. 1359, M. 1360, M. 1361, M. 1362, M. 1363, M. 1364, M. 1365, M. 1366, M. 1367, M. 1368, M. 1369, M. 1370, M. 1371, M. 1372, M. 1373, M. 1374, M. 1375, M. 1376, M. 1377, M. 1378, M. 1379, M. 1380, M. 1381, M. 1382, M. 1383, M. 1384, M. 1385, M. 1386, M. 1387, M. 1388, M. 1389, M. 1390, M. 1391, M. 1392, M. 1393, M. 1394, M. 1395, M. 1396, M. 1397, M. 1398, M. 1399, M. 1400, M. 1401, M. 1402, M. 1403, M. 1404, M. 1405, M. 1406, M. 1407, M. 1408, M. 1409, M. 1410, M. 1411, M. 1412, M. 1413, M. 1414, M. 1415, M. 1416, M. 1417, M. 1418, M. 1419, M. 1420, M. 1421, M. 1422, M. 1423, M. 1424, M. 1425, M. 1426, M. 1427, M. 1428, M. 1429, M. 1430, M. 1431, M. 1432, M. 1433, M. 1434, M. 1435, M. 1436, M. 1437, M. 1438, M. 1439, M. 1440, M. 1441, M. 1442, M. 1443, M. 1444, M. 1445, M. 1446, M. 1447, M. 1448, M. 1449, M. 1450, M. 1451, M. 1452, M. 1453, M. 1454, M. 1455, M. 1456, M. 1457, M. 1458, M. 1459, M. 1460, M. 1461, M. 1462, M. 1463, M. 1464, M. 1465, M. 1466, M. 1467, M. 1468, M. 1469, M. 1470, M. 1471, M. 1472, M. 1473, M. 1474, M. 1475, M. 1476, M. 1477, M. 1478, M. 1479, M. 1480, M. 1481, M. 1482, M. 1483, M. 1484, M. 1485, M. 1486, M. 1487, M. 1488, M. 1489, M. 1490, M. 1491, M. 1492, M. 1493, M. 1494, M. 1495, M. 1496, M. 1497, M. 1498, M. 1499, M. 1500, M. 1501, M. 1502, M. 1503, M. 1504, M. 1505, M. 1506, M. 1507, M. 1508, M. 1509, M. 1510, M. 1511, M. 1512, M. 1513, M. 1514, M. 1515, M. 1516, M. 1517, M. 1518, M. 1519, M. 1520, M. 1521, M. 1522, M. 1523, M. 1524, M. 1525, M. 1526, M. 1527, M. 1528, M. 1529, M. 1530, M. 1531, M. 1532, M. 1533, M. 1534, M. 1535, M. 1536, M. 1537, M. 1538, M. 1539, M. 1540, M. 1541, M. 1542, M. 1543, M. 1544, M. 1545, M. 1546, M. 1547, M. 1548, M. 1549, M. 1550, M. 1551, M. 1552, M. 1553, M. 1554, M. 1555, M. 1556, M. 1557, M. 1558, M. 1559, M. 1560, M. 1561, M. 1562, M. 1563, M. 1564, M. 1565, M. 1566, M. 1567, M. 1568, M. 1569, M. 1570, M. 1571, M. 1572, M. 1573, M. 1574, M. 1575, M. 1576, M. 1577, M. 1578, M. 1579, M. 1580, M. 1581, M. 1582, M. 1583, M. 1584, M. 1585, M. 1586, M. 1587, M. 1588, M. 1589, M. 1590, M. 1591, M. 1592, M. 1593, M. 1594, M. 1595, M. 1596, M. 1597, M. 1598, M. 1599, M. 1600, M. 1601, M. 1602, M. 1603, M. 1604, M. 1605, M. 1606, M. 1607, M. 1608, M. 1609, M. 1610, M. 1611, M. 1612, M. 1613, M. 1614, M. 1615, M. 1616, M. 1617, M. 1618, M. 1619, M. 1620, M. 1621, M. 1622, M. 1623, M. 1624, M. 1625, M. 1626, M. 1627, M. 1628, M. 1629, M. 1630, M. 1631, M. 1632, M. 1633, M. 1634, M. 1635, M. 1636, M. 1637, M. 1638, M. 1639, M. 1640, M. 1641, M. 1642, M. 1643, M. 1644, M. 1645, M. 1646, M. 1647, M. 1648, M. 1649, M. 1650, M. 1651, M. 1652, M. 1653, M. 1654, M. 1655, M. 1656, M. 1657, M. 1658, M. 1659, M. 1660, M. 1661, M. 1662, M. 1663, M. 1664, M. 1665, M. 1666, M. 1667, M. 1668, M. 1669, M. 1670, M. 1671, M. 1672, M. 1673, M. 1674, M. 1675, M. 1676, M. 1677, M. 1678, M. 1679, M. 1680, M. 1681, M. 1682, M. 1683, M. 1684, M. 1685, M. 1686, M. 1687, M. 1688, M. 1689, M. 1690, M. 1691, M. 1692, M. 1693, M. 1694, M. 1695, M. 1696, M. 1697, M. 1698, M. 1699, M. 1700, M. 1701, M. 1702, M. 1703, M. 1704, M. 1705, M. 1706, M. 1707, M. 1708, M. 1709, M. 1710, M. 1711, M. 1712, M. 1713, M. 1714, M. 1715, M. 1716, M. 1717, M. 1718, M. 1719, M. 1720, M. 1721, M. 1722, M. 1723, M. 1724, M. 1725, M. 1726, M. 1727, M. 1728, M. 1729, M. 1730, M. 1731, M. 1732, M. 1733, M. 1734, M. 1735, M. 1736, M. 1737, M. 1738, M. 1739, M. 1740, M. 1741, M. 1742, M. 1743, M. 1744, M. 1745, M. 1746, M. 1747, M. 1748, M. 1749, M. 1750, M. 1751, M. 1752, M. 1753, M. 1754, M. 1755, M. 1756, M. 1757, M. 1758, M. 1759, M. 1760, M. 1761, M. 1762, M. 1763, M. 1764, M. 1765, M. 1766, M. 1767, M. 1768, M. 1769, M. 1770, M. 1771, M. 1772, M. 1773, M. 1774, M. 1775, M. 1776, M. 1777, M. 1778, M. 1779, M. 1780, M. 1781, M. 1782, M. 1783, M. 1784, M. 1785, M. 1786, M. 1787, M. 1788, M. 1789, M. 1790, M. 1791, M. 1792, M. 1793, M. 1794, M. 1795, M. 1796, M. 1797, M. 1798, M. 1799, M. 1800, M. 1801, M. 1802, M. 1803, M. 1804, M. 1805, M. 1806, M. 1807, M. 1808, M. 1809, M. 1810, M. 1811, M. 1812, M. 1813, M. 1814, M. 1815, M. 1816, M. 1817, M. 1818, M. 1819, M. 1820, M. 1821, M. 1822, M. 1823, M. 1824, M. 1825, M. 1826, M. 1827, M. 1828, M. 1829, M. 1830, M. 1831, M. 1832, M. 1833, M. 1834, M. 1835, M. 1836, M. 1837, M. 1838, M. 1839, M. 1840, M. 1841, M. 1842, M. 1843, M. 1844, M. 1845, M. 1846, M. 1847, M. 1848, M. 1849, M. 1850, M. 1851, M. 1852, M. 1853, M. 1854, M. 1855, M. 1856, M. 1857, M. 1858, M. 1859, M. 1860, M. 1861, M. 1862, M. 1863, M. 1864, M. 1865, M. 1866, M. 1867, M. 1868, M. 1869, M. 1870, M. 1871, M. 1872, M. 1873, M. 1874, M. 1875, M. 1876, M. 1877, M. 1878, M. 1879, M. 1880, M. 1881, M. 1882, M. 1883, M. 1884, M. 1885, M. 1886, M. 1887, M. 1888, M. 1889, M. 1890, M. 1891, M. 1892, M. 1893, M. 1894, M. 1895, M. 1896, M. 1897, M. 1898, M. 1899, M. 1900, M. 1901, M. 1902, M. 1903, M. 1904, M. 1905, M. 1906, M. 1907, M. 1908, M. 1909, M. 1910, M. 1911, M. 1912, M. 1913, M. 1914, M. 1915, M. 1916, M. 1917, M. 1918, M. 1919, M. 1920, M. 1921, M. 1922, M. 1923, M. 1924, M. 1925, M. 1926, M. 1927, M. 1928, M. 1929, M. 1930, M. 1931, M. 1932, M. 1933, M. 1934, M. 1935, M. 1936, M. 1937, M. 1938, M. 1939, M. 1940, M. 1941, M. 1942, M. 1943, M. 1944, M. 1945, M. 1946, M. 1947, M. 1948, M. 1949, M. 1950, M. 1951, M. 1952, M. 1953, M. 1954, M. 1955, M. 1956, M. 1957, M. 1958, M. 1959, M. 1960, M. 1961, M. 1962, M. 1963, M. 1964, M. 1965, M. 1966, M. 1967, M. 1968, M. 1969, M. 1970, M. 1971, M. 1972, M. 1973, M. 1974, M. 1975, M. 1976, M. 1977, M. 1978, M. 1979, M. 1980, M. 1981, M. 1982, M. 1983, M. 1984, M. 1985, M. 1986, M. 1987, M. 1988, M. 1989, M. 1990, M. 1991, M. 1992, M. 1993, M. 1994, M. 1995, M. 1996, M. 1997, M. 1998, M. 1999, M. 2000, M. 2001, M. 2002, M. 2003, M. 2004, M. 2005, M. 2006, M. 2007, M. 2008, M. 2009, M. 2010, M. 2011, M. 2012, M. 2013, M. 2014, M. 2015, M. 2016, M. 2017, M. 2018, M. 2019, M. 2020, M. 2021, M. 2022, M. 2023, M. 2024, M. 2025, M. 2026, M. 2027, M. 2028, M. 2029, M. 2030, M. 2031, M. 2032, M. 2033, M. 2034, M. 2035, M. 2036, M. 2037, M. 2038, M. 2039, M. 2040, M. 2041, M. 2042, M. 2043, M. 2044, M. 2045, M. 2046, M. 2047, M. 2048, M. 2049, M. 2050, M. 2051, M. 2052, M. 2053, M. 2054, M. 2055, M. 2056, M. 2057, M. 2058, M. 2059, M. 2060, M. 2061, M. 2062, M. 2063, M. 2064, M. 2065, M. 2066, M. 2067, M. 2068, M. 2069, M. 2070, M. 2071, M. 2072, M. 2073, M. 2074, M. 2075, M. 2076, M. 2077, M. 2078, M. 2079, M. 2080, M. 2081, M. 2082, M. 2083, M. 2084, M. 2085, M. 2086, M. 2087, M. 2088, M. 2089, M. 2090, M. 2091, M. 2092, M. 2093, M. 2094, M. 2095, M. 2096, M. 2097, M. 2098, M. 2099, M. 2100, M. 2101, M. 2102, M. 2103, M. 2104, M. 2105, M. 2106, M. 2107, M. 2108, M. 2109, M. 2110, M. 2111, M. 2112, M. 2113, M. 2114, M. 2115, M. 2116, M. 2117, M. 2118, M. 2119, M. 2120, M. 2121, M. 2122, M. 2123, M. 2124, M. 2125, M. 2126, M. 2127, M. 2128, M. 2129, M. 2130, M. 2131, M. 2132, M. 2133, M. 2134, M. 2135, M. 2136, M. 2137, M. 2138, M. 2139, M. 2140, M. 2141, M. 2142, M. 2143, M. 2144, M. 2145, M. 2146, M. 2147, M. 2148, M. 2149, M. 2150, M. 2151, M. 2152, M. 2153, M. 2154, M. 2155, M. 2156, M. 2157, M. 2158, M. 2159, M. 2160, M. 2161, M. 2162, M. 2163, M. 2164, M. 2165, M. 2166, M. 2167, M. 2168, M. 2169, M. 2170, M. 2171, M. 2172, M. 2173, M. 2174, M. 2175, M. 2176, M. 2177, M. 2178, M. 2179, M. 2180, M. 2181, M. 2182, M. 2183, M. 2184, M. 2185, M. 2186, M. 2187, M. 2188, M. 2189, M. 2190, M. 2191, M. 2192, M. 2193, M. 2194, M. 2195, M. 2196, M. 2197, M. 2198, M. 2199, M. 2200, M. 2201, M. 2202, M. 2203, M. 2204, M. 2205, M. 2206, M. 2207, M. 2208, M. 2209, M. 2210, M. 2211, M. 2212, M. 2213, M. 2214, M. 2215, M. 2216, M. 2217, M. 2218, M. 2219, M. 2220, M. 2221, M. 2222, M. 2223, M. 2224, M. 2225, M. 2226, M. 2227, M. 2228, M. 2229, M. 2230, M. 2231, M. 2232, M. 2233, M. 2234, M. 2235, M. 2236, M. 2237, M. 2238, M. 2239, M. 2240, M. 2241, M. 2242, M. 2243, M. 2244, M. 2245, M. 2246, M. 2247, M. 2248, M. 2249, M. 2250, M. 2251, M. 2252, M. 2253, M. 2254, M. 2255, M. 2256, M. 2257, M. 2258, M. 2259, M. 2260, M. 2261, M. 2262, M. 2263, M. 2264, M. 2265, M. 2266, M. 2267, M. 2268, M. 2269, M. 2270, M. 2271, M. 2272, M. 2273, M. 2274, M. 2275, M. 2276, M. 2277, M. 2278, M. 2279, M. 2280, M. 2281, M. 2282, M. 2283, M. 2284, M. 2285, M. 2286, M. 2287, M. 2288, M. 2289, M. 2290, M. 2291, M. 2292, M. 2293, M. 2294, M. 2295, M. 2296, M. 2297, M. 2298, M. 2299, M. 2300, M. 2301, M. 2302, M. 2303, M. 2304, M. 2305, M. 2306, M. 2307, M. 2308, M. 2309, M. 2310, M. 2311, M. 2312, M. 2313, M. 2314, M. 2315, M. 2316, M. 2317, M. 2318, M. 2319, M. 2320, M. 2321, M. 2322, M. 2323, M. 2324, M. 2325, M. 2326, M. 2327, M. 2328, M. 2329, M. 2330, M. 2331, M. 2332, M. 2333, M. 2334, M. 2335, M. 2336, M. 2337, M. 2338, M. 2339, M. 2340, M. 2341, M. 2342, M. 2343, M. 2344, M. 2345, M. 2346, M. 2347, M. 2348, M. 2349, M. 2350, M. 2351, M. 2352, M. 2353, M. 2354, M. 2355, M. 2356, M. 2357, M. 2358, M. 2359, M. 2360, M. 2361, M. 2362, M. 2363, M. 2364, M. 2365, M. 2366, M. 2367, M. 2368, M. 2369, M. 2370, M. 2371, M. 2372, M. 2373, M. 2374, M. 2375, M. 2376, M. 2377, M. 2378, M. 2379, M. 2380, M. 2381, M. 2382, M. 2383, M. 2384, M. 2385, M. 2386, M. 2387, M. 2388, M. 2389, M. 2390, M. 2391, M. 2392, M. 2393, M. 2394, M. 2395, M. 2396, M. 2397, M. 2398, M. 2399, M. 2400, M. 2401, M. 2402, M. 2403, M. 2404, M. 2405, M. 2406, M. 2407, M. 2408, M. 2409, M. 2410, M. 2411, M. 2412, M. 2413, M. 2414, M. 2415, M. 2416, M. 2417, M. 2418, M. 2419, M. 2420, M. 2421, M. 2422, M. 2423, M. 2424, M. 2425, M. 2426, M. 2427, M. 2428, M. 2429, M. 2430, M. 2431, M. 2432, M. 2433, M. 2434, M.

Where weed growth is present at application time, SENCOR should be applied with Gramoxone or other contact herbicide. Refer to the other product label registered for additional directions, rates, and weed species controlled.

Weeds Controlled:

Broadleaves

Chickweed, Common
(*Stellaria media*)
Herb
(*Lamium amplexicaule*)
K. Weed
(*Chenopodium album*)
Mustard, Blue or Purple
(*Chorispora tenella*)
Mustard, Jim Hill
(*Sisymbrium albidum*)
Mustard, Tansy
(*Oenothera biennis*)

Mustard, Weeds
(*Erysimum repandum*)
Mustard, Wild
(*Barbarea vulgaris*)
Purslane, Field, Fat
(*Portulaca oleraceae*)
Ragwort
(*Senecio jacobinae*)
Russet thistle
(*Helianthus annuus*)
Wild radish
(*Raphanus sativus*)

Grasses

Heatgrass
(*Echinochloa crusgalli*)
L. Grass
(*Lolium temulentum*)

Wheat, Green
(*Triticum aestivum*)

NOTE

After Harvest Application (Fall Fallow) SENCOR may be applied to wheat stubble after harvest in the Fall. Apply 1 to 1.5 lb. SENCOR per acre broadcast before weeds emerge. Use higher rate for improved weed control, or for weeds designated as requiring the higher rate for control. Rastali (1/2 inch or more) is necessary for herbicide activation.

Do not plant crops in treated areas until at least 10 days after SENCOR application.

SENCOR may be applied to corn, soybeans, and other crops in the fall or spring. Apply 1 to 1.5 lb. SENCOR per acre broadcast before weeds emerge. Use higher rate for improved weed control, or for weeds designated as requiring the higher rate for control. Rastali (1/2 inch or more) is necessary for herbicide activation.

Spring Application (Summer Fallow) SENCOR may be applied to corn, soybeans, and other crops in the spring. Apply 1 to 1.5 lb. SENCOR per acre broadcast before weeds emerge. Use higher rate for improved weed control, or for weeds designated as requiring the higher rate for control. Rastali (1/2 inch or more) is necessary for herbicide activation.

BEST AVAILABLE COPY

For more information, contact your local SENCOR distributor.

PRECAUTION AND RESTRICTIONS

SENCOR is a restricted use herbicide. It is not to be used on crops or in areas where it may cause injury to crops, livestock, or wildlife. It is not to be used on crops or in areas where it may cause injury to crops, livestock, or wildlife.

FOR WEED CONTROL IN A FALLOW ROTATION WITH BARLEY AND WHEAT

Colorado, Kansas, Montana, Nebraska, and Wyoming only.

SENCOR is a restricted use herbicide. It is not to be used on crops or in areas where it may cause injury to crops, livestock, or wildlife. It is not to be used on crops or in areas where it may cause injury to crops, livestock, or wildlife.

Weeds Controlled

Broadleaves

Chickweed, Common
(*Stellaria media*)
Herb
(*Lamium amplexicaule*)
K. Weed
(*Chenopodium album*)
Mustard, Blue or Purple
(*Chorispora tenella*)
Mustard, Jim Hill
(*Sisymbrium albidum*)
Mustard, Tansy
(*Oenothera biennis*)

Mustard, Weeds
(*Erysimum repandum*)
Mustard, Wild
(*Barbarea vulgaris*)
Purslane, Field, Fat
(*Portulaca oleraceae*)
Ragwort
(*Senecio jacobinae*)
Russet thistle
(*Helianthus annuus*)
Wild radish
(*Raphanus sativus*)

431-46

Grasses		
Cheatgrass (<i>Bromus secalinus</i>)	*Foxtail, Green (<i>Setaria viridis</i>)	*Wild Oats (<i>Avena fatua</i>)
Downy Brome (<i>Bromus tectorum</i>)	*Wheat, Volunteer (<i>Triticum</i> spp.)	

NOTE: Since control of these weeds may vary, use the higher rate for control of weeds designated as requiring the higher rate for control. Rainfall is not a factor in determining the herbicide activation.

After Harvest Application: (Fall Fallow) SENCOR 4 may be applied to the stubble after harvest in the Fall. Apply 1 1/4 to 1 1/2 pints per acre broadcast before weeds emerge. Use the higher rate for longer weed control or for weeds designated as requiring the higher rate for control. Rainfall is not a factor in determining the herbicide activation.

Spring Application (Summer Fallow) SENCOR 4 may be applied to the stubble in the Spring. Apply 1 1/4 to 1 1/2 pints per acre broadcast before weeds emerge in the Spring. Use the higher rate for longer weed control or for weeds designated as requiring the higher rate for control. Rainfall is not a factor in determining the herbicide activation. Wheat or barley can be planted the day after Spring application.

PRECAUTIONS AND RESTRICTIONS

Do not plant Spring seeded cereals in the fallow area for fallow.

Where SENCOR 4 was applied in the Fall, do not apply SENCOR in the Spring.

CROP ROTATION RECOMMENDATIONS:

Waiting Period After SENCOR Herbicide Application			
4 Months	8 Months	12 Months	18 Months
Barley	Barley	Barley	Barley
Wheat	Wheat	Wheat	Wheat
Oats	Oats	Oats	Oats
Truck	Truck	Truck	Truck
Hay	Hay	Hay	Hay
Alfalfa	Alfalfa	Alfalfa	Alfalfa
Legume	Legume	Legume	Legume
Grass	Grass	Grass	Grass
Other	Other	Other	Other

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS. HAZARDS TO HUMANS AND DOMESTIC ANIMALS. CAUTION.

May be harmful if swallowed. Avoid contact with skin and eyes. Wash thoroughly with soap and water after handling. Do not eat, drink, or smoke while handling. Do not get on clothing or in shoes. Do not use near water. Do not use near food or feed. Do not use near children or pets.

To Physician: In case of poisoning, give information on label. **Symptoms of Poisoning:** Nausea, vomiting, diarrhea, dizziness, headache, weakness, loss of consciousness.

STATEMENTS OF PRACTICAL TREATMENT.

If in eyes: Flush immediately with water. **If swallowed:** Vomiting should be induced. **If on skin:** Wash thoroughly with soap and water. Do not use near water. Do not use near food or feed. Do not use near children or pets.

ENVIRONMENTAL HAZARDS.

May be harmful to fish and aquatic life. Do not use near water. Do not use near food or feed. Do not use near children or pets. Do not use near crops or livestock.

GROUND WATER ADVISORY: Metolachlor is a herbicide which may be found in ground water. Do not use near water. Do not use near food or feed. Do not use near children or pets.

should not be applied methidathion where the water table (ground water) is close to the surface and where there are any vegetables. Use on well-drained soils such as loamy sands. For more information, agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

RE-ENTRY STATEMENTS.

Do not apply this product to such a market as to the fruit or vegetable, or to ornamentals or other perennials. The area being treated must be vacated by unprotected persons. Do not re-enter treated areas within the re-entry time period. Do not re-enter treated areas. Because certain states may require more restrictive re-entry times for various crops treated with this product, consult your state department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings should include those statements found under the label heading "Precautionary Statements" and the Re-entry Statements included on this label. Written or oral warnings must be given to all workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given to all workers who are expected to be in a treated area or in an area about to be treated with this product. Written warnings must include the following: "Do not enter treated area until sprays have dried. If on skin, remove clothing immediately and wash skin immediately with soap and water. If eyes are contaminated, wash with flowing water for at least 15 minutes. If swallowed, vomiting should be induced (see other precautionary statements on label)."

STORAGE AND DISPOSAL.

Pesticide Disposal

Do not contaminate food, feed, or other products. Do not dispose of this product in a way that will cause harm to humans or the environment.

Container Disposal

Do not reuse this container for any other purpose. Do not dispose of this container in a way that will cause harm to humans or the environment. Do not burn this container. Do not use this container for any other purpose.

BEST AVAILABLE COPY