

314-7036.YLD

Reg # 3125 - 314

PM-25

Base Pre-Reg. (7036)

182

U.S. LABEL

Date of Draft: 5/25/90 (Pre-Reg)(H)

Reason to Issue: To add suppression of downy brome in Winter wheat.

EPA Reg. No. 3125-314

SENCOR® 4
Flowable Herbicide

ACTIVE INGREDIENT:

4-Amino-6-(1,1-dimethylethyl)-3-(methylthio)-1,2,4-triazin-5(4H)-one 41%

INERT INGREDIENTS: 59%
100%

AMENDMENT TO CURRENTLY REGISTERED LABELING

REVISE TO READ AS FOLLOWS:

282

SENCOR 4 Flowable herbicide


WINTER WHEAT

FOR USE IN IDAHO, MONTANA, NEVADA, OREGON, UTAH, AND WASHINGTON ON THE VARIETIES CENTURK, CHEYENNE, DAWS, FARO, GAINES, HATTON, HILL 81, HYSLOP, LUKE, McDERMID, MORO, NUGAINES, PAHA, PECK, SPRAGUE, STEPHENS, TYEE, WANSER, WESTON, AND WINALTA

SENCOR 4 flowable herbicide, in a tank mix combination with one of the herbicide products specified below, may be applied to wheat after the crop has reached the 2-leaf stage for control of the following broadleaf weeds:

- Common chickweed (Stellaria media)
- Henbit (Lamium amplexicaule)
- Prickly lettuce (Lactuca serriola)
- Blue mustard (Chorispora tenella)
- Field pennycress (Thlaspi arvense)
- Tarweed fiddleneck (Amsinckia lycopsoides)

Suppression only: Downy brome (Bromus tectorum)



RECOMMENDED APPLICATIONS

| CROP | SENCOR 4 Fluid Ounces Per Acre | REMARKS |
|-----------------|--------------------------------------|--|
| Winter Wheat | 3 to 4 1/2 | Use only in a tank mix combination for broadleaf weed control.* The specified dosage per acre may be added to other herbicides registered for use on wheat after the crop has reached the 2-leaf stage, but not during winter dormancy. See the appropriate sections of this label and the label for other herbicide products that are tank mixed with SENCOR 4 for directions, application rates, mixing procedures, special precautions and restrictions before using. |

*May be used with: Bronate, Bucril, Banvel, 2,4-D, MCPA, Glean, Ally, Finesse, Harmony, or Harmony Extra.

Observe all cautions and limitations on labeling of all products used in mixtures.

SENCOR 4 Flowable herbicide


WINTER WHEAT

FOR USE IN IDAHO, MONTANA, NEVADA, OREGON, UTAH, AND WASHINGTON ON THE VARIETIES CENTURK, CHEYENNE, DAWS, FARO, GAINES, HATTON, HILL 81, HYSLOP, LUKE, McDERMID, MORO, NUGAINES, PAHA, PECK, SPRAGUE, STEPHENS, TYEE, WANSER, WESTON, AND WINALTA

SENCOR 4 flowable herbicide, in a tank mix combination with one of the herbicide products specified below, may be applied to wheat after the crop has reached the 2-leaf stage for control of the following broadleaf weeds:

| | |
|--------------------|----------------------------------|
| Common chickweed | (<u>Stellaria media</u>) |
| Henbit | (<u>Lamium amplexicaule</u>) |
| Prickly lettuce | (<u>Lactuca serriola</u>) |
| Blue mustard | (<u>Chorispora tenella</u>) |
| Field pennycress | (<u>Thlaspi arvense</u>) |
| Tarweed fiddleneck | (<u>Amsinckia lycopsoides</u>) |

Suppression only: Downy brome (Bromus tectorum)



RECOMMENDED APPLICATIONS

| CROP | SENCOR 4 Fluid Ounces Per Acre | REMARKS |
|-----------------|--------------------------------------|--|
| Winter Wheat | 3 to 4 1/2 | Use only in a tank mix combination for broadleaf weed control.* The specified dosage per acre may be added to other herbicides registered for use on wheat after the crop has reached the 2-leaf stage, but not during winter dormancy. See the appropriate sections of this label and the label for other herbicide products that are tank mixed with SENCOR 4 for directions, application rates, mixing procedures, special precautions and restrictions before using. |

*May be used with: Bronate, Bucril, Banvel, 2,4-D, MCPA, Glean, Ally, Finesse, Harmony, or Harmony Extra.

Observe all cautions and limitations on labeling of all products used in mixtures.