


U.S. LABEL

Reason to Issue: To add claim for tank  
mix with Amber<sup>®</sup> herbicide for use on  
barley and wheat.

Date of Draft: 8/10/92 (Pre-Reg)(H)

EPA Reg. No. 3125-314



SENCOR 4  
Flowable Herbicide

ACTIVE INGREDIENT:

4-Amino-6-(1,1-dimethylethyl)-3-  
(methylthio)-1,2,4-triazin-5(4H)-one .... 41%

INERT INGREDIENTS: ..... 59%  
100%

AMENDMENT TO CURRENTLY REGISTERED LABELINGREVISE TO READ AS FOLLOWS:

## SENCOR 4 Flowable Herbicide

---

### CEREALS

(Spring and Winter Barley and Winter Wheat)

For suppression or control of certain grass and broadleaf weeds, apply SENCOR 4 flowable herbicide postemergence to barley (Winter and Spring) and Winter wheat (see the appropriate section of this label for geographic limitations):

1. Barley (Spring and Winter) and Winter Wheat in Idaho, Montana, Nevada, Oregon, Utah, and Washington.
2. Dryland (Non-irrigated) Winter wheat in Kansas, Oklahoma, and Texas.
3. Dryland (Non-irrigated) Winter wheat only (Arkansas, Georgia, Kentucky, Louisiana, Mississippi, Tennessee, and Missouri "Bootheel").

### Mixing

See the "General Information" section in the front of this label for specific mixing procedures. Do not tank mix SENCOR 4 with pesticides other than those specified below. When tank-mixing, carefully follow the instructions on this label. Refer to the other product label registered for use on barley and wheat for additional directions, rates, weeds controlled and restrictions.

### Application

Refer to the "General Information" section in the front of this label for detailed information on the application of SENCOR 4. Do not exceed the rates specified below.

Apply SENCOR 4 when crop is healthy and actively growing. Weed control may not be observed for 2 to 4 weeks after application; under very dry conditions, weed control may not be apparent for 4 to 6 weeks. Moisture (at least 1/2 inch) is needed within 2 to 3 weeks after application to move SENCOR into the weed root zone. Lack of adequate moisture after application may result in poor control. Control or suppression of listed weeds is dependent on weed size. Control or suppression can be reduced if broadleaf weeds are taller than 1 inch or grasses have more than 2 leaves.

If SENCOR 4 is applied more than once per crop season, allow at least 45 days between applications.

SENCOR 4 may be tank mixed with Ally, Amber, Finesse, Glean, Harmony, Harmony Extra, 2,4-D, MCPA, Igran, Banvel, Bronate, or Buctril herbicides. Refer to the other product labels registered for use on cereals for additional directions, rates, and weed species controlled. Observe all cautions and limitations on labeling of all products used in mixtures.