

YLD28

PM21
3125-318
pg 1 of 1

U.S. LABEL

Reason to Issue: To modify references
to control of summer patch and
fusarium blight on turf.

Date of Draft: 11/1/85 (Pre-Reg) (S)

EPA Reg. No. 3125-318

® BAYLETON

ACCEPTED

APR 7 1986

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 3125-318

25% Wettable Powder

SYSTEMIC FUNGICIDE

ACTIVE INGREDIENT:

1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H
-1,2,4-triazol-1-yl)-2-butanone 25%

Revise references to summer patch and fusarium blight as follows:

RECOMMENDED APPLICATIONS
TURF GRASS DISEASE CONTROL

DISEASE	Ozs. of BAYLETON 25% W.P. Per 1,000 Sq. Ft.* <u>Preventive Rates Only</u>	REMARKS
Fusarium Blight (<u>Fusarium culmorum</u>) (<u>Fusarium poae</u>) Summer Patch (<u>Phialophora graminicola</u>) (Except California)	2	Apply first application in mid-June or 30 days prior to time this blight normally becomes evident. Applications should be made on 30 day intervals.
Fusarium Blight (<u>Fusarium culmorum</u>) (<u>Fusarium poae</u>) Summer Patch (<u>Phialophora graminicola</u>) (California only)	2 to 4	Apply first application in mid-June or 30 days prior to time this blight normally becomes evident. Applications should be made at the 2 oz. rate on 15 day intervals or at the 4 oz. rate on 30 day intervals.

Note: *Apply the specified amount of BAYLETON 25% W.P. in 2 to 4 gallons of spray per 1,000 sq. ft. For equivalent rates per acre see Conversion Table below. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. Do not use clippings for animal feed.


YLD28

U.S. LABEL

Reason to Issue: To modify references
to control of summer patch and
fusarium blight on turf.

Date of Draft: 11/1/85 (Pre-Reg) (S)

EPA Reg. No. 3125-318



® BAYLETON

25% Wettable Powder

SYSTEMIC FUNGICIDE

ACTIVE INGREDIENT:

1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H
-1,2,4-triazol-1-yl)-2-butanone 25%

Revise references to summer patch and fusarium blight as follows:

RECOMMENDED APPLICATIONS
TURF GRASS DISEASE CONTROL

DISEASE	Ozs. of BAYLETON 25% W.P. Per 1,000 Sq. Ft.*	REMARKS
	<u>Preventive Rates Only</u>	
Fusarium Blight (<u>Fusarium culmorum</u>) (<u>Fusarium poae</u>) Summer Patch (<u>Phialophora graminicola</u>) (Except California)	2	Apply first application in mid-June or 30 days prior to time this blight normally becomes evident. Applications should be made on 30 day intervals.
Fusarium Blight (<u>Fusarium culmorum</u>) (<u>Fusarium poae</u>) Summer Patch (<u>Phialophora graminicola</u>) (California only)	2 to 4	Apply first application in mid-June or 30 days prior to time this blight normally becomes evident. Applications should be made at the 2 oz. rate on 15 day intervals or at the 4 oz. rate on 30 day intervals.

Note: *Apply the specified amount of BAYLETON 25% W.P. in 2 to 4 gallons of spray per 1,000 sq. ft. For equivalent rates per acre see Conversion Table below. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. Do not use clippings for animal feed.