

SEP 25 1992

John S. Thornton  
Miles Inc.  
P.O. Box 4913  
Kansas City, MO 64120

Dear Mr. Thornton:

Subject: Label Revisions  
Bayleton 25% Wettable Powder  
EPA Registration No. 3125-318  
Your Submission Dated September 14, 1992

The amendment referred to above, submitted in connection with registration under section 3(c)(7)(A) of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable provided that you:

1. Submit/cite all data required for registration/-reregistration of your product under FIFRA section 3(c)(5) or 4(a) when the Agency requires all registrants of similar products to submit such data.

2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

- In the environmental hazards section delete:

Do not use on crops grown for food or forage.


3. Submit one (1) copy of your final printed labeling before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

2013

-2-

A stamped copy of the label is enclosed for your records.

Sincerely yours,



Cynthia Giles-Parker  
Product Manager (22)  
Fungicide-Herbicide Branch  
Registration Division (H7505C)

Enclosure

PM 22 3125-318

20/13

**U.S. LABEL (S)**

**Reason to Issue:** To propose use on Bermuda grass decline; propose use on cedar trees for Didymascella thujina control; to add rate for Melampsora spp. control; change reference marks from asterisks to letters; to revise corporate designation; delete ® and trademark acknowledgement statement; 9/3/92 Draft; to revise in accordance with EPA comments (Environmental Hazards, Directions for Use, & Remarks).

Date of Draft: 9/3/92 Pre-Reg (S)  
Supersedes Pre-Reg Draft Dated: 6/25/92

ACCEPTED  
with COMMENTS  
In EPA Letter Dated:

SEP 25 1992

Under the Federal Insecticide, Fungicide, and Rodenticide Act  
Registration for the pesticide  
Registration Number EPA Reg. No.

3125-318

EPA Reg. No. 3125-318

**BAYLETON**

**25% WETTABLE POWDER**

**SYSTEMIC FUNGICIDE**

FOR CONTROL OF CERTAIN DISEASES ON FLOWERS, FOLIAGE PLANTS, SHRUBS,  
TURF AND SHADE TREES

**ACTIVE INGREDIENT:**

1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanone .....	25%
INERT INGREDIENTS .....	75%
	<u>100%</u>

U.S. Patent No. 3,912,752

EPA Reg. No. 3125-318

EPA Est. 3125-MO-1

**STOP - READ THE LABEL BEFORE USE**

KEEP OUT OF THE REACH OF CHILDREN

**CAUTION**

See Rear Panel for Statements of Practical Treatment and Other Precautionary Statements

NET WEIGHT \_\_\_\_\_ POUNDS

Miles Inc.  
Specialty Products  
Box 4913, Kansas City, MO 64120-0013

Note: Underlined letters in chemical nomenclature should be italicized when printed, not underlined.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Harmful if swallowed, inhaled, or absorbed through the skin. Causes eye irritation. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**STATEMENTS OF PRACTICAL TREATMENT**

If swallowed: Call a physician or Poison Control Center. Vomiting should be induced. Administer water freely and induce vomiting by giving one dose (1/2 oz. or 15 ml.) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person. Professional medical assistance should be secured immediately.

If in eyes: Flush with plenty of water. Call a physician if irritation persists.

If on skin: Wash with plenty of soap and water. Get medical attention.

**SYMPTOMS OF POISONING:** The compound does not cause any definite symptoms that would be diagnostic. Poisoning is accompanied by hyperactivity followed by sedation

**NOTE TO PHYSICIAN:** No specific antidote. Treat symptomatically.

**ENVIRONMENTAL HAZARDS**

Do not use on crops grown for food or forage. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

Do not make applications when weather conditions favor drift from target area.

**AVISO:**

**PRECAUCION AL USUARIO:** Si usted no puede leer o entender inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

**(TO THE USER:** If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**IMPORTANT:** Read these entire Directions and Conditions of Sale before using BAYLETON 25% Wettable Powder systemic fungicide.

**CONDITIONS OF SALE:** THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE APPROPRIATE FOR THE CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED UNDER DIFFERENT ENVIRONMENTAL CONDITIONS BOTH INDOORS AND OUTDOORS UNDER CONDITIONS SIMILAR TO THOSE THAT ARE ORDINARY AND CUSTOMARY WHERE THE PRODUCT IS TO BE USED. INSUFFICIENT CONTROL OF PESTS OR PLANT INJURY MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL CONDITIONS, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO ANIMALS, MAN, AND DAMAGE TO THE ENVIRONMENT. MILES OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL ENVIRONMENTAL CONDITIONS, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF MILES AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

**APPLICATION:**

BAYLETON fungicide is absorbed rapidly and works systemically from within the plant. Good coverage and wetting of the foliage is necessary. Rainfall or sprinkler irrigation, within 1/2 hour after application does not decrease effectiveness. Control may be less effective on plants suffering from drought stress. Therefore, in order to achieve maximum control, plants should be maintained in a vigorously growing state through good cultural practices.

Apply in all cases when plants are fully established and actively growing. Applications should be applied at recommended intervals to maintain disease control.

Do not use on crops grown for food or forage.

**BEST AVAILABLE COPY**

**USE IN CHEMIGATION SYSTEMS ON TURF ONLY:**

Apply BAYLETON 25% Wettable Powder only through solid set irrigation systems. Do not apply this product through any other type of irrigation system.

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back towards the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump water stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where the pesticide distribution is adversely affected.

USE IN CHEMIGATION SYSTEMS ON TURF ONLY (Continued):

Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the areas intended for treatment.

Crop injury or lack of effectiveness can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise. ✓

Pre-mix the required amount of BAYLETON 25% Wettable Powder, as determined under "Recommended Applications", in sufficient water to uniformly inject the entire mixture during the last 5 minutes of the irrigation cycle using a positive pressure pumping system. Continuous agitation of the mixture in the holding tank is required to maintain suspension of the product. The injection must occur during the last 5 minutes of the irrigation cycle.

1/01

**RECOMMENDED APPLICATIONS  
TURF GRASS DISEASE CONTROL**

DISEASE	OZS OF BAYLETON 25% WP PER 1,000 SQUARE FOOT*		REMARKS
	Preventive	Curative	
Dollar Spot ( <u>Sclerotinia homoeocarpa</u> )	1	-	<b>PREVENTIVE RATE:</b> Apply recommended rate on 30 day intervals. Protective activity of BAYLETON may extend for as long as 60 days dependent upon environmental conditions.
	0.5	-	<b>PREVENTIVE RATE:</b> (Except California): Apply recommended rate on 14 day intervals. Protective activity of BAYLETON may extend for as long as 30 days dependent upon environmental conditions.
	-	2	<b>CURATIVE RATE:</b> To control existing infections, apply the curative rate. Subsequent applications should be applied on a preventive schedule and rate.
Copper Spot ( <u>Gloeocercospora sorghi</u> ) Powdery Mildew ( <u>Epysiphe graminis</u> ) Corticium Red Thread ( <u>Laelisaria fuciformis</u> ) Rusts ( <u>Puccinia spp.</u> ) Brown Patch/ Rhizoctonia Bl. t ( <u>Rhizoctonia solani</u> ) (Suppression)	1	2	<b>PREVENTIVE RATE:</b> Apply at 15 to 30 day intervals. When environmental conditions favor light to moderate disease development, use a longer interval. Protective activity of BAYLETON can be greater than 30 days depending on environmental conditions.  <b>CURATIVE RATE:</b> To control existing infections, apply the curative rate. Subsequent applications should be applied on a preventive schedule and rate.

**BEST AVAILABLE COPY**

RECOMMENDED APPLICATIONS (Continued) TURF GRASS DISEASE CONTROL			
DISEASE	OZS OF BAYLETON 25% WP PER 1,000 SQUARE FOOT*		REMARKS
Anthracnose ( <u>Colletotrichum</u> <u>graminicola</u> )	<u>Preventive</u> 2	<u>Curative</u> -	<u>PREVENTIVE RATE:</u> Apply at 30 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 45 days.
	1	-	<u>PREVENTIVE RATE</u> (Except California): Apply at 21 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 30 days.
	-	2	<u>CURATIVE RATE:</u> To control existing infections, apply the curative rate. Subsequent applications should be applied on a preventive schedule and rate.
Southern Blight ( <u>Sclerotium rolfsii</u> )	<u>Preventive</u> 1 to 4	<u>Curative</u> 4	<u>PREVENTIVE:</u> Begin applications prior to the appearance of disease symptoms. Depending on anticipated disease severity, apply 2 to 4 oz rates at 14 day intervals for the initial 2 to 3 treatments. Apply subsequent treatments of 1 to 2 oz at 14 to 28 day intervals.  <u>CURATIVE:</u> To control existing infections, apply 4 oz at 14 day intervals for the initial 2 to 3 treatments followed by 1 to 2 oz at 14 to 28 day intervals.
Stripe Smut ( <u>Ustilago</u> <u>striiformis</u> )	<u>Preventive Rates Only</u> 2		Make a single application in spring as grass growth begins.
Fusarium Blight ( <u>Fusarium culmorum</u> ) ( <u>Fusarium poae</u> ) Summer Patch ( <u>Phaeophora</u> <u>graminicola</u> ) ( <u>Magnaporthe poae</u> )	2 to 4		Apply first application in the Spring 30 to 60 days before initial symptoms normally appear. Repeat applications at 30 day intervals as needed.

9/12/12

RECOMMENDED APPLICATIONS (Continued) TURF GRASS DISEASE CONTROL		
DISEASE	OZS OF BAYLETON 25% WP PER 1,000 SQUARE FOOT*	REMARKS
<p><u>Gaeumannomyces</u> <u>graminis</u> var. <u>graminis</u></p> <p>(Such as Bermudagrass decline, take all patch on centipede grass)</p>	<p><u>Preventive</u>    <u>Curative</u></p> <p>2                      4</p>	<p>Immediately after the fungicide is applied, the area should be thoroughly irrigated to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone. The objective is to water the fungicide into the crown and root zone.</p> <p><u>PREVENTIVE RATE:</u> Begin applications 2 to 4 weeks prior to the appearance of disease symptoms. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information. Apply subsequent applications at 21 to 28 day intervals.</p> <p><u>CURATIVE RATE:</u> To control existing infections, apply 4 oz at 14 to 21 day intervals for the initial 1 or 2 treatments followed by 2 oz at 21 to 28 day intervals. Cultural control practices such as aerification, topdressing, reseeding, and fertilization must be implemented prior to or at the same time the fungicide is applied. Refer to your local County Extension Service for this information.</p>

**BEST AVAILABLE COPY**

RECOMMENDED APPLICATIONS (Continued) TURF GRASS DISEASE CONTROL		
DISEASE	OZS OF BAYLETON 25% WP PER 1,000 SQUARE FOOT*	REMARKS
Gray Snow Mold/ Typhula Blight ( <u>Typhula incarnata</u> )  ( <u>Except California</u> )	<u>Preventive Rates Only</u>  4	Apply in fall 30 days prior to turf dormancy. If turf breaks dormancy during winter months a second application should be made. Do not apply over snow cover, or when turf is dormant.
Pink Snow Mold/ Fusarium Patch ( <u>Microdochium nivalis</u> )  ( <u>Except California</u> )	2 to 4	Apply before conditions favorable for infection occur. Re-application should be made as needed at a 60 to 90 day interval. Do not apply over snowcover, or when turf is dormant. Use higher rate in areas with a history of severe disease damage.
Gray Snow Mold/ Typhula Blight ( <u>Typhula incarnata</u> ) Pink Snow Mold/ Fusarium Patch ( <u>Microdochium nivalis</u> )  ( <u>California only</u> )	Single Application  5 to 8	Apply in late fall before first snow cover. Do not apply over snow cover. Use the high rate in areas with a history of severe snowmold damage.

Note: \*Apply the specified amount of BAYLETON 25% Wettable Powder in 2 to 4 gallons of spray per 1,000 sq. ft. For equivalent rates per acre see Conversion Table below. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. Do not use clippings for animal feed.

Conversion Table  
for  
BAYLETON 25% Wettable Powder

<u>Ounces per 1000 sq. ft.</u>	=	<u>pounds per acre</u>
1	=	2 3/4
2	=	5 1/2
4	=	11
5	=	<u>13 3/4</u>

ORNAMENTAL PLANT DISEASE CONTROL

PLANTS (Disease Controlled Reference)

DISEASES

Flowering & Foliage Plants

(Outdoor) (Except California)

- Ageratum (2b, 3, 4)
- Aster (4)
- Begonia (3)
- Canna (4)
- Carnation (3, 4)
- Chrysanthemum (3, 4)
- Dahlia (3)
- Delphinium (3)
- Dianthus (4)
- Four O'Clock (4)
- Geranium (3, 4)
- Hollyhock (3, 4)
- Iris (2c)
- Marigold (2b, 4)
- Nephtytis (2a)
- Pansy (3, 4)
- Petunia (3, 4)
- Phlox (2b, 3, 4)
- Rose (3)
- Salvia (3, 4)
- Sedum (3)
- Snapdragon (3, 4)
- Sunflowers (3, 4)  
(ornamental only)
- Sweet peas (3)
- Zinnia (2b, 3)

Shade Trees (Except California)

- Ash (3)
- Aspen (3, 4)
- Birch (3, 4)
- Buckeye (3)
- Chestnut (3)
- Cottonwood (3, 4)
- Elm (3)
- Fir (4)
- Locust (3)
- Maple (3)
- Oak (3)
- Pine (4, 5)
- Poplar (3, 4)
- Russian Olive (2b, 4)
- Sycamore (3)
- Walnut (3)
- Willow (3, 4)

Hawaii Only

- Dendrobium (1c)

California Only Recommendations

- Azalea (1a, 3)
- Begonia (3)
- Cedar (2d)
- Crape myrtle (3)
- Euonymus (3)
- Geranium (3, 4)
- Hollyhock (3, 4)
- Iris (2c)
- Nephtytis (2a)
- Oak (3)
- Pine Trees (4, 5)
- Rose (3)
- Snapdragon (3, 4)
- Sweet peas (3)
- Sycamore (3)
- Viburnum (3)
- Willow (4)
- Zinnia (2b, 3)

Ornamental Shrubs & Trees

- (Except California)
- Amelanchier (3)
- Azalea (1a, 2f, 3)
- Barberry (3, 4)
- Buckthorn (4)
- Camellia (suppression of 1b)
- Cedar (2d)
- Crabapple (flowering) (3, 4)
- Crape myrtle (3)
- Dogwood (3)
- Euonymus (3)
- Gardenia (3)
- Hawthorn (3, 4)
- Hemlock (4e)
- Holly (3)
- Juniper (4)
- Leucothoe (2b)
- Lilac (3)
- Mock-Orange (3, 4)
- Mountain Laurel (1a, 2b, 3)
- Ninebark (3)
- Paulownia (3)  
(Empress Tree)
- Pear (Flowering) (3)
- Photinia (2e, 3, 4)
- Potentilla (4)  
(Cinquefoil)
- Privet (2b, 3)
- Pyracantha (3)
- Rhododendron (1a, 2b, 3)
- Spirea (3)
- Viburnum (3, 4)
- Vitex (2b)  
(Chase Tree)

Flowering & Foliage Plants

- (Greenhouse [D])
- (Except California)
- African Violet (3)
- Azalea (1a, 2f, 3)
- Calendula (3, 4)
- Carnation (3, 4)
- Chrysanthemum (3, 4)
- Cineraria (3)
- Crassula (3)
- Daisy (3, 4)
- Fern, Boston (4)
- Desmella spp.
- Geranium (3, 4)
- Gerbera (3)
- Grape Leaf Ivy (3)
- Kalanchoe (3)
- Rose (3)
- Snapdragon (3, 4)

California Only Recommendations

- Greenhouse [D]
- African Violet (3)
- Carnation (3, 4)
- Chrysanthemum (3, 4)
- Geranium (3, 4)
- Grape Leaf Ivy (3)
- Rose (3)

- (1) Flower Blight
  - a) Ovulinia spp. [A]
  - b) Sclerotinia spp. [A]
  - c) Collectotrichum [A]
- (2) Leaf Blight/Spots
  - a) Cephalosporium spp. [C]
  - b) Cercoospora spp.
  - c) Didymellina spp. [B]
  - d) Didymascella thujina [G]
  - e) Entomosporium spp. [C]
  - f) Exobasidium spp. [E]
- (3) Powdery Mildew
  - Erysiphe spp.
  - Microsphaera spp.
  - Oidium spp.
  - Podosphaera spp.
  - Phylactinia spp.
  - Sphaerotheca spp.
  - Uncinula spp.
- (4) Rusts
  - a) Coleosporium spp.
  - b) Cronartium spp. [B]  
(Fusiform)
  - c) Gymnosporangium spp.
  - d) Melampsora spp. [F]
  - e) Melampsora farlowii [A]
  - f) Melampsorium spp.
  - g) Peridermium spp. [B]
  - h) Phragmidium andersonii
  - i) Puccinia spp.
  - j) Uromyces spp.
  - k) Uredinopsis mirabilis [A]
- (5) Tip Blight
  - Sirococcus strobilinus [B]

BEST AVAILABLE COPY

## ORNAIMENTAL PLANT DISEASE CONTROL (Continued)

**APPLICATION RATES:** Except as noted for specific diseases, mix 1 to 2 ounces of BAYLETON 25% Wettable Powder in 50 gallons of water and apply as a full coverage foliage spray to the point of drip as needed.

[A] Mix 4 to 8 ounces of BAYLETON 25% Wettable Powder in 50 gallons of water and apply as a full-coverage foliar spray to the point of drip. Applications should begin at the expanded bud stage (color showing). Use multiple applications at 7 to 14 day intervals as needed dependent upon bloom periods.

[B] Mix 8 ounces of BAYLETON 25% Wettable Powder plus sufficient spreader sticker for good coverage in 50 gallons of water. Apply in a spray application to the point of run-off on an as needed basis during the early part of the season. Excessive rates or excessive applications may result in a shortening of the flower stalk on iris.

[C] Mix 4 to 8 ounces of BAYLETON 25% Wettable Powder in 50 gallons of water and apply as a full coverage foliar spray to point of run-off. Apply in early spring as growth starts and re-apply on a 14 to 21 day interval until new growth is fully expanded. Protect new growth that develops in late summer or fall as temperatures begin to drop.

[D] Greenhouse Applications

Winter Use -- 1 ounce of BAYLETON 25% Wettable Powder

Summer Use -- 2 ounces of BAYLETON 25% Wettable Powder

Mix specified amount of BAYLETON 25% Wettable Powder in 50 gallons of water and apply in a spray application to the point of drip. Intervals between applications should be no shorter than 30 days to avoid flower stalk length reduction. Excessive rates or applications may result in a shortening of the flower stalk.

[E] For control of *Exobasidium* flower and leaf gall, apply 2 ounces of BAYLETON 25% Wettable Powder in 50 gallons of water. Begin application at bud break and apply at 10 day intervals through infestation period. ✓

[F] For control of *Melampsora pinitorqua* (Pine Twisting Rust), apply a single application in spring during periods favorable for infection. Mix 16 ounces in 50 gallons of water and apply to shoots in the upper whorl of susceptible pine species. Make a single application per year as a full coverage application sprayed to runoff.

[G] For control of *Didymascella thuja*, Cedar Leaf Blight, apply 1 lb per acre in sufficient water to provide full coverage in nurseries, or 0.5 lb/100 gallons applied as a full coverage spray to ornamentals. Begin applications before disease appears in spring, and repeat at 60 day intervals through early fall.

**COMPATIBILITY:** BAYLETON is compatible with many registered insecticides and fungicides. To determine the compatibility of BAYLETON with specific products, the following procedure should be conducted. Pour the recommended proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least 5 minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible.

**BEST AVAILABLE COPY**

**ORNAMENTAL PLANT DISEASE CONTROL (Continued)**

**SPRAY ADDITIVES:** Use of various spray additives such as spreaders, extenders, trace elements or fertilizers should be evaluated prior to use. The label directions given here are based on data obtained with no additives; use of any product with BAYLETON may affect the result. Contact local university extension personnel prior to use of spray mix additives.

**RESTRICTIONS:** Edible portions of treated trees, such as nuts and syrup, should not be used for feed or food.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**Household:** Do not reuse empty container. Securely wrap original container in several layers of newspaper and put in trash collection.

**Pesticide Disposal (Except Household):** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal (Except Household):** Completely empty carton into application equipment. Then dispose of empty carton in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Material that cannot be used as directed should be disposed of as directed above. In spill or leak incidents, keep unauthorized people away. You may contact the Miles Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Miles Kansas City Emergency Response Telephone No. is 816-242-2582, or contact Chemtrec at 800-424-9300.