

BAYLETON 25% Wettable Powder

318-7724.YLD

PM 22 3125-318

1042

Base Pre-Reg (7724)

Reason to Issue: To change PHI on apples
and pears to 45 days.

Date of Draft: 3/01/93 (Pre-Reg)(T)
Supersedes Draft Dated: 2/17/92

BAYLETON

25% Wettable Powder

Systemic Fungicide

ACTIVE INGREDIENT:


1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanone	25%
INERT INGREDIENTS	75%
	100%

U.S. Patent No. 3,912,752
EPA Reg. No. 3125-318
EPA Est. 3125-MO-1

Net Weight: ____ Pounds

AMENDMENT TO PREVIOUSLY REGISTERED LABELING

SHADED TEXT HAS NOT PREVIOUSLY BEEN REGISTERED



MILES

2072

RECOMMENDED APPLICATIONS											
CROP	DISEASE	RATE OF BAYLETON 25% Wettable Powder	REMARKS (Also Refer to Directions for Use)								
Apples	Cedar-apple rust	<u>Grams/5 gallon¹</u> 1.4 to 5.6 (1 level teaspoonful to 4 1/2 level teaspoonfuls)	Apply specified dosage per 5 gallons of water as a dilute spray applied to the point of drip. Make the first application at the green-tip stage and continue applications at 7-to 14-day intervals, or as needed until terminal growth ceases. Use the higher label rate under heavy disease pressure, with extended spray intervals, and/or on highly disease susceptible varieties. APPLICATION SCHEDULE * <table><tr><th><u>Rate (gms)</u></th><th><u>Maximum No. of Applications</u></th></tr><tr><td>1.4</td><td>12</td></tr><tr><td>2.8</td><td>6</td></tr><tr><td>5.6</td><td>3</td></tr></table> * A maximum of 16.8 grams of BAYLETON 25% Wettable Powder may be applied per season. BAYLETON 25% Wettable Powder may be applied up to 45 days prior to crop harvest.	<u>Rate (gms)</u>	<u>Maximum No. of Applications</u>	1.4	12	2.8	6	5.6	3
	<u>Rate (gms)</u>	<u>Maximum No. of Applications</u>									
1.4	12										
2.8	6										
5.6	3										
	Powdery mildew										
Pears	Powdery mildew	<u>Grams/5 gallon¹</u> 1.4 to 5.6 (1 level teaspoonful to 4 1/2 level teaspoonfuls)	Apply specified dosage per 5 gallons of water as a dilute spray applied to the point of drip. Make the first application at the budburst stage and continue applications at 7-to 14-day intervals, or as needed. Use the higher label rates under heavy disease pressure, with extended spray intervals, and/or on highly disease susceptible varieties. APPLICATION SCHEDULE * <table><tr><th><u>Rate (gms)</u></th><th><u>Maximum No. of Applications</u></th></tr><tr><td>1.4</td><td>12</td></tr><tr><td>2.8</td><td>6</td></tr><tr><td>5.6</td><td>3</td></tr></table> * A maximum of 16.8 grams of BAYLETON 25% Wettable Powder may be applied per season. BAYLETON 25% Wettable Powder may be applied up to 45 days prior to crop harvest.	<u>Rate (gms)</u>	<u>Maximum No. of Applications</u>	1.4	12	2.8	6	5.6	3
<u>Rate (gms)</u>	<u>Maximum No. of Applications</u>										
1.4	12										
2.8	6										
5.6	3										

¹ Rates of BAYLETON 25% Wettable Powder are based on a standard of 400 gallons of dilute spray solution per acre.

Miles Inc.
Specialty Products
Box 4013, Kansas City, Mo 64120-0013