318-8568.YL

U. S. LABEL

Base Pre-Reg (8568)

Reason to Issue: To add tank-mixing instructions with Prostar 50WP for control of brown patch.

Date of Draft: 07/07/95(Pre-RegicT)

### ACCEPTED

AUG 1 1 1995

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTER. ED UNDER NO. 3135-318

## **BAYLETON**°

### 25% Wettable Powder Systemic Fungicide

For control of certain diseases on flowers, foliage plants, shrubs, turf, fruit, melons, vegetables and shade trees. Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, home gardens, or on golf courses or lawns and grounds.

#### **ACTIVE INGREDIENT:**

1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanone	25%
INERT INGREDIENTS	<u>75%</u>
·	100%

U.S. Patent No. 3,912,752 EPA Reg. No. 3125-318 EPA Est. 3125-MO-1

VALETON

Net Weight: **Pounds** 

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read these entire Directions and Conditions of Sale before using **BAYLETON 25% Wettable Powder systemic** 

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE APPROPRIATE FOR THE CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED UNDER DIFFERENT ENVIRONMENTAL CONDITIONS. **BOTH INDOORS AND OUTDOORS UNDER** CONDITIONS SIMILAR TO THOSE THAT ARE ORDINARY AND CUSTOMARY WHERE THE PRODUCT IS TO BE USED.

INSUFFICIENT CONTROL OF PESTS OR PLANT INJURY MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR **UNUSUAL CONDITIONS, OR FROM FAILURE** TO FOLLOW LABEL DIRECTIONS. ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO ANIMALS, MAN, AND DAMAGE TO THE ENVIRONMENT. BAYER OFFERS, AND THE **BUYER ACCEPTS AND USES, THIS PRODUCT** SUBJECT TO THE CONDITIONS THAT **EXTRAORDINARY** OR UNUSUAL **ENVIRONMENTAL CONDITIONS, OR FAILURE** TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF BAYER AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

Do not formulate this product into other enduse products.



APPLICATION: BAYLETON fungicide is absorbed rapidly and works systemically from within the plant. Good coverage and wetting of the foliage are necessary. Rainfall or sprinkler irrigation, within 30 minutes after application does not decrease effectiveness. Control may be less effective on plants suffering from drought stress. Therefore, in order to achieve maximum control, plants should be maintained in a vigorously growing state through good cultural practices.

Apply in all cases when plants are fully established and actively growing. Applications should be applied at

....

recommended intervals to maintain disease control.

BAYLETON 25 Turf and Ornamental Fungicide can be tankmixed with Prostar 50WP for use on turf in accordance with the more (most) restrictive of label limitations and procautions. No label dosage rates should be exceeded. This product carnot be mixed with any product comaining a label prohibition against such mixing.

RECOMMENDED APPLICATIONS TURF GRASS DISEASE CONTROL						
DISEASE	OZ OF BAYLETON 25% WP PER 1,000 SQUARE FEET		REMARKS			
Brown Petch/Riszoctone Biglit (Phirocronic entent) (Recommendations for tank-nex with Proster 50WP)	<b>Exercise:</b>	CARATON	Preventative: Tank treat with Proster 50WP and stoky on a 21 to 28 day achieving. Follow the directions on the Proster 50WP table when tank mixing:  Curetive: To control existing infections, tank mix BAYLETON 25 with the puretive rate of Proster 50WP. Subsequent applications should be applied on a preventive schedule using the preventive rate of Proster 50WP.			

Sayer Corporation
Specialty Products
Sox 4913, Kansas City, Mo 64120-0013

"

AYLETON

Reason to Issue: To add tank-mixing instructions with Prostar 50WP for control of brown patch.

## **BAYLETON®**

# 25% Wettable Powder Systemic Fungicide

Date of Draft: 07/07/95(Pre-Reg)(T)

ACCEPTED

AUG | 1 1995

under the federal insecticide fundicide and rodenticide act for economic poison registered under no. 3135 - 218

For control of certain diseases on flowers, foliage plants, shrubs, turf, fruit, melons, vegetables and shade trees. Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, home gardens, or on golf courses or lawns and grounds.

#### **ACTIVE INGREDIENT:**

1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1 <i>H</i> -1,2,4-triazol-1-yl)-2-butanone	25%
INERT INGREDIENTS	75%
	100%

U.S. Patent No. 3,912,752 EPA Reg. No. 3125-318 EPA Est. 3125-MO-1

Net Weight: \_\_\_ Pounds

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read these entire Directions and Conditions of Sale before using BAYLETON 25% Wettable Powder systemic fungicide.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE APPROPRIATE FOR THE CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED UNDER DIFFERENT ENVIRONMENTAL CONDITIONS BOTH INDOORS AND OUTDOORS UNDER CONDITIONS SIMILAR TO THOSE THAT ARE ORDINARY AND CUSTOMARY WHERE THE PRODUCT IS TO BE USED.

INSUFFICIENT CONTROL OF PESTS OR PLANT INJURY MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL CONDITIONS, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO ANIMALS, MAN, AND DAMAGE TO THE ENVIRONMENT. BAYER OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT **EXTRAORDINARY** UNUSUAL OR **ENVIRONMENTAL CONDITIONS, OR FAILURE** TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF BAYER AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

Do not formulate this product into other enduse products.



Unregistered label text is highlighted in redlining.

1.71 4

APPLICATION: BAYLETON fungicide is absorbed rapidly and works systemically from within the plant. Good coverage and wetting of the foliage are necessary. Rainfall or sprinkler irrigation, within 30 minutes after application does not decrease effectiveness. Control may be less effective on plants suffering from drought stress. Therefore, in order to achieve maximum control, plants should be maintained in a vigorously growing state through good cultural practices.

Apply in all cases when plants are fully established and actively growing. Applications should be applied at

recommended intervals to maintain disease control.

BAYLETON 25 Turf and Ornamental Fungicide can he tankmixed with Prostar 50WP for use on turf in accordance with the more (most) restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

RECOMMENDED APPLICATIONS TURF GRASS DISEASE CONTROL						
DISEASE	OZ OF BAYLETON 25% WP PER 1,000 SQUARE FEET		REMARKS			
Brown Patch/Rhizoctoma Blight (Rhizoctoma solarii) (Recommendations for tank-mix with Prostar 50WP.)	Preventative	Curative 3	Preventatives: Tank-mix with Proster SOWP and apply on a 21 to 28 day schedule; Follow use directions and restrictions on the Proster SOWP label when tank-mixing;  Curative: To control existing infections, tank-mix BAYLETON 25 with the curative rate of Proster SOWP. Subsequent applications should be applied on a preventive schedule using the preventive rate of Proster SOWP.			

## **BEST COPY AVAILABLE**

Sayer Corporation Specialty Products Sox 4913, Kansas City, Mo 64120-0013