

# BAYLETON 25% WP

318-8697 LD

U. S. LABEL

Base Pm Reg (8695)

Reason to Issue: To revise directions for use on bermudagrass decline and take all patch.

Date of Draft: 01/03/96(T)

## BAYLETON®

### 25% Wettable Powder Systemic Fungicide

For control of certain diseases on flowers, foliage plants, shrubs, turf, fruit, melons, vegetables and shade trees.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, home gardens, or on golf courses or lawns and grounds.

ACTIVE INGREDIENT:

1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanone . . . . . 25%

INERT INGREDIENTS . . . . . 75%

100%


U.S. Patent No. 3,912,752

EPA Reg. No. 3125-318

EPA Est. 3125-MO-1

Net Weight:      Pounds

#### AMENDMENT TO PREVIOUSLY REGISTERED LABELING



Unregistered label text is highlighted in redlining  
Text to be deleted is indicated by ~~strikethrough~~

# RECOMMENDED APPLICATIONS TURF GRASS DISEASE CONTROL

DISEASE	OZ OF BAYLETON 25% WP PER 1,000 SQUARE FEET*		REMARKS
Bermudagrass decline <i>(Gaeumannomyces graminis var. graminis)</i>  Take all patch <i>(Gaeumannomyces graminis var. avenae)</i>  (Except California)	<b>Preventive</b>  2 - 4	<b>Curative</b>  4	<p>Immediately after the fungicide is applied, the area should be thoroughly irrigated to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone. The objective is to water the fungicide into the crown and root zone.</p> <p><b>PREVENTIVE RATE:</b> Begin applications <del>2 to 4 weeks</del> prior to the appearance of disease symptoms. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information. Apply subsequent applications at 21- to 28-day intervals. For take all patch, applications in both spring and fall may be necessary.</p> <p><b>CURATIVE RATE:</b> To control existing infections, apply 4 oz <del>at 14- to 21-day intervals</del> for the initial <del>4 or 2</del> treatments followed by 2 to 4 oz at 21- to 28-day intervals. Cultural control practices such as aerification, topdressing, reseeding, and fertilization should <del>must</del> be implemented prior to or at the same time the fungicide is applied. Refer to your local County Extension Service for this information.</p>

Bayer Corporation  
Specialty Products  
Box 4913, Kansas City, Mo. 64120-0013