

Reg # 3125-320

PM-22

1984

	316	79696
18	<u>37</u>	<u>1</u>

	316	84329
MAY 25 1986	<u>37</u>	<u>1</u>

Mobay Chemical Corporation
Agricultural Chemicals Division
P.O. Box 4913
Hawthorn Road
Kansas City, MO 64120

Attention: Mr. Glenn Brussell

Gentlemen:

Subject: Bayleton 50% Wettable Powder
EPA Registration No. 3125-320
Use on Grasses Grown for Seed
Your Application Dated May 27, 1982

The amendment referred to above, submitted in connection with registration under FIFRA sec. 3(c)(7)(B), is acceptable, provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.

2. Submit production information (pounds or gallons produced) for this product for the fiscal year in which the use(s) on grasses grown for seed is (are) conditionally registered, in accordance with FIFRA sec. 29. The production information will be submitted to the Agency no later than November 15, following the end of the preceding fiscal year.

This information should be submitted to:

Process Coordination Branch
Registration Division (TS-767)
Environmental Protection Agency
Washington, D.C. 20460

3. Make the following labeling changes before you release the product for shipment bearing the amended labeling:

a. Under "Directions for Use" add a statement similar to the following:

CONCURRENCES							
SYMBOL ▶							
SURNAME ▶							
DATE ▶							

2 2 4

..2-

Note to user: For worker protection during application, wear a hat, long sleeved shirt, and long legged trousers or overalls. Protective clothing should be removed and laundered after use.


- b. Submit five copies of final printed labeling showing all accepted use patterns as of this date before you release the product for shipment.

4. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,



Henry M. Jacoby
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (TS-767)

Enclosure

U.S. LABEL

Reason to Issue: To allow feeding of chaff and straw from treated grass seed crops; 11/10/82 Draft: revise rotational crop statement; modify format.

Date of Draft: 11/10/82 (Pre-Reg)(S)

Supersedes Pre-Reg Draft Dated: 9/13/82

EPA Reg. No. 3125-320

® BAYLETON

ACCEPTED
with COMMENTS
in EPA Letter Dated:

50% Wettable Powder

FOR CONTROL OF SPECIFIED DISEASES ON CERTAIN
FIELD CROPS

MAR 25 1983

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

31250-320

ACTIVE INGREDIENT:

1-(4-Chlorophenoxy)-3,3-dimethyl-

1-(1H-1,2,4-triazol-1-yl)-2-

butanone 50%

AMFNDMENT TO PREVIOUSLY REGISTERED LABELING

APPLICATIONS

BAYLETON 50% Wettable Powder may be applied by ground or aircraft spray equipment. Check equipment calibration frequently. Complete coverage and thorough application is essential for the most effective results, especially when lower spray volumes are applied.

Grasses Grown for Seed

BAYLETON is absorbed rapidly and works systemically from within the plant. Good coverage and wetting of the foliage is necessary. For maximum yield protection early treatment of infections is necessary. Control may be less effective on plants suffering from drought stress.

MIXING

Add the recommended amount of BAYLETON 50% Wettable Powder into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the BAYLETON 50% Wettable Powder should be thoroughly dispersed prior to the addition of other materials.

COMPATIBILITY

BAYLETON is physically compatible with many registered insecticides, miticides, and fungicides. To determine the compatibility of BAYLETON with other products, the following procedure should be followed: Pour the recommended proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least five (5) minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible. For further information contact your local Mobay representative.

DOSAGE

Use the specified amount of BAYLETON 50% Wettable Powder per acre in sufficient water to give complete coverage of foliage. Use the higher recommended rates on the most disease susceptible varieties and/or with increasing disease pressure.

RECOMMENDED APPLICATIONS

Consult your State Agricultural Experiment Station or Extension Service Specialist or Mobay representative for guidance on the selection of specific dosage rates and timing of applications appropriate for your specific crop and disease. State regulations may contain additional restrictions or requirements. Check the Remarks and Restrictions sections of this label for additional directions.

CROP	DISEASE	Rate BAYLETON 50% WP	REMARKS
Grasses Grown for Seed Perennial ryegrass Kentucky bluegrass	Rusts (Puccinia species)	8 to 16 oz.	Apply specified dosage per acre in at least 20 gallons of water per acre with ground sprayers, or in 7 to 10 gallons of water per acre by aircraft. Apply when rust pustules become readily noticeable and are increasing in number in late spring or early summer. Rust control is especially important during the period from early head emergence through last flowering. Under severe rust pressure, use high rates (12 to 16 ozs. per acre) and/or more frequent applications to maintain rust control until the grass seed matures, but a maximum of 2 lbs. of BAYLETON 50% WP may be applied per year. THE LAST APPLICATION MAY BE MADE UP TO 5 DAYS OF HARVEST. CHAFF AND STRAW FROM TREATED AREA MAY BE USED FOR FEED PURPOSES, HOWEVER, DO NOT FORAGE OR CUT GREEN CROP OR USE SEED FOR FEED PURPOSES. REGROWTH MAY BE GRAZED 11 WEEKS AFTER LAST APPLICATION.

ROTATIONAL CROPS

Small grains, corn, sorghum, soybeans, beans and peas may be planted 35 days after the last application of BAYLETON, however forage or vines from these crops may not be used for food or feed. Root crops may be planted 120 days after the last application but tops must not be used for food or feed. All crops may be planted 12 months or later after the last application of BAYLETON without any restrictions.