

BAYL1

Reg # 3125 - 320

PM-22


12/1

U.S. LABEL

Reason to Issue: To claim powdery
mildew control on grasses grown
for seed.

Date of Draft: 4/15/83 (Pre-Reg) (S)

EPA Reg. No. 3125-320



® BAYLETON

50% Wettable Powder
Fungicide

ACTIVE INGREDIENT:

1-(4-Chlorophenoxy)-3,3-dimethyl-1H-
-1,2,4-triazol-1-yl)-2-butanone 50%

AMENDMENT

To Previously Registered Labeling

Revise as Follows:

RECOMMENDED APPLICATIONS

CROP	DISEASE	Rate BAYLETON 50% WP	REMARKS
Grasses grown for Seed			Apply specified dosage per acre in at least 20 gallons of water per acre with ground sprayers, or in 7 to 10 gallons of water per acre by aircraft.
Perennial ryegrass Kentucky bluegrass	Rusts (Puccinia species)	8 to 16 oz.	<u>Rusts:</u> Apply when rust pustules become readily noticeable and are increasing in number in late spring or early summer. Rust control is especially important during the period from early head emergence through last flowering. Under severe rust pressure, use high rates (12 to 16 ozs. per acre) and/or more frequent applications to maintain rust control until the grass seed matures.
	Powdery mildew	2 to 4 oz.	<u>Powdery Mildew:</u> One or two applications at the 2 to 4 oz rate applied at the first signs of disease and when new powdery mildew symptoms reappear are generally sufficient for disease control. In cases where severe mildew infections are present, use the higher rate. A maximum of 2 lbs. of BAYLETON 50% WP may be applied per acre per year. THE LAST APPLICATION MAY BE MADE UP TO 5 DAYS OF HARVEST. CHAFF AND STRAW FROM TREATED AREA MAY BE USED FOR FEED PURPOSES, HOWEVER, DO NOT FORAGE OR CUT GREEN CROP OR USE SEED FOR FEED PURPOSES. REGROWTH MAY BE GRAZED 1 WEEKS AFTER LAST APPLICATION.