

DEC 3 1993

John S. Thornton
Miles Inc.
P.O. Box 4913
Kansas City, MO 64120

Dear Mr. Thornton:

Subject: Delete Use on Barley and Grasses Grown for Seed
Bayleton 50% Wettable Powder
EPA Registration No. 3125-320
Your Submission Dated May 4, 1993

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended is acceptable provided that you:

1. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

a. Modify the mancozeb tank mix directions so they are in compliance with the EBDC PD-4 published in the Federal Register on March 2, 1992.

b. Move the statement "Do not use on other crops grown for food or forage" from the Environmental Hazards section to the Directions for Use.

2. Submit one (1) copy of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

The amended labeling supersedes all previously accepted ones.

-2-

This labeling does not bear the required Worker Protection statements as outlined in PR Notices 93-7 and 93-11. We note that you have submitted labels in accordance with the instructions in those Notices. The new protective language in those submissions acceptance on October 22, 1993 must be incorporated into this label for any shipment released after the deadlines that have been established. Note finished printed labels must also be submitted in accordance with the instructions of the WPS group.

Sincerely yours,

15

Cynthia Giles-Parker
Product Manager (22)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

BAYLETON 50% Wetttable Powder

Base Pre-Req. (7739)

Date of Draft: 3/22/93(Pre-Reg)(T)

BAYLETON

50% Wettable Powder

Fungicide

**FOR CONTROL OF SPECIFIED DISEASES ON CERTAIN FIELD,
FRUIT AND VEGETABLE CROPS**

1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1 <i>H</i> -1,2,4-triazol-1-yl)-2-butanone	50%
INERT INGREDIENTS	50%
	100%

Net Weight: **Pounds**

STOP - READ THE LABEL BEFORE USE

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

See Rear Panel for Statements of Practical Treatment and Other Precautionary Statements

ACCEPTED
W. J. C. M. VTS
100 N. Main St., Q. ed:

DEC 3 1993

3125-320

MILES 

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled, or absorbed through the skin. Prolonged or frequently repeated skin contact may cause allergic skin reactions in some individuals. Avoid

contact with skin, eyes or clothing. Avoid breathing dust or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENTS OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger, or, if available, by administering syrup of ipecac. If syrup of ipecac is available, administer 1 tablespoon (15 mL) of syrup of ipecac followed by 1 to 2 glasses of water. If vomiting does not occur within 20 minutes, repeat dose once. Do not induce vomiting or give anything by mouth to an unconscious person.

If on skin: Wash with plenty of soap and water. Get medical attention.

If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, mouth to mouth. Get medical attention.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

SYMPTOMS OF POISONING: The compound does not cause any definite symptoms that would be diagnostic. Poisoning is accompanied by hyperactivity followed by sedation.

NOTE TO PHYSICIAN: No specific antidote. Treat symptomatically.

ENVIRONMENTAL HAZARDS

Do not use on other crops grown for food or forage. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

Do not make applications when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read these entire Directions and Conditions of Sale before using BAYLETON 50% Wettable Powder fungicide.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE THE DIRECTIONS FOR CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED FOR A RANGE OF WEATHER CONDITIONS SIMILAR TO THOSE WEATHER CONDITIONS THAT ARE ORDINARY AND CUSTOMARY IN THE GEOGRAPHIC AREA WHERE THE PRODUCT IS USED. INSUFFICIENT CONTROL OF PESTS AND/OR INJURY TO THE CROP TO WHICH THE PRODUCT IS APPLIED MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL

WEATHER, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO OTHER CROPS, ANIMALS, MAN, OR THE ENVIRONMENT. MILES OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL WEATHER, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF MILES AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

BEST AVAILABLE COPY

DIRECTIONS FOR USE (Continued)

MIXING: Add the recommended amount of BAYLETON 50% Wettable Powder into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the BAYLETON 50% Wettable Powder should be thoroughly dispersed prior to the addition of other materials.

COMPATIBILITY: BAYLETON fungicide is physically compatible with many registered insecticides, miticides, and fungicides. To determine the compatibility of BAYLETON fungicide with other products, the following procedure should be followed: Pour the recommended proportions of the products into a suitable container of

water, mix thoroughly and allow to stand at least five (5) minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible. For further information contact your local Miles representative.

NOTE TO USER: For worker protection during mixing, loading, and during application, wear a hat, long sleeve shirt, and long legged trousers or overalls. In addition, during mixing and loading, wear rubber or neoprene gloves and a dust mask. Protective clothing should be laundered separately following application.

RECOMMENDED APPLICATIONS

BAYLETON 50% Wettable Powder may be applied in a minimum of 20 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment except as noted under "Remarks" for each crop. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage.

Do not apply this product through any type of irrigation system.

Consult your State Agricultural Experiment Station or Extension Service Specialist or Miles representative for guidance on the selection of specific dosage rates and timing of applications appropriate for your specific crop and disease. State regulations may contain additional restrictions or requirements. Check the Remarks and restrictions sections of this label for additional directions.

BEST AVAILABLE COPY

RECOMMENDED APPLICATIONS				
CROP	DISEASE	RATE OF BAYLETON 50% Wettable Powder		REMARKS (Also refer to Directions for Use)
FRUITS Apples	Cedar-apple rust Powdery mildew	<u>Oz/A</u>	<u>Oz/100 gal¹</u>	Make the first application at the green-tip stage and continue applications at 7- to 14-day intervals, or as needed until terminal growth ceases. Use the higher label rate under heavy disease pressure, with extended spray intervals, and/or on highly disease susceptible varieties. A maximum of 24 ounces of BAYLETON 50% W.P. may be applied per acre per season. BAYLETON 50% W.P. may be applied up to 45 days prior to crop harvest. Do not graze livestock in treated orchards.
		2 - 8	1/2 to 2	
Pears	Powdery mildew	2 to 8	1/2 to 2	Make the first application at the budburst stage and continue applications at 7- to 14-day intervals, or as needed. Use the higher label rates under heavy disease pressure, with extended spray intervals, and/or on highly disease susceptible varieties. A maximum of 24 oz of BAYLETON 50% W.P. may be applied per acre per season. BAYLETON 50% W.P. may be applied up to 45 days prior to crop harvest. Do not graze livestock in treated orchards.
Grapes	Powdery mildew	2 to 6 oz/A		GENERAL COMMENTS Use the higher label rates under heavy disease pressure, with extended spray intervals, and/or on highly disease susceptible varieties. Fields should be observed closely for early disease symptoms when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. A maximum of 18 ounces of BAYLETON 50% W.P. may be applied per acre per crop season. BAYLETON 50% W.P. may be applied up to 14 days before harvest. POWDERY MILDEW: Make the first application pre bloom and continue applications at 14- to 21-day intervals, or as needed. BLACK ROT Protective Schedule: Make the first application at 10-inch green shoot and continue applications at 7- to 14-day intervals through 5° Brix stage or until veraison (berry coloring) is complete. Under severe disease conditions and/or on highly susceptible varieties, make applications at 7-day intervals. Post-infection Schedule: During bloom apply within 72 hours after the beginning of an infection period. Do not make spray applications any closer than 7-day intervals. After bloom, continue applications at rates of 2 to 3 ounces of BAYLETON 50% W.P. at 7- to 14-day intervals, or as needed to maintain control.
	Black rot	Protective Schedule 2 to 3 oz/A Post-infection 3 to 4 oz/A		

¹ Rates of BAYLETON 50% Wettable Powder are based on a standard of 400 gallons of dilute spray per acre, or the equivalent amount of product per acre in a concentrate spray.

¹ Rates of BAYLETON 50% Wettable Powder are based on a standard of 400 gallons of dilute spray per acre, or the equivalent amount of product per acre in a concentrate spray.

RECOMMENDED APPLICATIONS (Continued)			
CROP	DISEASE	RATE OF BAYLETON 50% Wettable Powder	REMARKS (Also Refer to Directions for Use)
Pineapple (Fresh market only)	Fresh fruit rot <i>Ceratocystis paradoxa</i>	13-1/3 oz	Post-Harvest Treatment: Apply specified dosage in 100 gallons of water to fruit after harvest. Fruit should be dipped or sprayed for thorough coverage and allowed to drain. Limit dipping time to not more than 3 minutes.
Raspberries (California Only)	Powdery mildew	4 oz/A	Apply specified dosage per acre in not less than 20 gallons of water using ground equipment. Make the first application when disease symptoms first appear and additional applications at 4- to 6-week intervals thereafter as needed. A maximum of 7 total applications may be made in one year with not more than 2 applications within the same 30-day period. Do not harvest within 1 day of application.
SMALL GRAINS Wheat	Powdery mildew	2 to 6 oz/A	Field should be observed closely for early disease symptoms when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. Apply specified dosage per acre in a minimum of 10 gallons of water per acre by ground equipment or a minimum of 5 gallons of water per acre by air when disease symptoms first appear on leaves or stems. Additional applications should be made if new disease symptoms appear. The total amount of BAYLETON 50% W.P. should not exceed 16 oz per acre per crop season and the last application should not be made within 21 days of harvest. Powdery mildew: In areas having severe powdery mildew infection apply 4 to 6 oz of BAYLETON 50% W.P. per acre.
	Rusts (stripe, leaf, and stem) Speckled leaf blotch (<i>Septoria tritici</i>) Leaf and glume blotch (<i>Septoria nodorum</i>) Tan spot (<i>Pyrenophora tritici-repentis</i>)	4 to 8 oz/A	Rusts: In areas having severe rust infection apply 6 to 8 oz of BAYLETON 50% W.P. per acre. Leaf and glume blotch, speckled leaf blotch, and tan spot: BAYLETON 50% W.P., applied at 4 to 8 oz per acre at first sign of infection, will control light to moderate leaf and glume blotch, speckled leaf blotch, and tan spot infections. Applications for the control of glume blotch should be timed to coincide with early head emergence. Applications made prior to head emergence will not provide sufficient residual activity to control glume blotch. For control of severe infections, it is recommended that a registered fungicide with good activity against these diseases be tank-mixed with BAYLETON. Refer to the Tank Mixes section of this product label.

BEST AVAILABLE COPY

RECOMMENDED APPLICATIONS (Continued)			
CROP	DISEASE	RATE OF BAYLETON 50% Wettable Powder	REMARKS (Also refer to Directions for Use)
Pine (Seedlings) (Except California)	Pine Rust (Fusiform rust)	4 to 16 oz/A	Begin application prior to infection period and repeat as necessary at 2- to 3-week intervals depending upon disease pressure. Use lower rates in areas of low disease incidence and higher rates in areas of severe disease incidence. A maximum of 64 oz of BAYLETON 50% W.P. may be applied per acre per season. A spreader-sticker is needed to help adhere spray solution to the pine trees.
Pine Seed (Nurseries) (Except California)	Fusiform rust (<i>Cronartium queruum</i>)	20 oz	Mix specified dosage in 100 gallons of water. Cover seeds with the mixture and soak them, at room temperature (75°F), for 24 hours, stirring occasionally. Seed stratification, if necessary, should be completed before treatment with BAYLETON 50% Wettable Powder. Thoroughly air dry all seed before applying any additional treatment or before planting. A spreader sticker may be needed to help adhere BAYLETON to the pine seed. Do not use treated seed for food or feed purposes.
		2 oz	Apply specified dosage to 50 lb of thoroughly wetted pine seeds in a commercial treater or other suitable tumbler apparatus. Allow to mix for at least 10 minutes before applying bird repellent or other seed dressing materials. Thoroughly air dry seed before sowing. Do not use treated seed for food or feed purposes.
Sugar beets	Powdery mildew	BROADCAST 4 to 16 oz/A	<p>GENERAL COMMENTS: Apply the specified dosage per acre in a minimum of 10 gallons of water by ground and 5 gallons of water by air.</p> <p>Under severe disease conditions, use of the higher recommended rate may be necessary.</p> <p>Do not apply more than 16 oz of BAYLETON 50% Wettable Powder per acre per season.</p> <p>BROADCAST APPLICATIONS: Apply specified dosage per acre as a broadcast foliar spray when disease symptoms first appear. To maintain control or if reinfection occurs, additional applications may be made at 10- to 14-day intervals.</p> <p>Broadcast applications may be made up to 15 days before harvest.</p>
		BAND 8 to 16 oz/A	<p>BAND APPLICATIONS: Apply specified dosage per acre as a spray directed into the leaf whorl in a 2-inch band by turning a flat fan spray nozzle sideways to achieve a narrow band.</p> <p>Band applications may be made up to 30 days before harvest.</p>

BEST AVAILABLE COPY

RECOMMENDED APPLICATIONS (Continued)

CROP	DISEASE	RATE OF BAYLETON 50% Wettable Powder	REMARKS (Also refer to Directions for Use)
Cucurbits [Balsam pear (bitter melon), Chinese waxgourd (Chinese preserving melon), citron melon, cucumber, gherkin, gourds, edible melons including hybrids (including cantaloupe, casaba, crenshaw, honeydew melons, honey balls, mango melon, muskmelon, Persian melon), pumpkin squash, summer squash, winter watermelon, including hybrids]	Powdery mildew (<i>Erysiphe cichoracearum</i>)*	2 to 4 oz/A	Apply specified dosage per acre in a minimum of 10 gallons of water by ground and 5 gallons of water by air as a foliar spray when disease symptoms first appear. Repeat at 14-day intervals to maintain control if new disease symptoms appear. Under severe disease conditions use the 4 oz rate. Applications may be made up to day of harvest. A maximum of 16 oz of BAYLETON 50% W.P. may be applied per acre per season. *BAYLETON 50% W.P. will not give commercially acceptable control of <i>Sphaerotheca fuliginea</i> .
Christmas Trees (Except Concolor Fir)	Stem & Cone Rusts <i>Cronartium</i> spp. (Fusiform) <i>Peridermium</i> spp. <i>Endocronartium harknessii</i> (Gall) Tip blight <i>Sirococcus strobilinus</i>	8 oz/A	Apply specified dosage per acre or per 100 gallons of water as a full coverage, dilute spray as needed. Full coverage of the trees is essential for maximum control. Use of nonionic spray adjuvant is recommended. Time applications appropriately for the specific disease being controlled. A maximum of 64 oz of BAYLETON 50% W.P. may be applied per acre per season. For rusts, begin applications when the needles break through the fascicle sheath. Make additional applications at 14- to 21-day intervals. Stop when galls become pale to white in color. For tip blight, begin applications to coincide with bud break. Make two additional applications at 14-day intervals.

BEST AVAILABLE COPY

TANK MIXES

The following tank mix recommendations are made for the improved control of specific diseases. Before mixing BAYLETON 50% Wettable Powder with any other fungicide, read and carefully observe the precautionary statements, directions for use, and other information appearing on the EPA registered product labels.

CROP	Rate of 50% Wettable Powder and 50% Dry Flowable Formulations of Benlate®	For Improved Control of:	BAYLETON 50% W.P. Rate
Wheat	4 oz/A	Speckled Leaf blotch Leaf and glume blotch Leaf rust Tan spot	2 oz/A
CROP	Rate of 80% Wettable Powder* Formulations of Mancozeb	For Improved Control of:	BAYLETON 50% W.P. Rate
Grapes	1 1/2 to 4 lb/A	Bunch rot Dead arm Downy mildew	2 to 6 oz/A
Sugar beets	1 1/2 to 2 lb/A	Cercospora leaf spot	4 to 16 oz/A
Wheat	2 lb/A	Helminthosporium leaf spots Speckled leaf blotch Leaf and glume blotch Tan spot Rusts	2 oz/A

* The rates given are for the following 80% wettable powder formulations of mancozeb: Dithane M-45®, Penncozeb®, Clean Crop®, Mancozeb 80 WP and Mancozeb Fungicide by Riverside/Terra Corp. Only tank mixes with Dithane M-45 can be used in California.

ROTATIONAL CROPS

Small grains, corn, sorghum, soybeans, beans, peas and cucurbits may be planted 35 days after the last application of BAYLETON, however forage or vines from these crops may not be used for food or feed. Root crops

may be planted 120 days after the last application but tops must not be used for food or feed. All crops may be planted 12 months or later after the last application of BAYLETON without any restrictions.

BEST AVAILABLE COPY**RE-ENTRY STATEMENTS**

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

(The following may appear on supplemental labeling accompanying the product) Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product.

Oral warnings should include those statements found under the label heading "Precautionary Statements" and "Re-entry Statements" included on this label. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: WARNING, area treated with BAYLETON 50% Wettable Powder on (date of application). Do not enter without appropriate protective clothing until sprays have dried. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash with flowing water for at least 15 minutes. If swallowed, vomiting should be induced. (See Precautionary Statements on label).

STORAGE AND DISPOSAL

Pesticide Disposal: Do not contaminate water, food or feed by storage or disposal.

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticide as directed above. In spill or leak incidents, keep unauthorized people away. You may contact the Miles Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Miles Kansas City Emergency Response Telephone No. is 816-242-2582, or contact Chemtrec at 800-424-9300.

Benlate is a Reg. TM of E.I. DuPont de Nemours & Co. (Inc.)
Clean Crop is a Reg TM of Platte Chemical Company.
Dithane M-45 is a Reg. TM of Rohm and Haas Company.
Penncozeb is a Reg. TM of Elf Atochem North America.

BEST AVAILABLE COPY

Miles Inc.
Crop Protection Products
Box 4913, Kansas City, MO 64120