

U.S. LABEL

Reason to Issue: Revise claims for use of SENCOR in a tank mix with 2,4-D LVE on soybeans.

Date of Draft: 9/22/92 (Pre-Reg)(H)

EPA Reg. No. 3125-325

SENCOR DF
75% Dry Flowable Herbicide


ACTIVE INGREDIENT:

4-Amino-6-(1,1-dimethylethyl)-3-(methylthio)-1,2,4-triazin-5(4H)-one 75%

INERT INGREDIENTS: 25%
100%

AMENDMENT TO CURRENTLY REGISTERED LABELING

REVISE TO READ AS FOLLOWS:



3 of 3

SENCOR DF 75% Dry Flowable Herbicide

2,4-D LVE herbicide use rates for Reduced-Tillage Combinations

Method of Application	2,4-D LVE pt/ac ^{1,2,3,4,5,6}	Minimum Preplant Interval, Days
Preplant	1/2 1	7 30

¹The application rate of 2,4-D LVE herbicide is based on, but not restricted to, a product containing 3.8 pounds acid equivalent (a.e.)/gallon. The equivalent rates in pounds acid equivalent would be: 1/2 pt/ac = 0.24 lb a.e./ac and 1 pt/ac = 0.48 lb a.e./ac.

²2,4-D LVE herbicide is Isooctyl (2-ethylhexyl) ester of 2,4-Dichlorophenoxyacetic Acid (e.g., Esteron 99 or Salvo herbicide).

³Flat fan nozzles that minimize drift are recommended. Use a minimum of 10 gallons total spray volume per acre with ground equipment or a minimum of 5 gallons total spray volume per acre with aerial equipment.

⁴Make only one application of 2,4-D LVE herbicide per growing season, regardless of the application rate used.

⁵When 2,4-D LVE herbicide is applied, do not feed hay, forage, or fodder. Restrict livestock from grazing treated fields. Livestock should be restricted from feeding/grazing of treated cover crops.

⁶Refer to the 2,4-D LVE herbicide label for precautionary statements, restrictions and mixing instructions. Read the crop injury statements on the 2,4-D LVE herbicide label and use only if these statements are acceptable.

Refer to this label and the 2,4-D LVE, Dual, Lasso, and Prowl labels for additional weeds controlled, precautions, restrictions, limitations and sprayer clean-up information.

Observe all cautions and limitations on labeling of all products used in mixtures.