300 330

YLD28

U.S. LABEL

Reason to Issue: To move Hyperodes weevil from the Soil Insects list

to the Surface Insect list.

EPA Reg. No. 3125-330

Date of Draft: 4/15/86 (Pre-Reg) (S)

® OFTANOL 5% Granular INSECTICIDE

ACTIVE INGREDIENT:

1-Methylethyl 2-[[ethoxy[(1methylethyl)amino] phosphinothioyl]oxy]benzoate ...... 5%

**AMENDMENT** 

To Previously Registered Labeling

Delete: Hyperodes weevil from Soil Insects

Amend: Surface Insects as follows:

ACCEPTED

40,90 1 A1

JUN 9 1986

Under the Federal Insecticide, Fungicide, and Rodenticide Act, registered under resistered values as amended, for the pesticide

## RECOMMENDED APPLICATIONS

CROR	DECT	DOSAGE	DEWARKS
CROP	PEST	OFTANOL 5% G	REMARKS
Turf Grasses	SURFACE INSECTS Sod webworm larvae Chinch bug Hyperodes weevil	40 lbg/A or 0.9 lbs./ 1000 sq. ft.	Apply 1/4 to 1/2 inch of water immediately after treatment. Reapply as necessary for pest re-infestations. Do not apply more than 80 lbs. of OFTANOL 5% Granular per acre per year or an equivalent amount of 4 lbs. active ingredient per acre per year regardless of formulation used. Minimum interval between applications must be 30 days.  For Hyperodes Weevil Control: Apply between full bloom periods of forsythia and flowering dogwood, usually April 15; to May 15, and water-in as directed
	BEST AVAILABLE	COPY	Consult your local turf, state Agricultural Experiment Station or state Extension Service Specialist for more specific information regarding timing of application.