

338-6466.YLD

Base Pre-Reg (6466)

U.S. LABEL

Reason to Issue: To allow for application to cherries.

Date of Draft: 6/20/88 (Pre-Reg) (M)

EPA Reg. No. 3125-33E

**RESTRICTED USE PESTICIDE**  
**Due to Acute Toxicity**

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

**ACCEPTED**  
**AUG 28 1990**

Under the Federal Insecticide, Fungicide, and Rodenticide Act  
 Registration No. 3125-33E

GUTHION® 3 Flowable  
INSECTICIDE

ACTIVE INGREDIENT:  
O,O-Dimethyl S[(4-oxo-1,2,3-benzotriazin-3(4H)-yl)methyl] phosphorodithioate ..... 30%

AMENDMENT  
To Previously Registered Labeling

Revise as Follows:

RECOMMENDED APPLICATIONS

CROP	PEST	Pints GUTHION 3F	REMARKS
FRUIT			
Cherries (East of Rocky Mountains)	Eye-spotted bud moth Forbes scale Fruit flies Fruit tree leaf roller Lesser peach tree borer Plum curculio San Jose scale* Cherry leaf miner Mineola moth	2  (2/3 pints/ 100 gallons)	Apply specified dosage per acre as a full coverage spray.  Up to four applications may be applied per acre per crop season.  Allow at least 14 days between applications and 21 days between the last application and harvest.

\*In some areas, these species may have developed resistance to organophosphate insecticides. GUTHION insecticide used alone may not provide satisfactory control in those areas. Consult your local agricultural advisor or Cooperative Extension Service for recommendations.

GUTHION 3 Flowable

RECOMMENDED APPLICATIONS (Cont'd)

CROP	PEST	Pints GUTHION 3F	REMARKS
FRUIT (Cont'd)			
Cherries (West of Rocky Mountains)	Leafhoppers Fruit flies Fruit tree leaf roller Lesser peach tree borer Cherry leaf miner	1-1/3  (7 fluid ounces per 100 gallons)	Apply specified dosage per acre as a full coverage spray.  Up to two applications may be applied per acre per crop season.  Allow at least 14 days between applications and 7 days between last application and harvest.

338-6467.YLD

Base Pre-Reg (6467)

U.S. LABEL

Reason to Issue: To allow for application to potatoes.


Date of Draft: 6/20/88 (Pre-Reg) (M)

EPA Reg. No. 3125-338

**RESTRICTED USE PESTICIDE**  
**Due to Acute Toxicity**

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

GUTHION® 3 Flowable  
INSECTICIDE



ACTIVE INGREDIENT:  
0,0-Dimethyl S[(4-oxo-1,2,3-benzotriazin-3(4H)-yl)methyl] phosphorodithioate ..... 30%

AMENDMENT

To Previously Registered Labeling

Revise as follows:

RECOMMENDED APPLICATIONS

CROP	PEST	Pints GUTHION 3F	REMARKS
<u>VEGETABLES</u>			
Potatoes	Colorado potato beetle	1	Apply specified dosage per acre in sufficient water for complete coverage.
	Banded cucumber beetle Leaf miners	1 to 1-1/3	A total of 3 applications may be made per crop season regardless of rate, formulation or method of application.
	European corn borer Flea beetle Leafhoppers Spittlebugs Tarnished plant bug	1-1/3 to 2	Allow at least 7 days between applications and 7 days between last application and harvest.  NOTE: Resistance of Colorado potato beetle has occurred in some areas. Consult your local Extension Service or Mobay Sales Representative for details.
	Tuberworm	1-1/2 to 2	

338-6467.YLD

Base Pre-Reg (6467)

Reason to Issue: To allow for application to potatoes.


U.S. LABEL

Date of Draft: 6/20/88 (Pre-Reg) (M)

EPA Reg. No. 3125-338

**RESTRICTED USE PESTICIDE**  
**Due to Acute Toxicity**

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.



**GUTHION® 3 Flowable**  
**INSECTICIDE**

**ACTIVE INGREDIENT:**  
 0,0-Dimethyl S[(4-oxo-1,2,3-benzotriazin-3(4H)-yl)methyl] phosphorodithioate ..... 30%

**AMENDMENT**

To Previously Registered Labeling

Revise as Follows:

**RECOMMENDED APPLICATIONS**

CROP	PEST	Pints GUTHION 3F	REMARKS
VEGETABLES Potatoes	Colorado potato beetle	1	Apply specified dosage per acre in sufficient water for complete coverage.
	Banded cucumber beetle Leaf miners	1 to 1-1/3	A total of 3 applications may be made per crop season regardless of rate, formulation or method of application.
	European corn borer Flea beetle Leafhoppers Spittlebugs Tarnished plant bug	1-1/3 to 2	Allow at least 7 days between applications and 7 days between last application and harvest.  NOTE: Resistance of Colorado potato beetle has occurred in some areas. Consult your local Extension Service or Mobay Sales Representative for details.
	Tuberworm	1-1/2 to 2	