

PM 22

3125-340

10413

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division

OCT 22 1993

Melvin K Tolliver
MILES INC
AGRICULTURE DIVISION
8400 HAWTHORN RD BOX 4913
KANSAS CITY MO 64120

Subject: Label Amendment Submission of 06/24/93
to Comply with WPS Labeling Requirements
See Reg. No. 3125-340
BAYLETON 50% WETTABLE POWDER IN WATER SOLUBLE BAGS

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is accepted subject to the comments listed below.

Based on your certification, the Agency has accepted only those changes to your labeling which are necessary to comply with PR Notices 93-7 and 93-11, which reflect the WPS labeling requirements of 40 CFR part 156, subpart K. Any other labeling changes submitted in connection with this amendment application and not directly related to compliance with PR Notice 93-7 or 93-11 have neither been reviewed nor accepted by the Agency. If you wish to make any such changes, you must submit a separate amendment application proposing them. If your product registration is currently suspended, acceptance of this labeling amendment does not affect the suspension in any way.

A copy of your proposed labeling stamped, "Accepted with comments", is attached. Make any required changes described below and send three copies of final labeling as soon as it is available to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (H-7504C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, DC 20460-0001

BEST AVAILABLE COPY

MILES INC
EPA Reg. Number: 3125-340
BAYLETON 50% WETTABLE POWDER IN WATER SOLUBLE BAGS
Application date: 06/24/93
Original Submission


Hand or courier deliveries of final labels may be made to:

Document Processing Desk (FIN-LABEL)
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Please correct the typographical errors circled on the draft before printing final labeling.

In your final labeling, the "Agricultural Use Requirements" text must be contained in a clearly separate box. This box may be set apart by a line, by another graphical device, by a different color background, or in any other way that clearly distinguishes it from surrounding text.

Sincerely,



James Taylor
Director, Office of Pesticide Registration
U.S. Environmental Protection Agency

BEST AVAILABLE COPY

3013

BAYLETON 50% WP Fungicide WSP

SPECIAL LABELING REQUIRED FOR BAGS CONTAINING PVA WATER SOLUBLE PACKETS

340-7875.YLD

U. S. LABEL

Base Pre-Reg (7875)

Reason to Issue: To revise in accordance
with the Worker Protection Standard, PR Notice 93-7.

Date of Draft: 06/15/93(Pre-Reg)(T)

BAYLETON®

50% Wettable Powder

FUNGICIDE

IN WATER SOLUBLE PACKETS

FOR CONTROL OF SPECIFIED DISEASES ON CERTAIN FIELD,
FRUIT AND VEGETABLE CROPS

ACTIVE INGREDIENT:

1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanone	50%
INERT INGREDIENTS	50%
	100%

U.S. Patent No. 3,912,752
EPA Reg. No. 3125-340
EPA Est. 3125-MO-1

DEALERS SHOULD SELL IN ORIGINAL PACKAGES ONLY
CONTAINS FOUR 8-OUNCE WATER SOLUBLE PACKETS
Total Net Weight: 2 Pounds

KEEP WATER SOLUBLE PACKETS IN THIS CONTAINER AND STORE
IN A COOL DRY PLACE BUT NOT BELOW FREEZING (32°F)

STOP - READ THE LABEL BEFORE USE

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

See Rear Panel for Statements of Practical Treatment and Other Precautionary Statements

MILES 

ACCEPTED
with COMMENTS
in EPA Letter Dated

OCT 22 1993

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
3125-340

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled, or absorbed through the skin. Prolonged or frequently repeated skin contact may cause allergic skin reactions in some individuals. Avoid contact with skin, eyes or clothing. Avoid breathing dust or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Water-proof gloves
- Shoes plus socks
- Chemical-resistant headgear for overhead exposure

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement: When handlers use closed systems, enclosed cabs, or aircraft in a manner that does not result in exposure to handlers, the following engineering controls do not apply. 40 CFR 170.240(b)(4-5). The label for this product may be found on the product label.

User Safety Recommendations:

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

STATEMENTS OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger, or, if available, by administering syrup of ipecac. If syrup of ipecac is available, administer 1 tablespoon (15 mL) of syrup of ipecac followed by 1 to 2 glasses of water. If vomiting does not occur within 20 minutes, repeat the dose once. Do not induce vomiting or give anything by mouth to an unconscious person.

If on skin: Wash with plenty of soap and water. Get medical attention.

If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, mouth to mouth. Get medical attention.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

SYMPTOMS OF POISONING: The compound does not cause any definite symptoms that would be diagnostic. Poisoning is accompanied by hyperactivity followed by sedation.

NOTE TO PHYSICIAN: No specific antidote. Treat symptomatically.

ENVIRONMENTAL HAZARDS

Do not use on other crops grown for food or forage. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water

when disposing of equipment washwaters. Apply this product only as specified on this label.

Do not make applications when weather conditions favor drift from target area.

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. The Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Chemical-resistant headgear for overhead exposure

IMPORTANT: Read these entire Directions and Conditions of Sale before using BAYLETON 50% Wettable Powder fungicide.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE THE DIRECTIONS FOR CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED FOR A RANGE OF WEATHER CONDITIONS SIMILAR TO THOSE WEATHER CONDITIONS THAT ARE ORDINARY AND CUSTOMARY IN THE GEOGRAPHIC AREA WHERE THE PRODUCT IS USED. INSUFFICIENT CONTROL OF PESTS AND/OR INJURY TO THE CROP TO WHICH THE PRODUCT IS APPLIED MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL WEATHER, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL

DIRECTIONS MAY CAUSE INJURY TO OTHER CROPS, ANIMALS, MAN, OR THE ENVIRONMENT. MILES OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL WEATHER, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF MILES AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

MIXING: The enclosed packets are water soluble. Do not allow packets to become wet prior to mixing spray solution. Do not handle with wet hands. Reseal outer bag to protect remaining packets.

Ground Application Equipment: The order of mixing is important and should be followed.

1. The spray tank should be filled 1/4 to 1/3 full of water.
2. While recirculating and with the agitator running, add the required number of PVA packets, as determined under "Recommended Applications," into the spray tank as you continue filling the tank to the desired level of water. One packet contains 8 ounces of wettable powder. If it is determined, under "Recommended Applications," that 2 pounds of BAYLETON 50% Wettable Powder should be added to the spray tank, add four 8-ounce packets. The PVA packets should be completely dissolved in 5 to 10 minutes. If an adjuvant is needed, add prior to addition of PVA packets to the spray tank.
3. The addition of other pesticides should be after the PVA packets are completely dissolved, but prior to completion of filling the tank.
4. As with any pesticide, maintain adequate agitation prior to and during spraying.

Aerial Application Equipment: It is important to pre-mix the PVA packets and water before loading into the aerial spray tank. The mixing process should be made according to the same directions for ground application equipment.

CAUTION: Do not use PVA packets in a tank-mix with products that contain Boron or release free chlorine. The resultant reaction of PVA and boron or free chlorine is a plastic which is not soluble in water or solvents such as diesel oils, kerosene, gasoline or alcohol.

Do not attempt to use the PVA packets directly in diesel oils or summer spray type oils as in ULV or LV uses. PVA packets are water soluble not oil soluble.

NOTICE TO USER: For worker protection during application, wear a hat, long-sleeve shirt, and long-legged trousers or overalls. Protective clothing should be laundered separately following application.

BEST AVAILABLE COPY

RECOMMENDED APPLICATIONS

BAYLETON 50% Wettable Powder may be applied in a minimum of 20 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment except as noted under "Remarks" for each crop. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage.

Consult your State Agricultural Experiment Station or Extension Service Specialist or Miles representative for guidance on the selection of specific dosage rates and timing of applications appropriate for your specific crop and disease. State regulations may contain additional restrictions or requirements. Check the Remarks and restrictions sections of this label for additional directions.

Do not apply this product through any type of irrigation system.

RECOMMENDED APPLICATIONS (Continued)				
CROP	DISEASE	Rate of BAYLETON 50% Wettable Powder		REMARKS (Also refer to Directions for Use)
FRUITS		<u>oz/A</u>	<u>oz/100 gal¹</u>	
Apples	Cedar-apple rust Powdery mildew	2 to 8	1/2 to 2	Make the first application at the green-tip stage and continue applications at 7- to 14-day intervals, or as needed until terminal growth ceases. Use the higher label rate under heavy disease pressure, with extended spray intervals, and/or on highly disease susceptible varieties. A maximum of 24 ounces of BAYLETON mildew 50% W.P. may be applied per acre per season. BAYLETON 50% W.P. may be applied up to 45 days prior to crop harvest. Do not graze livestock in treated orchards.
Pears	Powdery mildew	2 to 8	1/2 to 2	Make the first application at the budburst stage and continue applications at 7- to 14 day intervals, or as needed. Use the higher label rates under heavy disease pressure, with extended spray intervals, and/or on highly disease susceptible varieties. A maximum of 24 oz of BAYLETON 50% W.P. may be applied per acre per season. BAYLETON 50% W.P. may be applied up to 45 days prior to crop harvest. Do not graze livestock in treated orchards.
¹ Rates of BAYLETON 50% Wettable Powder are based on a standard of 400 gallons of dilute spray per acre, or the equivalent amount of product per acre in a concentrate spray.				

BEST AVAILABLE COPY

RECOMMENDED APPLICATIONS			
CROP	DISEASE	Rate of BAYLETON 50% Wettable Powder	REMARKS
Grapes	Powdery Mildew	2 to 6 oz/A	<p>GENERAL COMMENTS: Use the higher label rates under heavy disease pressure, with extended spray intervals, and/or on highly disease susceptible varieties.</p> <p>Fields should be observed closely for early disease symptoms when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.</p> <p>A maximum of 18 ounces of BAYLETON 50% W.P. may be applied per acre per crop season. BAYLETON 50% W.P. may be applied up to 14 days before harvest.</p> <p>POWDERY MILDEW: Make the first application pre-bloom and continue applications at 14- to 21-day intervals, or as needed.</p>
	Black rot	<p>Protective Schedule 2 to 3 oz/A</p> <p>Post-infection 3 to 4 oz/A</p>	<p>BLACK ROT</p> <p>Protective Schedule: Make the first application at 10-inch green shoot and continue applications at 7- to 14- day intervals through 5° Brix stage or until veraison (berry coloring) is complete. Under severe disease conditions and/or on highly susceptible varieties, make applications at 7-day intervals.</p> <p>Post-infection Schedule: During bloom apply within 72 hours after the beginning of an infection period. Do not make spray applications any closer than 7-day intervals. After bloom, continue applications at rates of 2 to 3 ounces of BAYLETON 50% W.P. at 7- to 14-day intervals, or as needed to maintain control.</p>
Raspberries (California Only)	Powdery mildew	4 oz	<p>Apply specified dosage per acre in not less than 20 gallons of water using ground equipment. Make first application when disease symptoms first appear and additional applications at 4- to 6- week intervals thereafter as needed. A maximum of 7 total applications may be made in one year with no more than 2 applications within the same 30-day period. Do not harvest within 1 day of application.</p>

BEST AVAILABLE COPY

RECOMMENDED APPLICATIONS (Continued)			
CROP	DISEASE	Rate of BAYLETON 50% Wettable Powder	REMARKS (Also refer to Directions for Use)
SMALL GRAINS Wheat, Barley	Powdery mildew	2 to 6 oz/A	<p>Fields should be observed closely for early disease symptoms when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.</p> <p>Apply specified dosage per acre in a minimum of 10 gallons of water per acre by ground equipment or a minimum of 5 gallons of water per acre by air when disease symptoms first appear on leaves or stems. Additional applications should be made if new disease symptoms appear.</p> <p>The total amount of BAYLETON 50% W.P. should not exceed 16 oz per acre per crop season and the last application should not be made within 21 days of harvest.</p> <p>Powdery mildew: In areas having severe powdery mildew infection apply 4 to 6 oz of BAYLETON 50% W.P. per acre.</p>
	Rusts (stripe, leaf, and stem) Speckled leaf blotch (<i>Septoria tritici</i>) Leaf and glume blotch (<i>Septoria nodorum</i>) Tan spot (<i>Pyrenophora tritici-repentis</i>) Barley scald	4 to 8 oz/A	<p>Rusts: In areas having severe rust infection apply 6 to 8 oz of BAYLETON 50% W.P. per acre.</p> <p>Leaf and glume blotch, speckled leaf blotch, and tan spot: BAYLETON 50% W.P., applied at 4 to 8 oz per acre at first sign of infection, will control light to moderate leaf and glume blotch, speckled leaf blotch, and tan spot infections. Applications for the control of glume blotch should be timed to coincide with early head emergence. Applications made prior to head emergence will not provide sufficient residual activity to control glume blotch. For control of severe infections, it is recommended that a registered fungicide with good activity against these diseases be tank mixed with BAYLETON. Refer to the Tank Mixes section of this product label.</p> <p>Barley scald: Apply 4 to 8 oz of BAYLETON 50% W.P. at the first sign of scald symptoms.</p>

BEST AVAILABLE COPY

RECOMMENDED APPLICATIONS (Continued)			
Crop	Disease	Rate of BAYLETON 50% Wettable Powder	Remarks (Also refer to Directions for Use)
GRASSES GROWN FOR SEED Bermudagrass Perennial Ryegrass Bluegrass Tall Fescue Fine Fescue Orchard Grass Bentgrass Crested Wheatgrass	Rusts <i>(Puccinia spp.)</i>	PREVENTIVE RATE 4 to 8 oz per acre CURATIVE RATE 8 to 16 oz per acre	<p>Apply the specified dosage in 20 gallons of water or more per acre with ground sprayers, or in 7 or more gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control.</p> <p>Disease control may be enhanced by the addition of a wetting agent to spray mixtures of BAYLETON 50% W.P.</p> <p>A maximum of 2 lbs. of BAYLETON 50% W.P. may be applied per acre per year.</p> <p>The last application may be made up to 5 days before harvest. Chaff and straw from treated area may be used for feed purposes; however, do not forage or cut green crop or use seed for feed purposes. Regrowth may be grazed 11 weeks after last application.</p> <p>RUSTS Preventive Schedule: For optimum disease control, apply specified dosage of BAYLETON 50% W.P. as soon as weather conditions are favorable for rust development. Apply before rust pustules are present, and continue applications at 14-to 21-day intervals depending on disease pressure.</p>
	Powdery mildew	2 to 4 oz per acre	<p>Curative Schedule: Apply the specified dosage if rust pustules are already present. A second application 14-to 21-days later may be necessary. On highly susceptible grass varieties and/or under prolonged conditions favorable for disease development use rates of 12 to 16 oz per acre and a shorter interval between applications. Rust control is especially important during the period from early head emergence through last flowering.</p> <p>Disease control may not be adequate and yield loss may occur if rust infection is allowed to reach 5 percent or greater before treating. BAYLETON 50% W.P. will prevent young, developing pustules from producing spores. However, pustules and other symptoms caused by the rust fungus prior to the application of BAYLETON 50% W.P. will remain visible.</p> <p>POWDERY MILDEW: One or two applications at the 2 to 4 oz rate applied at the first signs of disease and when new powdery mildew symptoms reappear are generally sufficient for disease control. In cases where severe mildew infections are present, use the higher rate.</p>

BEST AVAILABLE COPY

RECOMMENDED APPLICATIONS (Continued)			
Crop	Disease	Rate of BAYLETON 5% Wettable Powder	Remarks (Also refer to Directions for Use)
Pine (Seedlings) (Except California)	Pine Rust (Fusiform rust)	4 to 16 oz	Begin application prior to infection period and repeat as necessary at 2 to 3 week intervals depending upon disease pressure. Use lower rates in areas of low disease incidence and higher rates in areas of severe disease incidence. A maximum of 64 ozs of BAYLETON 50% W.P. may be applied per acre per season. A spreader-sticker is needed to help adhere spray solution to the pine trees.
Sugar Beets	Powdery mildew	BROADCAST 4 to 16 oz/A	<p>GENERAL COMMENTS: Apply the specified dosage per acre in a minimum of 10 gallons of water by ground and 5 gallons of water by air.</p> <p>Under severe disease conditions, use of the higher recommended rate may be necessary.</p> <p>Do not apply more than 16 oz. of BAYLETON 50% Wettable Powder per acre per season.</p> <p>BROADCAST APPLICATIONS: Apply specified dosage per acre as a broadcast foliar spray when disease symptoms first appear. To maintain control or if reinfection occurs, additional applications may be made at 10- to 14-day intervals.</p> <p>Broadcast applications may be made up to 15 days before harvest.</p> <p>BAND APPLICATIONS: Apply specified dosage per acre as a spray directed into the leaf whorl in a 2-inch band by turning a flat fan spray nozzle sideways to achieve a narrow band.</p> <p>Band applications may be made up to 30 days before harvest.</p>
		BAND 8 to 16 oz/A	

BEST AVAILABLE COPY

RECOMMENDED APPLICATIONS (Continued)

CROP	DISEASE	Rate Of BAYLETON 50% Wettable Powder	REMARKS (Also refer to Directions for Use)
Cucurbits (Balsam pear (bitter melon), Chinese waxgourd (Chinese preserving melon), citron melon, cucumber, gherkin, gourds, edible melons including hybrids (including cantaloupe, casaba, crenshaw, honeydew melons, honey balls, mango melon, muskmelon, Persian melon), pumpkin squash, summer squash, winter watermelon, including hybrids)	Powdery mildew (<i>Erysiphe cichoracearum</i>)*	2 to 4 oz/A	Apply specified dosage per acre in a minimum of 10 gallons of water by ground and 5 gallons of water by air as a foliar spray when disease symptoms first appear. Repeat at 14-day intervals to maintain control if new disease symptoms appear. Under severe disease conditions use the 4 oz rate. Applications may be made up to day of harvest. A maximum of 16 oz of BAYLETON 50% W.P. may be applied per acre per season. *BAYLETON 50% W.P. will not give commercially acceptable control of <i>Sphaerotheca fuliginea</i> .

TANK MIXES

The following tank mix recommendations are made for the improved control of specific diseases. Before mixing BAYLETON 50% Wettable Powder with any other fungicide, read and carefully observe the precautionary statements, directions for use, and other information appearing on the EPA registered product labels.

CROP	Rate of 50% Wettable Powder and 50% Dry Flowable Formulations of Benlate*	For Improved Control of:	BAYLETON 50% W.P. Rate
Wheat	4 oz/A	Speckled Leaf blotch Leaf and glume blotch Leaf rust Tan spot	2 oz/A
CROP	Rate of 80% Wettable Powder* Formulations of Mancozeb	For Improved Control of:	BAYLETON 50% W.P. Rate
Grapes	1 1/2 to 4 lb/A	Bunch rot Dead arm Downy mildew	2 to 6 oz/A
Sugar beets	1 1/2 to 2 lb/A	Cercospora leaf spot	4 to 16 oz/A
Wheat	2 lb/A	Helminthosporium leaf spots Speckled leaf blotch Leaf and glume blotch Tan spot Rusts	2 oz/A

* The rates given are for the following 80% wettable powder formulations of mancozeb: Dithane M-45*, Penncozeb*, Clean Crop, Mancozeb 80 WP and Mancozeb Fungicide by Riverside/Terra Corp. Only tank mixes with Dithane M-45 can be used in California.

BEST AVAILABLE COPY

ROTATIONAL CROPS

Small grains, corn, sorghum, soybeans, beans, peas and cucurbits may be planted 35 days after the last application of BAYLETON, however forage or vines from these crops may not be used for food or feed. Root crops

may be planted 120 days after the last application but tops must not be used for food or feed. All crops may be planted 12 months or later after the last application of BAYLETON without any restrictions.

RE-ENTRY STATEMENTS

~~Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.~~

~~The following may appear on supplemental labeling accompanying the product: Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product.~~

~~Oral warnings should include these statements found under the label heading "Precautionary Statements" and "Re-entry Statements" included on this label. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: WARNING, area treated with BAYLETON 50% Wettable Powder on (date of application). Do not enter without appropriate protective clothing until sprays have dried. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash with flowing water for at least 15 minutes. If swallowed, vomiting should be induced. (See Precautionary Statements on label).~~

STORAGE AND DISPOSAL

Pesticide Disposal: Do not contaminate water, food or feed by storage or disposal.

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Do not store below freezing

(32° F). Exposure to moisture or excessive handling of water soluble packets may cause breakage. Store packets in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticide as directed above. In spill or leak incidents, keep unauthorized people away. You may contact the Miles Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Miles Kansas City Emergency Response Telephone No. is 816-242-2582, or contact Chemtrec at 800-424-9300.

Benlate is a Reg. TM of E.I. DuPont de Nemours & Co. (Inc.)
Clean Crop is a Reg TM of Platte Chemical Company.
Dithane M-45 is a Reg. TM of Rohm and Haas Company.
Penncozeb is a Reg. TM of Pennwalt Corporation.

Miles Inc.
Crop Protection Products
Box 4913, Kansas City, MO 64120

BEST AVAILABLE COPY

340-7875.YLD

Base Reg (7875a)

LABELING FOR 8 OUNCE PVA PACKET

BAYLETON
50% Wettable Powder
Fungicide

ACTIVE INGREDIENT:

1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanone 50%

EPA Reg. No. 3125-340

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

(See Outer Package for Directions and Precautions)

BEST AVAILABLE COPY