


U.S. LABEL

Reason to Issue: To move Hyperodes weevil to the surface insect section of label; to add recommendation for area fire ant control. Date of Draft: 4/15/86 (Pre-Reg) (S)

EPA Reg. No. 3125-350



LAWN FOOD AND INSECTICIDE

28-3-9 Fertilizer with OFTANOL Insecticide

ACTIVE INGREDIENT:

1-Methylethyl 2-[[ethoxy[(1-methylethyl)amino] phosphinothioyl]oxy]benzoate 1.25%

AMENDMENT

To Previously Registered Labeling

Move Hyperodes weevil to the soil insect section as follows:

- Sod webworm larvae: Apply 1 lb. 14 ozs. or 3 2/3 lbs. per 1000 sq. ft., (use higher rate for severe infestations) when insect damage is first observed. Apply 1/4 to 1/2 inch of water immediately after treatment. Re-apply as necessary for pest re-infestation (see restrictions)
Chinch bugs: Apply 1 lb. 14 ozs. or 3 2/3 lbs. per 1000 sq. ft., (use higher rate for severe infestations) when insect damage is first observed. Apply 1/4 to 1/2 inch of water immediately after treatment. Re-apply as necessary for pest re-infestation (see restrictions)
Hyperodes weevil: For Hyperodes weevil control: Apply in the Spring between full bloom periods of forsythia and flowering dogwood, usually April 15 to May 15 and water-in as directed above.
Fire ants: Area Fire Ant Control: Apply 3 2/3 lb. per 1000 sq. ft. uniformly over area to be treated for residual control. Water immediately after application using 1/2 to 1 inch of water. If established fire ant mounds are present treat the mounds with a drench treatment using an EPA accepted product prior to area application. If a mound drench treatment is made with a product containing OFTANOL isofenphos, then only one area treatment may be made with this product.