

YLD29

Base Pre-Reg (5910)

U.S. LABEL

1013  
PM 21

3125-363

Reason to Issue: To clarify curative rate remarks; reduce application interval for lower preventive rate for anthracnose and poa annua decline; to modify references to control of summer patch and fusarium blight on turf.

Date of Draft: 6/5/86 (Pre-Reg) (S)

EPA Reg. No. 3125-363

® BAYLETON

1% Granular

Turf Fungicide


ACTIVE INGREDIENT:

1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H  
-1,2,4-triazol-1-yl)-2-butanone ..... 1%

AMENDMENT

To Previously Registered Labeling

Revise as Follows:



2063

RECOMMENDED APPLICATIONS  
TURF GRASS DISEASE CONTROL

DISEASE	Lbs. of BAYLETON 1% Granular Per 1,000 Sq. Ft.*		REMARKS
	<u>Preventative</u>	<u>Curative</u>	
Dollar Spot	1.5	-	<u>PREVENTIVE RATE:</u> Apply recommended rate on 30 day intervals. Protective activity of BAYLETON may extend for as long as 60 days dependent upon environmental conditions.
	0.75	-	<u>PREVENTIVE RATE:</u> Apply recommended rate on 14 day intervals. Protective activity of BAYLETON may extend for as long as 30 days dependent upon environmental conditions.
	-	3	<u>CURATIVE RATE:</u> To control existing infections, apply the curative rate. Subsequent applications should be applied <u>at</u> a preventive rate and schedule.
Copper Spot Powdery Mildew Red Thread ( <u>Laetisaria</u> <u>fuciformis</u> )  Rusts  Brown Patch ( <u>Rhizoctonia</u> <u>solani</u> ) (Suppression)	1.5	3	<u>PREVENTIVE RATE:</u> Apply at 15 to 30 day intervals. When environmental conditions favor light to moderate disease development, use a longer interval. Protective activity of BAYLETON can be greater than 30 days depending on environmental conditions.  <u>CURATIVE RATE:</u> To control existing infections, apply the curative rate. Subsequent applications should be applied <u>at</u> a preventive rate and schedule.
Anthracnose Poa Annua Decline	3	-	<u>PREVENTIVE RATE:</u> Apply at 30 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 45 days.
	1.5	-	<u>PREVENTIVE RATE:</u> Apply at <u>21</u> day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 30 days.
	-	3	<u>CURATIVE RATE:</u> To control existing infections, apply the curative rate. Subsequent applications should be applied <u>at</u> a preventive rate and schedule.

RECOMMENDED APPLICATIONS  
TURF GRASS DISEASE CONTROL

DISEASE	Lbs. of BAYLETON 1% Granular Per 1,000 Sq. Ft.*	REMARKS
Fusarium Blight ( <u>Fusarium</u> <u>culmorum</u> ) ( <u>Fusarium poae</u> ) <u>Summer Patch</u> ( <u>Phialophora</u> <u>graminicola</u> )  (Except California)	Preventive Rates Only  3	Apply first application in mid-June or 30 days prior to time this blight normally becomes evident. Applications should be made on 30 day intervals.
Fusarium Blight ( <u>Fusarium</u> <u>culmorum</u> ) ( <u>Fusarium</u> <u>poae</u> ) <u>Summer Patch</u> ( <u>Phialophora</u> <u>graminicola</u> )  (California only)	3 to 6	Apply first application in mid-June or 30 days prior to time this blight normally becomes evident. Applications should be made at the 3 lb. rate on 15 day intervals or at the 6 lb. rate on 30 day intervals.

Note\*: Apply the specified amount of BAYLETON 1% Granular with a standard granular applicator accurately calibrated to uniformly apply recommended amount of product per 1,000 sq. ft. Make application when turf is dry and following the mowing operation. Do not mow until a minimum of .25 inch of water has been applied to treated area. Do not use clippings for animal feed.

-----

All copy except that which is double underlined has been previously registered.