

363-6113

Base Pre-Reg (6113)


U.S. LABEL

Reason to Issue: To propose re-application interval for curative rate.

Date of Draft: 5/21/87 (Pre-Reg) (S)

EPA Reg. No. 3125-363

BAYLETON®
1% Granular
Turf Fungicide



ACTIVE INGREDIENT:

1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H
-1,2,4-triazol-1-yl)-2-butanone 1%

AMENDMENT

To Previously Registered Labeling

Revise as shown

RECOMMENDED APPLICATIONS
TURF GRASS DISEASE CONTROL

DISEASE	Lbs. of BAYLETON 1% G. Per 1,000 Sq. Ft.*		REMARKS
	Preventative	Curative	
Dollar Spot	1.5	-	PREVENTIVE RATE: Apply recommended rate on 30 day intervals. Protective activity of BAYLETON may extend for as long as 60 days dependent upon environmental conditions.
	0.75	-	PREVENTIVE RATE: Apply recommended rate on 14 day intervals. Protective activity of BAYLETON may extend for as long as 30 days dependent upon environmental conditions.
	-	3	CURATIVE RATE: To control existing infections, apply the curative rate. Re-application should be no closer than 14 days. When infections have been controlled, refer to the preventive rate schedule for rates and timing.

262

BAYLETON 1% Granular Turf Fungicide

RECOMMENDED APPLICATIONS (Continued)
TURF GRASS DISEASE CONTROL

DISEASE	Lbs. of BAYLETON 1% G. Per 1,000 Sq. Ft.*		REMARKS
	Preventative	Curative	
Copper Spot Powdery Mildew Red Thread (Laetisaria fuciformis)	1.5	3	PREVENTIVE RATE: Apply at 15 to 30 day intervals. When environmental conditions favor light to moderate disease development, use a longer interval. Protective activity of BAYLETON can be greater than 30 days depending on environmental conditions.
Rusts			CURATIVE RATE: To control existing infections, apply the curative rate. <u>Re-application should be no closer than 14 days. When infections have been controlled, refer to the preventive rate schedule for rates and timing.</u>
Brown Patch (Rhizoctonia solani) (Suppression)			
Anthracnose Poa Annua Decline	3	-	PREVENTIVE RATE: Apply at 30 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 45 days.
	1.5	-	PREVENTIVE RATE: Apply at 21 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 30 days.
	-	3	CURATIVE RATE: To control existing infections, apply the curative rate. <u>Re-application should be no closer than 14 days. When infections have been controlled, refer to the preventive rate schedule for rates and timing.</u>

ALL COPY EXCEPT THAT WHICH IS UNDERLINED HAS BEEN PREVIOUSLY REGISTERED.

ACCEPTED

AUG 11 1987

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 3125-363

BAYLETON^c
1% Granular
Turf Fungicide

ACTIVE INGREDIENT:

1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H
-1,2,4-triazol-1-yl)-2-butanone

AMENDMENT

To Previously Registered Labeling

Revise as shown

RECOMMENDED APPLICATIONS
TURF GRASS DISEASE CONTROL

DISEASE	Lbs. of BAYLETON 1% G. Per 1,000 Sq. Ft.*		REMARKS
	Preventative	Curative	
Dollar Spot	1.5	-	PREVENTIVE RATE: Apply recommended rate on 30 day intervals. Protective activity of BAYLETON may extend for as long as 60 days dependent upon environmental conditions.
	0.75	-	PREVENTIVE RATE: Apply recommended rate on 14 day intervals. Protective activity of BAYLETON may extend for as long as 30 days dependent upon environmental conditions.
	-	3	CURATIVE RATE: To control existing infections, apply the curative rate. <u>Re-application should be no closer than 14 days.</u> <u>When infections have been controlled, refer to the preventive rate schedule for rates and timing.</u>

RECOMMENDED APPLICATIONS (Continued)
TURF GRASS DISEASE CONTROL

DISEASE	Lbs. of BAYLETON 1% G. Per 1,000 Sq. Ft.*		REMARKS
	Preventative	Curative	
Copper Spot Powdery Mildew Red Thread (Laetisaria fuciformis)	1.5	3	PREVENTIVE RATE: Apply at 15 to 30 day intervals. When environmental conditions favor light to moderate disease development, use a longer interval. Protective activity of BAYLETON can be greater than 30 days depending on environmental conditions.
Rusts			CURATIVE RATE: To control existing infections, apply the curative rate. <u>Re-application should be no closer than 14 days. When infections have been controlled, refer to the preventive rate schedule for rates and timing.</u>
Brown Patch (Rhizoctonia solani) (Suppression)			
Anthracnose Poa Annua Decline	3	-	PREVENTIVE RATE: Apply at 30 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 45 days.
	1.5	-	PREVENTIVE RATE: Apply at 21 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 30 days.
	-	3	CURATIVE RATE: To control existing infections, apply the curative rate. <u>Re-application should be no closer than 14 days. When infections have been controlled, refer to the preventive rate schedule for rates and timing.</u>

ALL COPY EXCEPT THAT WHICH IS UNDERLINED HAS BEEN PREVIOUSLY REGISTERED.