

364-6114.

Base Pre-Reg (6114)

U.S. LABEL

Reason to Issue: To propose re-application interval for curative rate.

Date of Draft: 5/21/87 (Pre-Reg) (S)

EPA Reg. No. 3125-364

BAYLETON®

0.5% Granular

Turf Fungicide

ACCEPTED

AUG 11 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 3125-364

## ACTIVE INGREDIENT:

1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H  
-1,2,4-triazol-1-yl)-2-butanone ..... 0.5%

## AMENDMENT

To Previously Registered Labeling

Revise as shown

RECOMMENDED APPLICATIONS  
TURF GRASS DISEASE CONTROL

DISEASE	Lbs. of BAYLETON 0.5% G. Per 1,000 Sq. Ft.*		REMARKS
	Preventative	Curative	
Dollar Spot	3	-	PREVENTIVE RATE: Apply recommended rate on 30 day intervals. Protective activity of BAYLETON may extend for as long as 60 days dependent upon environmental conditions.
	1.5	-	PREVENTIVE RATE: Apply recommended rate on 14 day intervals. Protective activity of BAYLETON may extend for as long as 30 days dependent upon environmental conditions.
	-	6	CURATIVE RATE: To control existing infections, apply the curative rate. <u>Re-application should be no closer than 14 days. When infections have been controlled, refer to the preventive rate schedule for rates and timing.</u>

2062

BAYLETON 0.5% Granular Turf Fungicide

RECOMMENDED APPLICATIONS (Continued)  
TURF GRASS DISEASE CONTROL

DISEASE	Lbs. of BAYLETON 0.5% G. Per 1,000 Sq. Ft.*		REMARKS
	Preventative	Curative	
Copper Spot Powdery Mildew Red Thread (Laetisaria fuciformis)  Rusts  Brown Patch (Rhizoctonia solani) (Suppression)	3	6	<p><b>PREVENTIVE RATE:</b> Apply at 15 to 30 day intervals. When environmental conditions favor light to moderate disease development, use a longer interval. Protective activity of BAYLETON can be greater than 30 days depending on environmental conditions.</p> <p><b>CURATIVE RATE:</b> To control existing infections, apply the curative rate. <u>Re-application should be no closer than 14 days. When infections have been controlled, refer to the preventive rate schedule for rates and timing.</u></p>
Anthracnose Poa Annua Decline	6	-	<p><b>PREVENTIVE RATE:</b> Apply at 30 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 45 days.</p>
	3	-	<p><b>PREVENTIVE RATE:</b> Apply at 21 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 30 days.</p>
	-	3	<p><b>CURATIVE RATE:</b> To control existing infections, apply the curative rate. <u>Re-application should be no closer than 14 days. When infections have been controlled, refer to the preventive rate schedule for rates and timing.</u></p>

ALL COPY EXCEPT THAT WHICH IS UNDERLINED HAS BEEN PREVIOUSLY REGISTERED.