

375-6441.YLD

Base Pre-Reg (6441)


U.S. LABEL

Reason to Issue: To add Dassel to sensitive soybean varieties.

Date of Draft: 6/13/88 (Pre-Reg) (H)

EPA Reg. No. 3125-375

SALUTE 4EC
Herbicide



ACTIVE INGREDIENTS:

Trifluralin, alpha, alpha, alpha-trifluoro-2, 6-dinitro-N,N-dipropyl-p-toluidine	28%
Metribuzin, 4-amino-6-(1,1-dimethylethyl)-3-(methylthio)-1,2,4-triazin-5(4H)-one.....	14%

INERT INGREDIENTS	58%
	<u>100%</u>

AMENDMENT TO PREVIOUSLY REGISTERED LABELING

REVISE TO READ AS FOLLOWS:

SPECIAL PRECAUTIONS

Injury to soybeans may occur when SALUTE 4EC is used under the following conditions:

1. When soils have a calcareous surface area or a pH of 7.5 or higher.
2. Due to the sensitivity of certain soybeans varieties, SALUTE 4EC is not recommended for use on sensitive varieties, such as Tracy, Semmes, Altona, Dassel, Vansoy, Coker 102, Coker 156, Portage, Maple Amber, Govan, NKS-1884, Terra-Vig 505, Terra-Vig 606, Vinton 81, Asgrow 6520, Agri Pro 55, Agri Pro 71 and Narow. Consult your Mobay representative or your seed supplier for information on the tolerance of newly released soybean varieties prior to use of SALUTE 4EC.
3. When applied in conjunction with soil-applied organic phosphate pesticides.
4. Over application or boom overlapping (may result in stand loss and soil residues).
5. Uneven application or improper incorporation can decrease the level of weed control and/or increase the level of injury.
6. When applied to any soil with less than 1/2% organic matter.
7. Soil incorporation deeper than recommended.
8. When sprayers are not calibrated accurately.
9. When heavy rains occur soon after application, especially in poorly drained areas where water may stand for several days.
10. When soybeans are planted less than 1 1/2" deep.
11. Where atrazine carryover is present.
12. When poor quality soybean seed is used.

375-6455.YLD

Base Pre-Reg (6455)

U.S. LABEL

Reason to Issue: To add claims for use in tank mixes with Commence, Squadron, or Triscept herbicides on soybeans.

Date of Draft: 6/20/88 (Pre-Reg) (H)

EPA Reg. No. 3125-375

SALUTE® 4EC

Herbicide

ACCEPTED
JUL 11 1988
 Under the Federal Insecticide, Fungicide, and Plant Growth Regulator Act, as amended, for the pesticide registered under EPA Reg. No. 3125-375

ACTIVE INGREDIENTS:


- Trifluralin, alpha, alpha, alpha-trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine 28%
- Metribuzin, 4-amino-6-(1,1-dimethylethyl)-3-(methylthio)-1,2,4-triazin-5(4H)-one 14%

INERT INGREDIENTS 58%

100%

AMENDMENT TO PREVIOUSLY REGISTERED LABELING

ADD THE FOLLOWING CLAIMS FOR SOYBEANS:



SALUTE 4EC herbicide

SALUTE 4EC plus Commence Tank Mix Application

SALUTE 4EC may be applied with Commence 5.25 EC as a preplant incorporated broadcast application for the control of certain broadleaf weeds and grasses in soybeans. Consult the Commence label for specific directions for use, recommendations and restrictions not specified on this label.

Mixing:

Refer to "Mixing" under the "General Information" section in the front of the SALUTE 4EC label.

Application:

For specific application information, refer to "Application" under the "General Information" section in the front of the SALUTE 4EC label.

Weeds Controlled:

In addition to weeds controlled by SALUTE 4EC alone, SALUTE 4EC plus Commence will provide control of shattercane and greater consistency in the control of velvetleaf, especially under heavy infestation.

SALUTE 4EC plus Commence Application Broadcast Rates

Soil Texture	SALUTE 4EC (Pts/Acre)	Commence 5.25 EC (Pts/Acre)
Coarse (loamy sand or sandy loam)	1 1/2	1
Medium (loam, silt loam, silt, sandy clay, sandy clay loam)	2	1
Fine (silty clay, silty clay loam,* clay, clay loam)	2 1/4	1

* Silty clay loam soils are transitional soils and may be classified as medium textured soils in some regions of the U.S.

Special Precautions:

Do not plant wheat, oats, barley, rye or alfalfa in the fall or following spring after application as crop injury may occur.

Do not use treated vines for feed or forage.

For additional precautions, restrictions, limitations and sprayer clean-up information, refer to the appropriate sections of this label and the Commence label.

Observe all cautions and limitations on labeling of all products used in mixtures.

Do not apply where weather conditions favor drift.

SALUTE 4EC herbicide

SALUTE 4EC plus Squadron® Tank Mix Application

SALUTE 4EC may be applied with Squadron 2.33 EC as a preplant incorporated broadcast application for the control of certain broadleaf weeds and grasses in soybeans. Consult the Squadron label for specific directions on use, recommendations, restrictions and any additional weeds not specified on this label.

Mixing:

Refer to "Mixing" under the "General Information" section in the front of the SALUTE 4EC label.

Application:

For specific application information refer to "Application" under the "General Information" section in the front of the SALUTE 4EC label.

Weeds Controlled:

In addition to weeds controlled by SALUTE 4EC alone, SALUTE plus Squadron will control the following broadleaf weeds: buffalobur, cocklebur, morningglory (pitted and smallflower), shattercane, sunflower.

SALUTE 4EC plus Squadron will provide suppression (reduce the competition) of morningglory (ivyleaf and tall) and red rice.

SALUTE plus Squadron Application
(Broadcast Rates)

Soil Texture	SALUTE 4EC (pts/Acre)	Squadron 2.33 EC (pts/acre)
Coarse (loamy sand or sandy loam)	1 1/2	1 1/2
Medium (loam, silt loam, silt, sandy clay, sandy clay loam)	2	1 1/2
Fine (silty clay, silty clay loam,* clay, clay loam)	2 1/4	1 1/2

*Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S.


Special Precautions

For additional precautions, restrictions, limitations and sprayer clean-up information, refer to the appropriate sections of this and the Squadron label.

Observe all cautions and limitations on labeling of all products used in mixtures.

Do not use treated vines for feed or forage.

In coarse soils, crop injury may occur in fall planted winter wheat.



SALUTE 4EC herbicide

SALUTE 4EC plus Triscept® Tank Mix Application

SALUTE 4EC may be applied with Triscept 3EC as a preplant incorporated broadcast application for the control of certain broadleaf weeds and grasses in soybeans. Consult the Triscept label for specific directions on use, recommendations, restrictions and any additional weeds not specified on this label.

Mixing:

Refer to "Mixing" under the "General Information" section in the front of the SALUTE 4EC label.

Application:

For specific application information refer to "Application" under the "General Information" section in the front of the SALUTE 4EC label.

Weeds Controlled:

In addition to weeds controlled by SALUTE 4EC alone, SALUTE plus Triscept will control the following broadleaf weeds: buffalobur, cocklebur, morningglory (pitted and smallflower), shattercane, sunflower.

SALUTE 4EC plus Triscept will provide suppression (reduce the competition) of morningglory (ivyleaf and tall) and red rice.

SALUTE plus Triscept Application
(Broadcast Rates)

Soil Texture	SALUTE 4EC (pts/Acre)	Triscept 3EC (pts/acre)
Coarse (loamy sand or sandy loam)	1 1/2	1
Medium (loam, silt loam, silt, sandy clay, sandy clay loam)	2	1 1/4
Fine (silty clay, silty clay loam,* clay, clay loam)	2 1/4	1 1/4

*Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S.

Special Precautions

For additional precautions, restrictions, limitations and sprayer clean-up information, refer to the appropriate sections of this and the Triscept label.

Observe all cautions and limitations on labeling of all products used in mixtures.

Do not use treated vines for feed or forage.

In coarse soils, crop injury may occur in fall planted winter wheat.