

U.S. LABEL

Reason to Issue: To revise walnut use pattern and add split application.

Date of Pre-Reg Draft: 7/27/92 (C)
Supersedes Draft Dated: 10/24/91

EPA Reg. No. 3125-378

**GUTHION
35% WETTABLE POWDER
CROP INSECTICIDE**

ACTIVE INGREDIENT:
0,0-Dimethyl S-[(4-oxo-1,2,3-benzotriazin-3(4H)-yl)methyl] phosphorodithioate 35%

INERT INGREDIENTS: 65%
100%

EPA Reg. No. 3125-378

ACCEPTED
FEB 23 1993
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 3125-378

AMENDMENT

To Previously Registered Labeling

Revise as follows:

RECOMMENDED APPLICATIONS

CROP	INSECT	POUNDS GUTHION 35% WP	REMARKS
<u>NUTS</u>			
Walnuts	Codling Moth Filbertworm Navel orangeworm Walnut husk fly Red-humped caterpillar	4-1/4 to 5-11/16 2-14/16 + 2-13/16	Apply specified dosage per acre as a full coverage spray by air or ground. A total of 2 applications may be made per crop season regardless of dosage or formulation used. Allow at least 14 days between applications except in the case of split application described below. Allow at least 21 days between last application and harvest. Do not graze livestock in treated groves for 21 days after treatment. Split Application: On large trees where coverage is difficult to obtain, an aerial application followed by a ground application may be made. In order to achieve control at this rate, the two applications must be made within 10 days. Allow 21 days from the last application until harvest.

BEST AVAILABLE COPY