

GUTHION 35% WP

PM 19 3125-378
 379-7733.YL

page 1 of 1
 Base Pre-Reg. (7733)

U.S. LABEL

Date of Pre-Reg Draft: 3/8/93 (C)

Reason to Issue: To revise almond use pattern by allowing 3 applications, 28 day PHI, adding split application, and alternate row spraying directions.

ACCEPTED
 MAY 17 1993
 Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 3125-378

GUTHION 35% WETTABLE POWDER CROP INSECTICIDE

ACTIVE INGREDIENT:
O,O-Dimethyl S-[(4-oxo-1,2,3-benzotriazin-3(4H)-yl)methyl] phosphorodithioate 35%
INERT INGREDIENTS: 65%
 100%

EPA Reg. No. 3125-378

AMENDMENT To Previously Registered Labeling

Revise as Follows:

RECOMMENDED APPLICATIONS			
CROP	INSECT	POUNDS GUTHION 35% WP	REMARKS
NUTS Almonds	Peach twig borer Navel orangeworm	4-1/4 to 5-11/16	Apply specified dosage per acre as an aerial, ground concentrate or dilute application. Use at least 20 gallons per acre by aircraft and up to 400 gallons per acre by ground equipment. Applications are limited to one dormant spray and/or two post-bloom sprays per season. Alternate Row Spraying: Alternate rows may be sprayed. However, due to the presence of untreated areas, satisfactory control may not be obtained. For reduction of pest numbers to occur, all rows must be treated within 10 days of loss.
		2-14/16 + 2-13/16	Split Application: On large trees where coverage is difficult to obtain, an aerial application followed by a ground application may be made. In order to achieve control at this rate, the two applications must be made within 10 days or less.

All Applications: Allow 120 days between the dormant application and the first post-bloom spray. Allow at least 28 days between post-bloom sprays at 3 to 4 lbs per acre or the first spray of a split application. The last application may be made up to 28 days before harvest. Do not graze livestock in treated groves for 21 days after treatment.

SPECIAL LABELING REQUIRED FOR CARTONS CONTAINING WATER SOLUBLE PACKETS

379-7732.YLD

Base Pre-Reg. (7732)

U.S. LABEL

Reason to Issue: To revise almond use pattern by adding split application and alternate row spraying directions.

Date of Pre-Reg Draft: 3/8/93 (C)

ACCEPTED
 MAY 17 1993
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 3125-379

GUTHION SOLUPAK 35% WETTABLE POWDER CROP INSECTICIDE IN WATER SOLUBLE PACKETS

ACTIVE INGREDIENT:

O,O-Dimethyl S-[(4-oxo-1,2,3-benzotriazin-3(4H)-yl)methyl] phosphorodithioate 35%

INERT INGREDIENTS: 65%

100%

EPA Reg. No. 3125-379

AMENDMENT To Previously Registered Labeling

Revise as Follows:

RECOMMENDED APPLICATIONS			
CROP	INSECT	POUNDS GUTHION 35% WP	REMARKS
NUTS Almonds	Peach twig borer Navel orangeworm	4-1/4 to 5-11/16	Apply specified dosage per acre as an aerial, ground concentrate or dilute application. Use at least 20 gallons per acre by aircraft and up to 400 gallons per acre by ground equipment. Applications are limited to one dormant spray and/or two post-bloom sprays per season. Alternate Row Spraying: Alternate rows may be sprayed. However, due to the presence of untreated areas, satisfactory control may not be obtained. For reduction of pest numbers to occur, all rows must be treated within 10 days or less.
		2-14/16 to 2-13/16	Split Application: On large trees where coverage is difficult to obtain, an aerial application followed by a ground application may be made. In order to achieve control at this rate, two applications must be made within 10 days or less.
<p>All Applications: Allow 120 days between the dormant application and the first post-bloom spray. Allow at least 28 days between post-bloom sprays at 3 to 4 lbs per acre or the first spray of a split application. The last application may be made up to 28 days before harvest. Do not graze livestock in treated groves for 21 days after treatment.</p>			

GUTHION SOLUPAK 35% WP