

3121-368

John S Thornton
Manager, Registrations
Bayer Corporation
P O Box 4913
8400 Hawthorne Road
Kansas City, MO 64120 0013

MAR 22 1996

Dear Mr Thornton

Subject EPA Reg No 3125-388
Elite 45 DF Foliar Fungicide
Amendments to Label
Correspondence Dated January 22, 1996

These label amendments are acceptable in accordance with FIFRA sec. 3(c)(7)(A) provided that:

1. You submit and/or cite all data required for registration/reregistration of your product under FIFRA sec 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. You submit five copies of the final printed label for the record

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e) Your release for shipment of the product constitutes acceptance of these conditions

A stamped copy of the label is enclosed for your records

Sincerely,

Connie B Welch
Product Manager (21)
Fungicide Herbicide Branch
Registration Division

(7505C) K Scanlon ks 03/15/96 ELITEAME LTR

CONCURRENCES

SYMBOL ▶	7505C							
SURNAME ▶	KS Scanlon							
DATE ▶	Mar 15 1996							

Reason to Issue: To revise use on cherry, peach and nectarine.

Date of Draft: 01/19/96 (T)

Elite[®] 45 DF

Foliar Fungicide

For control of specified diseases on banana, cherry, peach and nectarine.

ACTIVE INGREDIENT:

Tebuconazole, α -[2-(4-chlorophenyl)ethyl]- α -(1,1-dimethylethyl)

-1H-1,2,4-triazole-1-ethanol 45%

INERT INGREDIENTS 55%

100%

EPA Reg. No. 3125-388

EPA Est. 3125-MO-1

NET CONTENTS: ___ POUNDS

Elite 45 DF

AMENDMENT TO PREVIOUSLY REGISTERED LABELING

with ~~COMMERCIAL~~
In EPA Reg. No. 3125-388

MADE IN U.S.A. 1996

Unregistered use of this Fungicide is prohibited as amended. This fungicide registered under EPA Reg. No. 3125-388

Bayer Corporation
P.O. Box 4913
8400 Hawthorne Road
Kansas City, MO 64120-0013

MAR 22 1996

Dear Mr. Thornton:

Subject: EPA Reg. No. 3125-388
Elite 45 DF Foliar Fungicide
Amendments to Label
Correspondence Dated January 22, 1996


These label amendments are acceptable in accordance with FIFRA sec. 3(c)(7)(A) provided that:

1. You submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. You submit five copies of the final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely,



Connie B. Welch
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division

(7505C):K.Scanlon:ks:03/15/96:ELITEAME.LTR

CONCURRENCES

SYMBOL ▶	7505C							
SURNAME ▶	KScanlon							
DATE ▶	Mar 15, 1996							

Elite 45 DF

388-8707.YLD

U.S. LABEL

293
Base Pre-Reg. (8707)

Reason to Issue: To revise use on cherry, peach and nectarine.

Date of Draft: 01/19/96 (T)

Elite[®] 45 DF

Foliar Fungicide

For control of specified diseases on banana, cherry, peach and nectarine.

ACTIVE INGREDIENT:

Tebuconazole, α -[2-(4-chlorophenyl)ethyl]- α -(1,1-dimethylethyl)

-1H-1,2,4-triazole-1-ethanol 45%

INERT INGREDIENTS 55%

100%

EPA Reg. No. 3125-388

EPA Est. 3125-MO-1

NET CONTENTS: ___ POUNDS

AMENDMENT TO PREVIOUSLY REGISTERED LABELING

with ~~COMMON~~ COMMENTS
In EPA ~~REGISTRATION~~
MAY 1996
United States Environmental Protection Agency
Funds ~~are available~~
as amended ~~to~~ ~~the~~ ~~labeling~~
registered under ~~EPA Reg. No.~~
3125-388

RECOMMENDED APPLICATIONS				
CROP	DISEASE	RATE OF ELITE 45 DF		REMARKS
Cherry Peach Nectarine	Brown rot (blossom blight, fruit rot)	2 oz/100 gal	4 to 8 oz/acre*	<p>Blossom blight: Begin application at white bud on cherry or pink bud on peach and nectarine. Apply again at 50% bloom and at petal fall if conditions continue to be favorable for disease development.</p> <p>Fruit rot: Make Begin application at 21, 14 and 7 days two to three weeks before harvest and continue at 7-day intervals through the day of harvest.</p>
Cherry	Leaf Spot Powdery mildew			
<p>The blossom and fruit stages must be protected for optimum control of brown rot. If ELITE 45 DF is applied during only one of these stages, another registered fungicide should be applied to the other stage to provide optimum protection. Additional cover sprays during the early postbloom period are also important for preventing quiescent fruit infections in sweet cherry and peach.</p> <p>Leaf spot: Begin application at petal fall or when first leaves unfold and continue applications at 10-7 to 14-day intervals. Applications should be made at 7-day intervals early in the growing season when terminal growth is rapid and/or under severe disease conditions. A postharvest application may be made to maintain control and reduce overwintering inoculum.</p> <p>Powdery mildew: Follow leaf spot schedule until terminal growth ceases.</p> <p>GENERAL USE DIRECTIONS*: The amount of ELITE 45 DF required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees. For smaller trees, multiply 2 oz times the number of 100 gallons of spray solution required to thoroughly wet, to the point of runoff, one acre of the trees being treated. For concentrate sprays, apply the same amount of product per acre as would be applied in a dilute spray based on tree size and foliage volume.</p> <p>ELITE 45 DF may be applied up to and including the day of harvest (0-day PHI). Do not apply more than 3 lb of product per acre per crop season.</p> <p>ELITE 45 DF is a sterol demethylation inhibitor (DMI) fungicide. It may be applied in tank mix or alternate block sprays (2 or more sprays per block) alternated (every other spray application) with a non-DMI (protectant) fungicide as a resistance management strategy. Do not mix or alternate ELITE 45 DF with other DMI fungicides.</p>				

BEST AVAILABLE COPY