

PM-25

Reg # 3125-402

194

Please read instructions on reverse before completing form. Form Approved. OMB No. 2070-0060. Approval expires 05-31-98



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number
244907

Application for Pesticide - Section I

1. Company/Product Number 3125-402	2. EPA Product Manager Robert J. Taylor	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) SENCOR Solupak 75% DF Herbicide	PM# 25	
5. Name and Address of Applicant (Include ZIP Code) Bayer Corporation, Agriculture Div. 8400 Hawthorn Rd., P.O. Box 4913 Kansas City, MO 64120-0013 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

See attached.

NOTIFICATION

JAN 2 1997

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Metal Plastic Glass Paper Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container		
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Brian A. Dehart	Title Senior Registration Scientist	Telephone No. (Include Area Code) (816) 242-2381
2. Signature 		6. Date Application Received (Stamped)
3. Title Director, Product Registrations and Regulatory Affairs		
4. Typed Name John S. Thornton		
5. Date 11/27/96		

294

402-8894.YLD

U. S. LABEL

Base Pre-Reg (8894)

Reason To Issue: To include tolerant wheat varieties which were dropped off of last EPA label submitted for SENCOR SOLUPAK.

Date of Draft: 11/15/96 (Pre-Reg) (D)
Supersedes Pre-Reg Draft Dated: 02/21/96

Sencor Solupak 75%

Sencor® Solupak 75% Dry Flowable Herbicide

ACTIVE INGREDIENT:

4-Amino-6-(1,1-dimethylethyl)-3-(methylthio)-1,2,4-triazin-5(4H)-one	75%
INERT INGREDIENTS:	25%
	100%

EPA Reg. No. 3125-402

AMENDMENT To Currently Registered Label NOTIFICATION

REVISE TO READ AS FOLLOWS:

CEREALS

(Spring and Winter Barley and Winter Wheat)

SENCOR SOLUPAK herbicide is recommended for control or suppression of certain grasses and broadleaf weeds when applied postemergence to spring and winter barley or winter wheat. SENCOR SOLUPAK alone and several tank mixture treatments are recommended for use in the following states: AR, GA, ID, IL, IN, KS, KY, LA, MS, MO, MT, NV, OH, OK, OR, TN, TX, UT, WA.

Mixing: See the 'General Information' section of this label for specific mixing procedures. When tank mixing, carefully follow the instructions on this label. Refer to the other product labels registered for use in barley and winter wheat for additional use directions, rates, weeds controlled and restrictions.

Application: SENCOR SOLUPAK may be applied by aerial or ground application equipment. Use a minimum spray volume of 2 gpa by air and 10 gpa by ground. Uniform spray coverage is necessary to obtain optimum weed control and to minimize potential for crop injury. Do not exceed rates specified on this label. Do not apply SENCOR SOLUPAK through any type of irrigation equipment. Apply SENCOR SOLUPAK when the crop is healthy and actively growing. SENCOR SOLUPAK may be applied more than once per crop season. Allow a

minimum of 21 days between applications if wheat is actively growing or allow 45 days between applications if wheat is growing in adverse conditions, has entered dormancy or is stressed due to frost damage, disease, drought or excessive moisture. Do not use on soils containing less than 0.75% organic matter. Do not apply more than a total of 10.66 ounces SENCOR SOLUPAK (8 ounces active ingredient) per acre per year. On irrigated cereals, do not apply more than 0.5 inch of water for the first irrigation, the maximum amount for each additional irrigation should not exceed 1 inch. Allow a minimum of 14 days between the first irrigation and subsequent irrigations.

Performance Factors: Weed control may not be observed for 2 to 4 weeks under normal growth conditions and for 4 to 6 weeks under very dry conditions. Moisture (at least 1/2 inch) is required within 2 to 3 weeks after application to move SENCOR into the weed root zone. Lack of adequate moisture after application may result in poor or erratic weed control. Control or suppression of listed weeds is dependent on weed size at time of application. Control or suppression may be reduced if broadleaf weeds are taller than 1 inch or grasses have more than 2 leaves.

Text being added is highlighted by redlining.

394

SENCOR SOLUPAK 75% Dry Flowable Herbicide

Tank Mixtures: SENCOR SOLUPAK may be tank mixed with Ally, Amber, Finesse, Glean FC, Harmony Extra, 2,4-D, MCPA, Igran, Banvel/Banvel SGF, Bronate or Buctril herbicides. A nonionic surfactant containing at least 80% active ingredient may be used in SENCOR SOLUPAK tank mixes with sulfonylurea herbicides (Ally, Amber, Finesse, Glean FC and Harmony Extra). Do not use a crop oil concentrate or any adjuvant containing vegetable or petroleum oils with any SENCOR SOLUPAK mix as crop injury may result. Additional pesticides may also be tank mixed with SENCOR SOLUPAK unless specifically prohibited on the mix products' label. In some instances, combinations with organophosphate insecticides may cause temporary leaf yellowing and/or crop injury, especially when widely fluctuating day/night temperatures occur near application. Always refer to the other product labels registered for use on cereals for additional directions, rates and weed species controlled. Observe all precautions and limitations on labeling of all products used in mixtures.

Feeding Restrictions: Do not graze wheat within 14 days of SENCOR SOLUPAK application or harvest grain within 21 days after last application. Do not graze or harvest barley before crop maturity. For tank mix combinations, follow the most restrictive label.

Special Precautions: Cereal Injury - Crop injury may occur if SENCOR SOLUPAK is applied:

1. When the crop is under stress such as winter kill, frost damage, disease, drought or excessive moisture, severe grazing, or when these conditions follow the application.
2. In combination with fluid fertilizer especially with the addition of surfactant,
3. Prior to the growth stage specified on this label.
4. To soils high in lime or sodium, a pH greater than 7.7, calcareous, gravelly, thinly covered or exposed subsoil areas.
5. To fields where cereal seeds have been planted less than 1 inch deep.
6. To a non-winter hardy wheat or barley variety.
7. To a sensitive wheat or barley variety as listed below.
8. To frozen soil or crop still in winter dormancy.

Cereal Rotations Following Potatoes Treated with SENCOR SOLUPAK: If planting a sensitive cereal variety (listed under the wheat and barley variety tolerance portion of this label), following potatoes treated with SENCOR SOLUPAK or metribuzin containing products, refer to the potato section of the SENCOR SOLUPAK label for special cultural practices to follow.

APPLICATION RECOMMENDATIONS

SENCOR SOLUPAK alone or in a tank mix with labeled broadleaf herbicides may be applied by aerial or ground spray equipment as a broadcast postemergence spray.

RECOMMENDED POSTEMERGENCE BROADCAST APPLICATIONS OF SENCOR SOLUPAK				
CROP GROWTH STAGE	SOIL TEXTURE	SENCOR SOLUPAK RATE (oz/A)		REMARKS
		% ORGANIC MATTER		
		0.75 TO 2.0	OVER 2.0	
2 Leaf To 2 Tiller	Coarse	1 to 2	1 to 3	Use these rates on crops with secondary roots smaller than 1 inch. For dryland winter wheat (nonirrigated), apply the highest recommended rate to achieve maximum weed suppression/control.
	Medium	1 to 3	2 to 3	
	Fine	2 to 3	2 to 4	
3 Tiller To 4 Tiller	Coarse	3 to 4	4 to 5	Do not apply within 2 weeks after grazing or breaking of winter dormancy. Apply after the crop is at or beyond the 3 tiller growth stage but before jointing. Secondary roots should be developed and larger than 1 inch long. Do not apply before 75 days after planting. For dryland winter wheat (nonirrigated), apply the highest recommended rate to achieve maximum weed suppression/control.
	Medium	4 to 5	5 to 6	
	Fine	5 to 6	5 to 6	
Over 4 Tillers	Coarse	4 to 6	5 to 8	GEORGIA ONLY: Wheat must be planted before November 15 in the Piedmont area and Northern part of the state, and before December 1 in the Coastal Plain area.
	Medium	4 to 8	5 to 8	
	Fine	5 to 8	8 to 10-2/3	

WHEAT AND BARLEY VARIETAL TOLERANCE*

Wheat and barley varieties vary in their tolerance to SENCOR SOLUPAK. Varieties below are tolerant to and are

recommended for use with SENCOR SOLUPAK:

424

SENCOR SOLUPAK 75% Dry Flowable Herbicide

Winter Wheat: ~~Abe, Arthur, AS 7846, AS 7853, Baker Seed 32, Barbie VI, Basin, Batum, Bayles, Becker, Binter, V, Buchshot DS 2368, Caldwell, Cardinal, Cashup, Centurk, Cherokee, Cheyenne, Clark, Coker 747, Coker 762, Coker 797, Coker 68-15, Coker 9134, Coker 9543, Coker 9904, Coker 9907, Daws, DB 533W, DB 562W, DB 580W, Doublecrop, Dusty, Dyna-gro 426, Dynasty, Excel, Faro, FFR 525W, Florida 302, FS 432, FS 433, FS 435, Gains, Garst 64, Georgia 100, Genie V, Hatton, Hawk, Hill 81, Howell, Hunter, Hyak, Hyslop, Katie VI, KY 16-2, Larned, Lewis 833, Lewjain, Lisa, Longhorn, Luke, Madsen, Magnum, Malcom, McDermid, McNair 1003, McNair 1813, Molly, Moro, Neely, Nelson, Newton, Norstar, Norwin, Nugaines, Oasis, Omega 78, Paha, Peck, Pike, PI 2157, PI 2180, PI 2510, PI 2545, PI 2548, PI 2550, PI 2552, PI 2555, PI 2566, PI 2571, PI 2580, PI 2684, Quantum 577, Redwin, Rocky, Saluda, Sawyer, SC 104, Siouxiand, Sprague, Southern Belle, Stacy, Stallion, Stephens, TAM W101, TAM 105, TE 877, TE 2548, TE SR204, Tiber, Tomahawk, Traveler, Tres, Tyee, Tyler, Veme, Victory, Wakefield, Wanser, Weston, Winalta, Wrangler.~~

Barley: Advance, Boyer, Clark, Compana, Hannchen, Hector, Hesk, Hudson, Lud, Luther, Kamiak, Klages, Olympic, Piroline, Steptoe and Triumph.

The following cereal varieties are sensitive to SENCOR SOLUPAK and are not recommended for use:

Winter Wheat: AT 90W, AT 91W, Arapaho, Baker Seed 33, Century, Cimarron, Coker 833, Coker 916, Coker 983, Coker 9024, Coker 9105, Coker 9323, Coker 9474, Coker 9835, Coker 9766, Coker 9877, FFR 555, Florida 304, Freedom, FS 417, FS 423, FS 425, FS 430, Gore, Hickory, Julie III, KY-49-25, Linden, Mesa, Mustang, Pacer, PI XW 522, PI 2551, PI 2163, Princeton 733, PSR W71, PSR 226, PSR 278, Rosen, Savannah, Sierra, TAM 107, TR 101, TR 1011, TR 8822, Triumph 64, Vona, Wings, Winridge, Yamhill.

Spring/Durum Wheat: Avoid use on Spring wheat and Durum wheat varieties.

Barley: Glenn, Morex, Moravian 3, Larker, Summit, Bracken, Anheuser Busch B2601 and varieties with Morex parentage.

Varieties Not Listed: To avoid possible crop injury on any variety not mentioned in this label, contact a Bayer representative or herbicide expert for a variety recommendation prior to treatment or treat a small strip of the unlisted variety with the recommended SENCOR SOLUPAK rate to ascertain crop tolerance before treating an entire field.

* Abbreviated names of vendors: AS (Agseco), AT (Agratech), DB (Diener Bros.), FS (Growmark FS), PI (Pioneer), PSR (Hybritech), SC (J.M. Schultz), TE (Terra), and TR (Terral).

WEEDS CONTROLLED			
Used at recommended rates, SENCOR SOLUPAK will control many annual broadleaf weeds. Control is best when applied to young, actively growing weeds. Weeds controlled by SENCOR SOLUPAK include:			
Bittercress	Evening Primrose, Cutleaf	Knotweed, Prostrate	Pigweed, spp.
Catchfly, Conical (Sand)	Falseflax, Smallseed	Lambsquarter, Common	Pineappleweed
Catchweed (Madwort)	Fiddleneck, Tarweed	Lettuce, Miners	Polemonium, Annual (Jacob's Ladder)
Chickweed, Common	Filaree, Redstem	Mustard, Blue	Radish, Wild
Chickweed, Mouseear	Geranium, Carolina	Mustard, Wild	Shepardspurse
Comcockle	Gromwell, spp.	Pennycress, Field	Speedwell, Ivyleaf
Dogfennel (Mayweed)	Henbit	Pepperweed, Virginia	Turnip, Wild

WEEDS SUPPRESSED					
SENCOR SOLUPAK control of the following weeds varies from poor to excellent depending on time of application, stage of growth at application, temperatures and soil moisture conditions following treatment. For maximum effect on these weeds, apply the highest recommended rate at the earliest growth stage timing for each particular soil type and organic matter. Suppression is a reduction in weed size and growth as compared to a non-treated area in the same field.					
BROADLEAVES		GRASSES			
Buckwheat, Wild*	Mustard, Tumble (Jim Hill)*	Barley, Hare (Wild)	Bluegrass, Bulbous	Cheat*	Whitlowgrass, Spring
Buttercup, spp.	Tanymustard	Barley, Little	Brome, Downy*	Foxtail, spp*	(Vernal) , ,
Cowcockle	Thistle, Russian	Blackgrass	Brome, Japanese*	Oat, Wild*	Windgrass
Kochia*	Vetch, Winter	Bluegrass, Annual	Brome, Riggut*	Rescuegrass*	, , , ,
Lettuce, Pricky					, , , ,

* Use the highest recommended SENCOR SOLUPAK rate for maximum weed suppression.

BEST AVAILABLE COPY

Bayer Corporation
Crop Protection Products
Box 4913
Kansas City, MO 64120-0013

