

PM

13

3125-403

Page 187

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

3108486086
1615447984
17/13

2 - MAY 1995

Ms. Julie Spagnoli
Bayer Corp.
PO Box 4913
8400 Hawthorn Rd.
Kansas City, MO 64120-0013

Dear Ms. Spagnoli:

Subject: Labeling Amendments; Add Turf Use Around Commercial Structures
Tempo 2 Lawn and Ornamental Insecticide
EPA Registration No. 3125-352
Tempo 20 WP Lawn and Ornamental Insecticide
EPA Registration No. 3125-403
Tempo 20 WP Lawn and Ornamental Insecticide in Water Soluble Packets
EPA Registration No. 3125-395
Your Applications Dated April 10, 1995

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

- Under "Applications to Ornamentals" revise the statement, "...to the point of run-off" to read as follows: "...to the point of run-off but do not allow run-off to occur."
- Under Environmental Hazards delete "For terrestrial uses." This phrase incorrectly implies that your product has aquatic uses.

Submit three copies of your final printed labeling before you release the product for shipment. A stamped copy of the labeling is enclosed for your records.

Sincerely,

George T. LaRocca
Product Manager 13

1087

CONCURRENTS
Insecticide & Rodenticide Branch
Registration Division (H7505C)

SYMBOL							
SURNAME	Helmut						
DATE	5-2-95						

TEMPO 20 WP Lawn and Ornamental Insecticide

403-8488.YLD

2 9 7
Base Pre-Reg (8488)

U.S. LABEL

Reason to Issue: To propose use on landscape and recreational turf. 04/04/95 Draft: To exclude golf course use. To change company name.

Date of Draft: 04/04/95(Pre-Reg)(S)
Supersedes Pre-Reg Draft Dated: 07/15/93

TEMPO 20 WP

Lawn and Ornamental Insecticide

**FOR COMMERCIAL USE ONLY
FOR BROAD-SPECTRUM CONTROL OF INSECT PESTS INFESTING
RESIDENTIAL AND COMMERCIAL LAWNS, TREES AND ORNAMENTALS**

ACTIVE INGREDIENT:

Cyfluthrin, cyano(4-fluoro-3-phenoxyphenyl)methyl 3-(2,2-dichloroethenyl)- 2,2-dimethylcyclopropanecarboxylate	20%
INERT INGREDIENTS:	80%
	100%

EPA File Symbol No. 3125-403
EPA Est. No. 3125-MO-1

TOTAL NET CONTENTS: 600 GRAMS

STOP - READ THE LABEL BEFORE USE

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

See Rear Panel for Statements of Practical Treatment and Other Precautionary Statements


PRECAUCION AL USUARIO: Si usted no puede leer o entender inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)



ACCEPTED
with COMMENTS
in EIA Letter Dated
2 - MAY 1995

Under the Federal Insecticide, Fungicide, and Rodenticide Act, this product is registered under EPA Reg. No. 3125-403



2

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Harmful if swallowed, inhaled or absorbed through the skin. Do not get in eyes, on skin, or on clothing. Avoid breathing dust or spray mist.

Do not contaminate feed or food. Keep out of reach of children.

Do not allow children or pets to enter treated areas until surfaces are dry. Wash thoroughly with soap and warm water after handling. Remove contaminated clothing and wash before reuse.

SYMPTOMS OF POISONING: In case of poisoning, call physician or Poison Control Center immediately. Have patient lie down and keep quiet.

STATEMENTS OF PRACTICAL TREATMENT

If in eyes: Hold eyelids open and flush with plenty of water. Call a physician if irritation persists. **If swallowed,** call a physician or Poison Control Center. Administer water freely and induce vomiting by giving one dose (1/2 oz or 15 ml) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is

clear. Never give anything by mouth to an unconscious person. Avoid alcohol. **If on skin:** Wash thoroughly with soap and water. Get medical attention if irritation occurs. **If inhaled,** remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.

TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply TEMPO 20 WP Insecticide or allow it to drift onto crops or weeds on which bees are actively foraging. Additional information may be obtained by consulting your Cooperative Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read these entire Directions and Conditions of Sale before using TEMPO 20 WP Insecticide.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE THE DIRECTIONS FOR CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED FOR A RANGE OF WEATHER CONDITIONS SIMILAR TO THOSE WEATHER CONDITIONS THAT ARE ORDINARY AND CUSTOMARY IN THE GEOGRAPHIC AREA WHERE THE PRODUCT IS USED. INSUFFICIENT CONTROL OF PESTS AND/OR

INJURY TO THE CROP TO WHICH THE PRODUCT IS APPLIED MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL WEATHER, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO OTHER CROPS, ANIMALS, MAN, OR THE ENVIRONMENT. BAYER OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL WEATHER, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF BAYER AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

Do not formulate this product into other end-use products.

4 9 7

TEMPO 20 WP Insecticide will not stain or cause damage to any painted or varnished household surface, plastic, fabric or other surface where water applied alone causes no damage.

Add the appropriate amount of product when filling sprayer tank with water, shake or agitate as necessary to mix. Diluted spray mixture can be stored overnight and applied the following day; however, mixture should be agitated prior to application to prevent uneven distribution of product.

One level spoonful (as provided) or one level tablespoonful equals 6 grams. Do not re-use measuring utensils for feed, food, or drinking water purposes.

When used at recommended rates and as directed under **RECOMMENDED APPLICATIONS**, TEMPO 20 WP Insecticide will control the designated pests on trees, shrubs, foliage plants and flowers in outdoor landscaped areas such as parks, recreational areas, athletic fields, institutional grounds, etc., and interior plantscapes (such as in hotels, shopping malls, office buildings, etc.) where these plants are grown. TEMPO 20 WP Insecticide also controls designated pests on landscape and recreational turf, including home lawns, business and office complexes, shopping complexes, multi-family residential complexes, airports, cemeteries, parks, playgrounds and athletic fields, and other landscape turfgrass.

NOTE: Not for use in commercial greenhouses, nurseries, sodfarms or grass grown for seed. Not for use on golf course turf.

APPLICATION TO ORNAMENTALS: TEMPO 20 WP Insecticide mixes readily with water and may be used in all types of spray equipment. Mix product with the required amount of water and apply as a dilute spray application to the point of run-off. When spraying hard-to-

wet foliage such as holly, pine or ivy, the addition of a spreader/sticker may enhance both knock-down and residual activity. If concentrate or mist type spray equipment is used, an equivalent amount of product should be used on the area sprayed, as would be used in a dilute application. Phytotoxicity testing has been carried out on a wide range of ornamental plants under various environmental conditions and no phytotoxicity has been observed. It is advised to pre-spray a selection of ornamentals if local use experience is unavailable. Time applications to flowering plants during periods when pollinating insects are not present, such as early morning or late evening.

APPLICATIONS ON TURF: TEMPO 20 WP Insecticide mixes readily with water and/or various fertilizer solutions and can be used in appropriate spray equipment to treat listed turfgrass. Applications should be made with accurately calibrated equipment and applied uniformly, using a coarse, large droplet, spray applications procedure. Do not allow drift onto ponds, streams, or lakes. For optimum control of armyworms, cutworms and sod webworms, delay irrigation and mowing for at least 24 hours following the application. Keep children and pets off treated areas until grass have dried following application.

COMPATIBILITY: TEMPO 20 WP Insecticide has been found to be compatible with all commonly used fungicides, miticides, liquid fertilizers and other insecticides. Check physical compatibility using the correct proportion of products in a small jar if local experience is unavailable.

Do not apply this product through any type of irrigation system.

4

5 9 7

RECOMMENDED APPLICATIONS			
CROP	PEST	DOSAGE TEMPO 20 WP	REMARKS
Landscape Ornamentals (Including Trees, Shrubs, Evergreens, Flowers, Foliage Plants)	Armyworms Azalea caterpillars Bagworms Bristly rose slugs California oakworms Cankerworms Crickets* Cutworms Elm spanworms Flies* Fungus gnats Gypsy moth larvae Lace bugs Leaf-feeding caterpillars Midges* Mosquitoes* Oleander moth larvae Pillbugs Pine shoot moths Pine tip moths Redhumped caterpillars Sawfly larvae Sowbugs Spiders* Striped oakworms Tent caterpillars Tussock moth larvae Walnut caterpillars Webworms Yellownecked caterpillars	36 grams (1.3 oz) Per 100 Gal.	Applications should be started prior to establishment of high pest populations and reapplication made as necessary. Good spray coverage is necessary to provide the most effective level of control. Addition of a spreader/sticker at recommended rates may enhance control of insects on certain species of ornamentals having hard-to-wet surfaces. *The use of TEMPO 20 WP Insecticide for the control of crickets, spiders, midges, flies and mosquitoes is limited to ornamental areas and areas adjacent to buildings. Apply as a general spray, concentrating on damp areas beneath shrubbery, around foundations and in areas of tall grass or weeds. Do not apply to any food crop. Remove any animal feeding dishes prior to treatment and do not allow spray or drift to contact fish-bearing water.

RECOMMENDED APPLICATIONS			
CROP	PEST	DOSAGE TEMPO 20 WP	REMARKS
Landscape Ornamentals (Continued) (Including Trees, Shrubs, Evergreens, Flowers, Foliage Plants)	Ants* Aphids Boxelder bugs Budworms Casebearers Clover mites Cockroaches** Elm leaf beetles Flea beetles Grasshoppers Japanese beetles (adult) June beetles (adult) Leafhoppers Leafrollers Leaf skeletonizers Mealybugs Orchid weevil Pear psylla Peppertree psyllid Plant bugs Scale insects (crawler stages) Spittlebugs Striped beetles Thrips Ticks (including deer ticks) Tussock moth larvae Wasps* Whiteflies	54 grams (1.9 oz) Per 100 Gal.	Applications should be started prior to establishment of high pest populations and reapplication made as necessary. Good spray coverage is necessary to provide the most effective level of control. Addition of a spreader/sticker at recommended rates may enhance control of insects on certain species of ornamentals having hard-to-wet foliage. *The use of TEMPO 2 Insecticide for the control of ants and wasps is limited to ornamental areas and areas adjacent to buildings. Apply as a general spray, concentrating on damp areas beneath shrubbery, around foundations and in areas of tall grass or weeds. Do not apply to any food crop. Remove any animal feeding dishes prior to treatment and do not allow spray or drift to contact fish-bearing water. **Outdoor cockroach control (including Asian cockroaches) is obtained by spraying hiding or resting places such as under shrubbery, in ground covers and mulching materials, in hollow trees, etc.
CROP	PEST	DOSAGE TEMPO 20 WP	REMARKS
LANDSCAPE AND RECREATIONAL TURFGRASS (including home lawns, business and office complexes, shopping complexes, multi-family residential complexes, airports, cemeteries, parks, playgrounds and athletic fields, and other landscape turfgrass	Ants Armyworms Crickets Cutworms Earwigs Grasshoppers Japanese beetles (adult) Sod webworms Ticks (including deer ticks)	5 grams per 1000 sq ft or 7.7 oz/A	Apply when pests first appear. Reapply when necessary based on pest reinfestations, not exceeding 6 applications per year. Delay watering or mowing for 24 hours for optimum control of armyworms, cutworms, and sod webworms. Keep children and pets off treated areas until lawn has dried following the application. * Aids on control only.
	Bluegrass billbug (adult) Chinch bugs Chiggers Fire ants* Fleas Hyperodes weevil (adult) Mole crickets Scorpions	7 grams per 1,000 sq ft or 10.75 oz/A	

RESTRICTIONS FOR TURFGRASS APPLICATION

Do not apply this product by aerial application or by any type of chemigation system. Do not apply when the average windspeed is greater than 15 mph.

Apply product using spray nozzles which produce a coarse droplet size, such as flood jet nozzles. Do not apply when turfgrass areas are water-logged or the soil is saturated with water (i.e., will not accept irrigation). Use this product only as directed on label.

STORAGE AND DISPOSAL

Not for Storage In or Around the House

Pesticide Disposal: Do not contaminate water, food, or feed by storage or disposal.

Pesticide wastes are hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

Container Disposal: Do not use container in connection with food, feed, or drinking water. Completely empty container into application equipment. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Storage: Store in cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original containers and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticide as directed above. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer Kansas City Emergency Response Team Telephone No. is 816-242-2582, or contact Chemtrec at 800-424-9300.

Sold by:
Bayer Corporation
Specialty Products
Box 4913, Kansas City, MO 64120-0013