

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS 767) WASHINGTON, DC 20460	EPA REGISTRATION NO. 3125-413	DATE OF ISSUANCE 12/1/76
NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> REGISTRATION <input type="checkbox"/> REREГИSTRATION <i>(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)</i>	TERM OF ISSUANCE Conditional	
NAME OF PESTICIDE PRODUCT Bayleton 25% Wettable Powder System Fungicide in Water Soluble Packets		
NAME AND ADDRESS OF REGISTRANT (Include ZIP code)		
<div style="border: 1px solid black; padding: 10px; min-height: 100px;"> <div style="position: relative; height: 100px;"> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 100%; border: 1px solid black;"></div> </div> </div>		
NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.		
On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.		
A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.		
Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.		
This product is conditionally registered in accordance with Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) section 3(c)(7)(A) provided that you:		
<ol style="list-style-type: none"> 1. Submit/cite all data required for registration/-reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable response required for reregistration of your product under FIFRA section 4. 2. Make the labeling changes listed below before you release the product for shipment: <ol style="list-style-type: none"> a. Add the phrase "EPA Registration No. 3125-413." b. In the environmental hazards section update the statements to include the following statements: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. c. Move "Do not use on crop grown for forage or food" to the directions for use section. 		
<input type="checkbox"/> ATTACHMENT IS APPLICABLE		
SIGNATURE OF APPROVING OFFICIAL 		DATE

3. Submit one (1) copy of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

4. Submit within 15 months from the date of this Notice of Registration a Storage Stability Study conducted in accordance with Guideline Reference Number 63-17.

5. Submit within 15 months from the date of this Notice of Registration sample analyses for five representative batches including the raw analytical data to support the nominal concentrations as required in PR Notice 91-2.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Cynthia Giles-Parker
Product Manager (22)
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

LABELING FOR PRODUCT IN WATER SOLUBLE PACKETS

318-7425.YLD

Base Pre-Reg. (7425)

U.S. LABEL(S)

Reason to Issue: To propose new
product in water soluble packets.
12/11/91 Draft: To revise according
to internal review comments.

Date of Draft: 12/11/91 (Pre-Reg)(S)
Supersedes Pre-Reg Draft Dated 7/11/91 and
9/11/91

EPA File Symbol No. 3125-XXX

BAYLETON®
25% Wettable Powder
SYSTEMIC FUNGICIDE
IN WATER SOLUBLE PACKETS

FOR CONTROL OF CERTAIN DISEASES ON FLOWERS, FOLIAGE PLANTS, SHRUBS,
TURF AND SHADE TREES

ACTIVE INGREDIENT:

~~delete~~ ① 1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H
1,2,4-triazol-1-yl)-2-butanone 25% ✓

INERT INGREDIENTS 75% ✓
100% ✓

KEEP WATER SOLUBLE PACKETS IN THIS
CONTAINER AND STORE IN A COOL PLACE
BUT NOT BELOW FREEZING (32°F)

U.S. Patent No. 3,912,752

EPA File Symbol No. 3125-XXX

EPA Est. 3125-MO-1

STOP - READ THE LABEL BEFORE USE

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

See Rear Panel for Statements of Practical Treatment
and Other Precautionary Statements

SELL IN ORIGINAL PACKAGES ONLY

CONTAINS FOUR 11 OUNCE WATER SOLUBLE PACKETS

TOTAL NET CONTENTS 2-3/4 POUNDS

ACCEPTED
with COMMENTS
In EPA Letter Dated

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the purpose of
being registered under EPA Reg. 40.100.

3125-413

MOBAY CORPORATION
A Bayer USA INC COMPANY
AGRICULTURAL CHEMICALS DIVISION
SPECIALTY PRODUCTS GROUP
Box 4913, Kansas City, MO 64120

Note: Underlined letters in chemical nomenclature should be italicized when printed,
not underlined.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Exposure to wettable powder can cause eye irritation. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENTS OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center. Vomiting should be induced. Administer water freely and induce vomiting by giving one dose (1/2 oz or 15 ml) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person. Professional medical assistance should be secured immediately.

If in eyes: Flush with plenty of water. Call a physician if irritation persists.

If on skin: Wash with plenty of soap and water. Get medical attention.

SYMPTOMS OF POISONING: The compound does not cause any definite symptoms that would be diagnostic. Poisoning is accompanied by hyperactivity followed by sedation.

NOTE TO PHYSICIAN: No specific antidote. Treat symptomatically.

ENVIRONMENTAL HAZARDS

Do not use on crops grown for food or forage. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not make applications when weather conditions favor drift from target area.

AVISO:

PRECAUCION AL USUARIO: Si usted no puede leer o entender ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read these entire Directions and Conditions of Sale before using BAYLETON 25% Wettable Powder systemic fungicide.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE APPROPRIATE FOR THE CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED UNDER DIFFERENT ENVIRONMENTAL CONDITIONS BOTH INDOORS AND OUTDOORS UNDER CONDITIONS SIMILAR TO THOSE THAT ARE ORDINARY AND CUSTOMARY WHERE THE PRODUCT IS TO BE USED. INSUFFICIENT CONTROL OF PESTS OR PLANT INJURY MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL CONDITIONS, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO ANIMALS, MAN, AND DAMAGE TO THE ENVIRONMENT. MOBAY OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL ENVIRONMENTAL CONDITIONS, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF MOBAY AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

APPLICATION:

BAYLETON fungicide is absorbed rapidly and works systemically from within the plant. Good coverage and wetting of the foliage is necessary. Rainfall or sprinkler irrigation, within 1/2 hour after application does not decrease effectiveness. Control may be less effective on plants suffering from drought stress. Therefore, in order to achieve maximum control, plants should be maintained in a vigorously growing state through good cultural practices.

In all cases apply when plants are fully established and actively growing. Applications should be made at recommended intervals to maintain disease control.

MIXING: The enclosed packets containing BAYLETON 25% Wettable Powder are water soluble. Do not allow packets to become wet prior to adding to the spray tank. Do not handle with wet hands. Reseal outer carton to protect remaining packets.

To prepare the spray mixture, drop the required number of unopened packets, as determined under "Recommended Applications", into the spray tank while filling the water to the desired level. Operate the agitator while mixing. Depending on the water temperature and the degree of agitation, the packets should be completely dissolved within a few minutes from the time they were added to the water.

BEST AVAILABLE COPY

USE IN CHEMIGATION SYSTEMS ON TURF ONLY:

Apply BAYLETON 25% Wettable Powder only through solid set irrigation systems. Do not apply this product through any other type of irrigation system.

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back towards the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where the pesticide distribution is adversely affected.

Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the areas intended for treatment.

Turf injury or lack of effectiveness can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Pre-mix the required amount of BAYLETON 25% Wettable Powder, as determined under "Recommended Applications", in sufficient water to uniformly inject the entire mixture during the last 5 minutes of the irrigation cycle using a positive pressure pumping system. Continuous agitation of the mixture in the holding tank is required to maintain suspension of the product. The injection must occur during the last 5 minutes of the irrigation cycle.

RECOMMENDED APPLICATIONS
TURF GRASS DISEASE CONTROL

DISEASE	Ozs of BAYLETON 25% Wettable Powder Per 1,000 Sq Ft*	REMARKS
Dollar Spot (<u>Sclerotinia</u> <u>homoeocarpa</u>)	<u>Preventive</u> <u>Curative</u>	<u>PREVENTIVE RATE:</u> Apply recommended rate at 30-day intervals. Protective activity of BAYLETON may extend for as long as 60 days, depending upon environmental conditions.
	1 -	<u>CURATIVE RATE:</u> To control existing infections, apply the curative rate. Subsequent applications should be applied on a preventive schedule and rate.
Copper Spot (<u>Gloeocercospora</u> <u>sorghii</u>) Powdery Mildew (<u>Erysiphe graminis</u>) Corticium Red Thread (<u>Laetisaria</u> <u>fuciformis</u>) Rusts (<u>Puccinia spp.</u>) Brown Patch/ Rhizoctonia Blight (<u>Rhizoctonia solani</u>) (Suppression)	1 2	<u>PREVENTIVE RATE:</u> Apply at 15- to 30-day intervals. When environmental conditions favor light to moderate disease development, use a longer interval. Protective activity of BAYLETON can be greater than 30 days depending on environmental conditions.
	2 -	<u>CURATIVE RATE:</u> To control existing infections, apply the curative rate. Subsequent applications should be applied on a preventive schedule and rate.
Anthracnose (<u>Colletotrichum</u> <u>graminicola</u>)	2 -	<u>PREVENTIVE RATE:</u> Apply at 30-day intervals and repeat as necessary for seasonal control. Depending upon environmental conditions, residual control may be extended to 45 days.
	- 2	<u>CURATIVE RATE:</u> To control existing infections, apply the curative rate. Subsequent applications should be applied on a preventive schedule and rate.

BAYLETON 25% Wettable Powder Fungicide in Water Soluble Packets

RECOMMENDED APPLICATIONS (Continued) TURF GRASS DISEASE CONTROL

DISEASE	Oz of BAYLETON 25% Wettable Powder Per 1,000 Sq Ft*		REMARKS
	<u>Preventive</u>	<u>Curative</u>	
Southern Blight (<u>Sclerotium rolfsii</u>)	1 to 4	4	<u>PREVENTIVE</u> : Begin applications prior to the appearance of disease symptoms. Depending on anticipated disease severity, apply 2 to 4 oz rates at 14-day intervals for the initial 2 to 3 treatments. Apply subsequent treatments of 1 to 2 oz at 14- to 28-day intervals. <u>CURATIVE</u> : To control existing infections, apply 4 oz at 14-day intervals for the initial 2 to 3 treatments followed by 1 to 2 oz at 14- to 28-day intervals.
Stripe Smut (<u>Ustilago striiformis</u>)	<u>Preventive Rates Only</u> 2		Make the first application in the spring just before the turf breaks dormancy, followed by a second application just prior to the summer heat stress period and a third application when the cool nighttime temperatures of the late summer or early fall return.
Fusarium Blight (<u>Fusarium culmorum</u>) (<u>Fusarium poae</u>) Summer Patch (<u>Magnaporthe poae</u>)	2 to 4		Apply first application in the Spring, 30 to 60 days before initial symptoms normally appear. Repeat applications at 30-day intervals as needed.
Gray Snow Mold/ Typhula Blight (<u>Typhula incarnata</u>) (<u>Except California</u>)	4		Apply in fall 30 days prior to turf dormancy. If turf breaks dormancy during winter months, a second application should be made. Do not apply over snow cover, or when turf is dormant.
Pink Snow Mold/ Fusarium Patch (<u>Microdochium nivalis</u>) (<u>Except California</u>)	2 to 4		Apply before conditions favorable for infection occur. Re-application should be made as needed at a 60- to 90-day interval. Do not apply over snowcover, or when turf is dormant. Use higher rate in areas with a history of severe disease damage.

Note: *Apply the specified amount of BAYLETON 25% Wettable Powder using 2 to 4 gallons of spray per 1,000 sq ft. For equivalent rates per acre, see Conversion Table below. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. Do not use clippings for animal feed.

Conversion Table for
BAYLETON 25% Wettable Powder

<u>Oz/1,000 ft²</u>	<u>Pounds per Acre</u>	<u>Number of 11 oz PVA Packets per Acre</u>
1.0	2.75	4
2.0	5.5	8
4.0	11.0	16

BAYLETON 25% Wettable Powder Fungicide

ORNAMENTAL PLANT DISEASE CONTROL		DISEASES
PLANTS (Disease controlled reference)		
<u>Flowering and Foliage Plants (Outdoor)</u> (Except California)	<u>Ornamental Shrubs and Trees</u> (Except California)	(1) Flower Blight a. <u>Ovulinia</u> spp. [A] b. <u>Sclerotinia</u> spp. [A] c. <u>Collectotrichum</u> [A]
Ageratum (2b,3,4)	Amelanchier (3)	
Aster (4)	Azalea (1a,2e,3)	
Begonia (3)	Barberry (3,4)	
Canna (4)	Buckthorn (4)	
Carnation (3,4)	Camellia (suppression of 1b)	
Chrysanthemum (3,4)	Crabapple (flowering) (3,4)	(2) Leaf Blight/Spots a. <u>Cephalosporium</u> spp. [C] b. <u>Cercospora</u> spp. c. <u>Didymella</u> spp. [B] d. <u>Entomosporium</u> spp. [C] e. <u>Exobasidium</u> spp. [E]
Dahlia (3)	Crape myrtle (3)	
Delphinium (3)	Dogwood (3)	
Dianthus (4)	Euonymus (3)	
Four O'Clock (4)	Gardenia (3)	
Geranium (3,4)	Hawthorn (3,4)	
Hollyhock (3,4)	Hemlock (4d,e)	
Iris (2c)	Holly (3)	
Marigold (2b,4)	Juniper (4)	
Nephtytis (2a)	Leucothoe (2b)	
Pansy (3,4)	Lilac (3)	
Petunia (3,4)	Mock-Orange (3,4)	(3) Powdery Mildew <u>Erysiphe</u> spp. <u>Microsphaera</u> spp. <u>Oidium</u> spp. <u>Podosphaera</u> spp. <u>Phyllactinia</u> spp. <u>Sphaerotheca</u> spp. <u>Uncinula</u> spp.
Phlox (2b,3,4)	Mountain Laurel (1a,2b,3)	
Rose (3)	Ninebark (3)	
Salvia (3,4)	Paulownia (3)	
Sedum (3)	(Empress tree)	
Snapdragon (3,4)	Pear (Flowering) (3)	
Sunflowers (3,4)	Photinia (2d,3,4)	
(ornamental only)	Potentilla (4)	
Sweet peas (3)	(Cinquefoil)	
Zinnia (2b,3)	Privet (2b,3)	
	Pyracantha (3)	
	Rhododendron (1a,2b,3)	(4) Rusts a. <u>Coleosporium</u> spp. b. <u>Cronartium</u> spp. [B] (Fusiform) c. <u>Gymnosporangium</u> spp. d. <u>Melampsora</u> spp. [A] e. <u>Melampsora</u> farlowii [A] f. <u>Melampsoridium</u> spp. g. <u>Peridermium</u> spp. [B] (Gall) h. <u>Phragmidium</u> andersonii i. <u>Puccinia</u> spp. j. <u>Uromyces</u> spp. k. <u>Uredinopsis</u> mirabilis [A]
<u>Shade Trees</u> (Except California)		
Ash (3)	Spirea (3)	
Aspen (3,4)	Viburnum (3,4)	
Birch (3,4)	Vitex (2b)	
Buckeye (3)	(Chaste-tree)	
Chestnut (3)		
Cottonwood (3,4)	<u>Flowering and Foliage Plants</u> (Greenhouse [D]) (Except California)	
Elm (3)	African Violet (3)	
Fir (4)	Azalea (1a,2e,3)	
Locust (3)	Calendula (3,4)	
Maple (3)	Carnation (3,4)	
Oak (3)	Chrysanthemum (3,4)	
Pine (4,5)	Cineraria (3)	
Poplar (3,4)	Crassula (3)	
Russian Olive (2b,4)	Daisy (3,4)	
Sycamore (3)	Fern, Boston (4)	
Walnut (3)	<u>Desmella</u> spp.	
Willow (3,4)	Geranium (3,4)	
	Gerbera (3)	
<u>Hawaii Only</u>	Grape Leaf Ivy (3)	
Dendrobium (1c)	Kalanchoe (3)	(5) Tip Blight <u>Sirococcus</u> strobilinus [B]
	Rose (3)	
	Snapdragon (3,4)	
<u>California Only</u>		
<u>Recommendations</u>	<u>California Only</u> <u>Recommendations</u>	
Azalea (1a,3)	Greenhouse [D]	
Begonia (3)	African Violet (3)	
Crape myrtle (3)	Carnation (3,4)	
Euonymus (3)	Chrysanthemum (3,4)	
Geranium (3,4)	Geranium (3,4)	
Hollyhock (3,4)	Grape Leaf Ivy (3)	
Iris (2c)	Rose (3)	
Nephtytis (2c)		
Oak (3)		
Pine trees (4,5)		
Rose (3)		
Snapdragon (3,4)		
Sweet peas (3)		
Sycamore (3)		
Willow (4)		
Zinnia (2b,3)		

ORNAMENTAL PLANT DISEASE CONTROL (Cont'd)

APPLICATION RATES: Except as noted for specific diseases, mix 11 ounces of BAYLETON 25% Wettable Powder in 275 to 550 gallons of water and apply as a full coverage foliage spray to the point of drip as needed.

- [A] Mix 11 ozs of BAYLETON 25% Wettable Powder in 68.75 to 137.5 gallons of water and apply as a full-coverage foliar spray to the point of drip. Use multiple applications at 7- to 14-day intervals as needed dependent upon bloom periods. Applications should begin at the expanded bud stage (color showing or at bud break on Hemlock). Use 4 applications at 14-day intervals for Hemlock rust.
- [B] Mix 11 oz of BAYLETON 25% Wettable Powder plus sufficient spreader sticker for good coverage in 68.75 gallons of water. Apply in a spray application to the point of run-off on an as needed basis during the early part of the season. Excessive rates or excessive applications may result in a shortening of the flower stalk on iris.
- [C] Mix 11 oz of BAYLETON 25% Wettable Powder in 68.75 to 137.5 gallons of water and apply as a full coverage foliar spray to point of run-off. Apply in early spring as growth starts and re-apply on a 14- to 21 day interval until new growth is fully expanded. Protect new growth that develops in late summer or fall as temperatures begin to drop.
- [D] Greenhouse Applications of BAYLETON 25% Wettable Powder

Winter Use -- 1 oz in 50 gallons of water or 11 oz in 550 gallons of water

Summer Use -- 2 oz in 50 gallons of water or 11 oz in 275 gallons of water

Mix specified amount of BAYLETON 25% Wettable Powder in water and apply in a spray application to the point of drip. Intervals between applications should be no shorter than 30 days to avoid flower stalk length reduction. Excessive rates or applications may result in a shortening of the flower stalk.

- [E] For control of Exobasidium flower and leaf gall, apply 11 ounces of BAYLETON 25% Wettable Powder in 275 gallons of water. Begin application at bud break and apply at 10-day intervals through infestation period.

COMPATIBILITY: BAYLETON is compatible with many registered insecticides and fungicides. To determine the compatibility of BAYLETON with specific products, the following procedure should be conducted. Pour the recommended proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least 5 minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible.

SPRAY ADDITIVES: Use of various spray additives such as spreaders, extenders, trace elements or fertilizers should be evaluated prior to use. The label directions given here are based on data obtained with no additives; use of any product with BAYLETON may affect the result. Contact local university extension personnel prior to use of spray mix additives.

RESTRICTIONS: Edible portions of treated trees, such as nuts and syrup, should not be used for feed or food.

STORAGE AND DISPOSAL

- ✓ Do not contaminate water, food or feed by storage or disposal.
- ✓ Household: Do not reuse empty container. Securely wrap original container in several layers of newspaper and put in trash collection.
- ✓ Pesticide Disposal (Except Household): Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
- ✓ Container Disposal (Except Household): Completely empty carton into application equipment. Then dispose of empty carton in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.
- ✓ Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Do not store below freezing (32°F). Exposure to moisture or excessive handling of water soluble packets may cause breakage. Store packets in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Material that cannot be used as directed should be disposed of as directed above. In spill or leak incidents, keep unauthorized people away. You may contact the Mobay emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Mobay Kansas City Emergency Response Telephone No. is 816-242-2582, or contact Chemtrec at 800-424-9300.

BAYLETON is a Reg. TM of Bayer AG, Germany.

BAYLETON®

25% Wettable Powder

SYSTEMIC FUNGICIDE

ACTIVE INGREDIENT:

1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H
1,2,4-triazol-1-yl)-2-butanone

EPA File Symbol No. 3125-XXX

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

(See Outer Package for Directions and Precautions)

Net Contents 11 Ounces