

BAYLETON 25 T&O

413-8541.YLD

PM 22 2125-413

U. S. LABEL

Base Pre-Reg (8541)

10/2

Reason to Issue: To revise name for bermudagrass decline and take all patch.

Date of Draft: 06/07/95(Pre-Reg)(T)

BAYLETON® 25

Turf and Ornamental Fungicide

Systemic Fungicide in Water Soluble Packets

For control of certain diseases on flowers, foliage plants, shrubs, turf and shade trees. Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

ACTIVE INGREDIENT:

1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanone 25%


INERT INGREDIENTS 75%
100%

Keep water soluble packets in this container and store in a cool place but not below freezing (32°F).

U. S. Patent No. 3,912,752
EPA Reg. No. 3125-413
EPA Est. 3125-MO-1

SELL IN ORIGINAL PACKAGES ONLY
CONTAINS FOUR 11 OUNCE WATER SOLUBLE PACKETS
Total Net Contents: 2.75 Pounds

AMENDMENT TO
PREVIOUSLY REGISTERED LABELING



Unregistered label text is highlighted in redlining.
Text to be deleted is indicated by ~~strikeout~~.

Compared to 4/13/95 Accepted label. p

10/2

2 of 2

RECOMMENDED APPLICATIONS - TURF GRASS DISEASE CONTROL		
DISEASE	OZ OF BAYLETON 25 TURF AND ORNAMENTAL PER 1,000 SQ FEET*	REMARKS
<p><i>Gaeumannomyces</i> <i>-graminis var. graminis</i></p> <p>(Such as Bermuda grass decline, take-all patch on centipede grass)</p> <p><i>Bermudagrass decline</i> (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>)</p> <p><i>Take-all patch</i> (<i>Gaeumannomyces graminis</i> var. <i>avenae</i>)</p> <p>(Except California)</p>	<p>Preventive Curative</p> <p>2 4</p>	<p>Immediately after the fungicide is applied, the area should be thoroughly irrigated to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone. The objective is to water the fungicide into the crown and root zone.</p> <p>PREVENTIVE RATE: Begin applications 2 to 4 weeks prior to the appearance of disease symptoms. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information. Apply subsequent applications at 21- to 28-day intervals.</p> <p>CURATIVE RATE: To control existing infections, apply 4 oz at 14- to 21-day intervals for the initial 1 or 2 treatments followed by 2 oz at 21- to 28-day intervals. Cultural control practices such as aerification, topdressing, reseeding, and fertilization must be implemented prior to or at the same time the fungicide is applied. Refer to your local County Extension Service for this information.</p>

2