416-8308.YLD

Base Pre-Reg. (8308)

U.S. LABEL

Reason to Issue: To require irrigation after application to golf courses.

Date of Draft: 09/01/94(Pre-Reg)(S) Supersedes Pre-Reg Draft Dated: 04/20/94

MERIT 0.62 G

Insecticide

For Systemic Insect Control In Turfgrass And Landscape Ornamentals

ACTIVE INGREDIENT:

Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine INERT INGREDIENTS

0.62% 99.38% 100.00%

EPA Reg. No. 3125-416

NET CONTENTS: ____ Pounds

SEP 20 1994

A Reg. No.

STOP - Read The Label Before Use **KEEP OUT OF REACH OF CHILDREN**

CAUTION

PRECAUCION AL USUARIO: Si usted no puede leer o entender inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing.

Wash thoroughly with soap and water after

STATEMENTS OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger, or, if available, by administering syrup of ipecac. If syrup of ipecac is available, administer 1 tablespoonful (15 ml) of syrup of ipecac followed by 1 to 2 glasses of water. If vomiting does not occur within 20 minutes, repeat the dose once.

Do not induce vomiting or give anything by mouth to an unconscious person, if on skin: Wash thoroughly, with, soap and, water. Get medical attention is iditation occurs. If In eyes: Hold eyelids open and flush with plenty of water. Call a physician if irritation persists.

TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically.

ERIT

ECTICID

q 4

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read these entire DIRECTIONS FOR USE, GENERAL INFORMATION, AND CONDITIONS OF SALE before using MERIT 0.62 G Insecticide.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE APPROPRIATE FOR THE CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED UNDER DIFFERENT ENVIRONMENTAL CONDITIONS BOTH INDOORS AND OUTDOORS UNDER CONDITIONS SIMILAR TO THOSE THAT ARE ORDINARY AND CUSTOMARY WHERE THE PRODUCT IS TO BE USED.

INSUFFICIENT CONTROL OF PESTS OR PLANT INJURY MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL CONDITIONS, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO ANIMALS, MAN, AND DAMAGE TO THE ENVIRONMENT. MILES OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL ENVIRONMENTAL CONDITIONS, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF MILES AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

Do not formulate this product in other end-use products.

GENERAL INFORMATION

APPLICATION TO TURFGRASS

MERIT 0.62 G Insecticide can be used for the control of soil inhabiting pests of turfgrass, such as Northern & Southern masked chafers, Cyclocephala borealis, C. immaculata, and/or C. lurida; Asiatic garden beetle, Maladera castanea; European chafer, Rhizotrogus majalis; May or June beetles, Phyllophaga spp.; Japanese beetle, Popillia japonica; Oriental beetle, Anomala orientalis: Billbugs, Spherophorus spp.; Annual bluegrass weevil, Hyperodes spp.; Black turfgrass ataenius. .4taenius spretulus; and Aphodius spp. MERIT 0.62 G Insecticide can be used as directed on turfgrass in sites such as home lawns, business and office complexes, shopping complexes, multi-family residential complexes, golf courses, airports, cemeteries, parks, playgrounds, and athletic fields. MERIT Insecticide can not be used on comi roial sod farms.

The active ingredient in MERIT 0.62 G Insecticide has sufficient residual activity so that applications can be made preceding the egolisying activity of the target pests. High levels of control can be achieved when applications are made preceding or during the egg laying period. The need for an application can be based on historical monitoring of the site, provious records or experiences, current season adult tropping or other methods. Optimum control will be achieved when applications are made prior to egg hatch of the target pest, followed by sufficient irrigation of cainfall to move the active ingredient through the thatch.

Applications should not be made when turfgrass areas are waterlogged or the soil is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist. The treated turf must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile. Applications cannot exceed a total of 65 lb (0.4 lb of active ingredient) per acre per year.

APPLICATION TO ORNAMENTALS

MERIT 0.62 G Insecticide can be used as a broadcast application on ornamentals in commercial and usidential landscapes. MERIT 0.62 G is a systemic product and will be translocated upward within the plant system. Optimum control will be achieved when applications are followed by sufficient mechanical incorporation, irrigation or rainfall to move the active ingredient into the soil. Applications should be made prior to anticipated pest infestation to achieve optimum levels of control.

Applications of MERIT 0.62 G Insecticide cannot exceed a total of 65 lb (0.4 lb of active ingredient) per acre per year.

NOTE: Not for use in commercial greenhouses or nurseries or on sod farms.

BEST AVAILABLE COPY

APPLICATION EQUIPMENT

Apply MERIT 0.62 G Insecticide uniformly over the area being treated with normally used granular application equipment. Both drop-type and rotary-type spreaders may be used to apply MERIT 0.62 G Insecticide. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly and accurately.

SPREADER SETTINGS FOR MERIT 0.62 G INSECTICIDE

The following chart can be used as a guide in establishing an accurate rate of MERIT 0.62 G Insecticide. Application rates will vary dependent upon equipment wear and condition of spreader. Carefully calibrate all the equipment used prior to commercial use of this product.

(Chart to Come)

RECOMMENDED APPLICATIONS				
CROP	PEST	DOSAGE MERIT 0.62 G	REMARKS	
Turf Grasses	Larvae of: Japanese beetle Black turfgrass ataenius Northern masked chafer Southern masked chafer European chafer Oriental beetle Asiatic garden beetle Phyllophaga spp. Billbugs Annual bluegrass weevil	40 to 65 lb per acre or 0.9 to 1.5 lb per 1000 sq ft	For optimum control, make application prior to egg hatch of the target pest. Be sure to read "APPLICATION EQUIPMENT" Section of this label. SEE NOTE SEE GENERAL INFORMATION SECTION OF THIS LABEL	
	Larvae of: Japanese beetle Black turfgrass ataenius Northern masked chafer Southern masked chafer European chafer Oriental beetle Asiatic garden beetle Phyllophaga spp.	50 lb per acre or 1.2 lb per 1000 sq ft		
	Larvae of: Billbugs Annual bluegrass weevil	50 to 65 lb per acre or 1.2 to 1.5 lb per 1000 sq ft		

Consult your local State Agricultural Experiment Station, or State Extension Turf Specialists for more specific information regarding tirning of application.

NOTE: For optimum control, irrigation or rainfall should occur within 24 hours after application to move the active ingredient through the thatch. On golf courses, irrigate treated areas following application. Do not apply more than 65 lb (0.4 lb of active ingredient) per acre per year. Avoid mowing turf or lawn area until after irrigation or rainfall has occurred so that christenity of explication will not be affected.

RECOMMENDED APPLICATIONS FOR USE ON ORNAMENTALS For use only in and around industrial and commercial buildings and residential areas (including trees, shrubs, evergreens, flowers, foliage plants and ground covers).					
CROP	PEST	DOSAGE MERIT 0.62 G	REMARKS		
Landscape · Plantings	White grub lervee (such as Japanese beetle larvae, Chafers, Phyllophaga spp. Asiatic garden beetle, Oriental beetle)	1.2 to 1.5 lb per 1000 sq ft	Apply as a broadcast treatment. Prior to application, remove black plastic or other weed barrier material that will not permit water permeability. NOTE: For optimum control mechanical incorporation, irrigation or rainfall should occur within 24 hours after application to move the active ingredient into soil. Refer to label sections on application equipment and spreader settings. SEE GENERAL INFORMATION		
Flowers, Groundcovers	Aphids Lecebugs Leafhoppers Leafminers Mealybugs Scale Insects Thrips Whiteflies				

RESTRICTIONS

Avoid runoff or puddling of irrigation water following application. Avoid application of MERIT 0.62 G Insecticide to areas which are waterlogged or saturated, which will not allow penetration into the root zone of the plant.

Do not apply more than 65 lb (0.4 lb of active ingredient) per acre per year. Do not plant any food crop within one year of a treatment with MERIT 0.62 G Insecticide.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty container into application equipment. Dispose of empty container by incineration in a sanitary landfill, or if allowed by the state and local authorities, by burning. If burned, stay out of smoke.

Stold in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and

out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If material is spilled for any reason or cause, carefully contain any spilled material to prevent non-target contamination. Do not walk through spilled material and dispose of as directed for pesticides above. Refer to Precautionary Statements on label for hazards associated with the handling of this material. In spill or leak incidents, keep unauthorized people away. You may contact the Miles Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Miles Kansas City Emergency Response telephone number is 816-242-2582, or contact Chemtrec at 800-424-9300.

Miles Inc.: Specialty Products
Box 4913, Khnnas City, MO 64120-0013

