

419-8296.YLD

U.S. LABEL

Base Pre-Reg (8296)

Reason to Issue: To add groundwater advisory and
remove trade name BAYLETON.

Date of Draft: 08/23/94(Pre-Reg)(T)

SUMMIT S

Flowable Fungicide

FOR CONTROL OF SPECIFIED DISEASES ON CERTAIN FIELD AND FRUIT CROPS

ACTIVE INGREDIENT:

Triadimefon, 1-(4-Chlorophenoxy)-3,3-dimethyl-1-(2H-1,2,4-triazol-1-yl)-2-butanone ..	2.1%
Sulfur	53.7%
INERT INGREDIENTS	44.2%
	100%

Contains 0.25 lb of triadimefon and 6.4 lb of sulfur per gallon.

EPA Reg. No. 3125-419

Net Contents: ___ Gallon(s)


AMENDMENT TO CURRENTLY REGISTERED LABELING REVISE AS FOLLOWS

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposal of equipment washwaters. Apply this product only as specified on this label.

Do not make applications when weather conditions favor drift from target area.

GROUND WATER ADVISORY: This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.


MILES
BEST AVAILABLE COPY

ay

RECOMMENDED APPLICATIONS			
CROP	DISEASE	RATE OF SUMMIT S Flowable Fungicide	REMARKS (Also refer to Directions for Use)
Grape	Powdery mildew	1 to 2 quarts per acre	<p>Apply recommended dosage as a preventive spray in a minimum of 50 gallons of solution per acre by ground equipment. Increase spray volume as vine growth increases to ensure thorough coverage of the foliage and fruit.</p> <p>Make the first application prebloom to the new cane growth. Continue applications at 10- to 14-day intervals to maintain disease control.</p> <p>Use the higher label rate under heavy disease pressure, with 14-day spray intervals, and/or on varieties which are highly disease susceptible.</p> <p>Do not apply to grape varieties which are sensitive to sulfur.</p> <p>SUMMIT S may be applied up to 14 days before harvest. A maximum of 9 quarts of product may be applied per acre per crop season. If other products containing BAYLETON triadimefon are used during the season, do not exceed a total of 9 ounces active of triadimefon per acre per season.</p>
Sugar beet	Powdery mildew	1 to 4 quarts per acre	<p>Apply recommended dosage as a broadcast spray at first sign of disease symptoms. Use the higher label rate under heavy disease pressure. Additional applications may be made at 10- to 14-day intervals to maintain disease control or if reinfection occurs.</p> <p>SUMMIT S may be applied up to 15 days before harvest. A maximum of 8 quarts of product may be applied per acre per crop season. If other products containing BAYLETON triadimefon are used during the season, do not exceed a total of 8 ounces active of triadimefon per acre per season.</p>

RESTRICTIONS

If **BAYLETON** or other products containing triadimefon are used during the season, the total amount of triadimefon should not exceed the maximum season total stated on the

BAYLETON label for each crop. Each quart of SUMMIT S contains 1 ounce of triadimefon active ingredient.

Do not use on other crops grown for food or forage.

Miles Inc.
Crop Protection Products
Box 4913, Kansas City, MO 64120

BEST AVAILABLE COPY